

Matroid Polytopes: Algorithms, Theory, and Applications

By

DAVID CARLISLE HAWS

B.S. (University of California, Davis) 2004

DISSERTATION

Submitted in partial satisfaction of the requirements for the degree of

DOCTOR OF PHILOSOPHY

in

Mathematics

in the

OFFICE OF GRADUATE STUDIES

of the

UNIVERSITY OF CALIFORNIA

DAVIS

Approved:

Jesús A. De Loera (*chair*)

Matthias Köppe

Eric Babson

Committee in Charge

2009

Contents

Abstract	iii
Acknowledgments	iv
Chapter 1. What are Matroids and who are their polytopes?	1
1.1. Matroids	1
1.2. Matroid Polytopes & Polymatroids	7
Chapter 2. Volume and Ehrhart Polynomial Computation	17
2.1. Computing the Ehrhart Polynomials	18
2.2. Preliminaries on Rational Generating Functions.	18
2.3. On the Tangent Cones of Matroid Polytopes	20
2.4. Polymatroids	30
2.5. The construction of a short multivariate rational generating function	32
2.6. Polynomial-time specialization of rational generating functions in varying dimension	35
Chapter 3. Results on the Algebraic Combinatorics of Matroid Polytopes	43
3.1. Algebraic Properties of h^* -vectors and Ehrhart polynomials of Matroid Polytopes	43
3.2. Unimodular Simplices of Matroid Polytopes	53
3.3. Two Dimensional Faces of Matroid Polytopes	64
Chapter 4. Applications to Optimization Through the Structure of Matroid Polytopes	67
4.1. Description of the Algorithms and Heuristics	67
4.2. Software Implementation	79
4.3. Computational Results	80
Bibliography	126

Abstract

This dissertation presents new results on three different themes all related to matroid polytopes. First we investigate properties of Ehrhart polynomials of matroid polytopes, independence matroid polytopes, and polymatroids. We prove that for fixed rank their Ehrhart polynomials are computable in polynomial time. The proof relies on the geometry of these polytopes as well as a new refined analysis of the evaluation of Todd polynomials.

Second, we discuss theoretical results regarding the algebraic combinatorics of matroid polytopes. We discuss two conjectures about the h^* -vector and coefficients of Ehrhart polynomials of matroid polytopes and provide theoretical and computational evidence for their validity. We also explore a variant of White's conjecture which states that every matroid polytope has a regular unimodular triangulation. We provide extensive computational evidence supporting this new conjecture and propose a combinatorial condition on simplices sufficient for unimodularity. Lastly we discuss properties of two dimensional faces of matroid polytopes.

Finally, motivated by recent work on algorithmic theory for non-linear and multicriteria matroid optimization, we have developed algorithms and heuristics aimed at practical solutions of large instances of these difficult problems. Our methods primarily use the local adjacency structure inherent in matroid polytopes to pivot to feasible solutions which may or may not be optimal. We also present a modified breadth-first-search heuristic that uses adjacency to enumerate a subset of feasible solutions. We present other heuristics, and provide computational evidence supporting these new techniques. We implemented all of our algorithms in the software package MOCHA (Matroids Optimization Combinatorial Heuristics and Algorithms).

Acknowledgments

First and foremost to my advisor Jesús De Loera who started me on this journey as an undergraduate. He guided me around the pitfalls, encouraged me, mentored me, provided me with problems and kept challenging me at every turn. His patience was remarkable and he was my number one cheerleader. Most of all, he taught me that mathematics without passion is not math at all.

To my wife Tami Joy whose support and love made this all possible. Her utmost confidence in me is amazing. It lifted my spirits and kept me going. She can always make me laugh, and is always there to lean on.

My thanks to Matthias Köppe for sharing his mathematical and computational skills. His careful and thoughtful approach to mathematics set a great example and he always made himself available for any math or computer questions.

To my family and friends for all their support, understanding and love. My thanks to the UC Davis mathematics department staff who were very helpful and instrumental in my success.

Thanks to Francisco Santos for all our insightful conversations on triangulations of matroid polytopes and more.

My appreciation to Eric Babson for serving on my qualification exam and dissertation committee.

Support for this dissertation was provided by VIGRE (DMS-01-35345, DMS-0636297), University of California at Davis (DMS-0608785) and a gift from IBM.

CHAPTER 1

What are Matroids and who are their polytopes?

1.1. Matroids

Matroids are combinatorial objects that naturally encapsulate the idea of independence. They are found in matrices, graphs, transversals, point configurations, hyperplane arrangements, greedy optimization, and pseudosphere arrangements to name a few. One of the reasons matroids have become fundamental objects in pure and applied combinatorics are their many equivalent axiomatizations. For an excellent guide and thorough treatment of matroids we recommend [Oxl92, Wel76, Sch03], which we refer to for most all of our definitions and theory. We now offer a brief introduction to matroids and their associated polytopes. We begin with the most basic and prominent matroid definition (axiomatization). First we adopt the notation $X + y := X \cup \{y\}$ and $X - y := X \setminus \{y\}$. A pair $(S, \mathcal{I})$ is called a *matroid* if S is a finite set and $\mathcal{I}$ is a nonempty collection of subsets of S (denoted 2^S) satisfying:

$$(1.1) \quad \begin{aligned} (i) \quad & \text{if } I \in \mathcal{I} \text{ and } J \subseteq I, \text{ then } J \in \mathcal{I} \\ (ii) \quad & \text{if } I, J \in \mathcal{I} \text{ and } |I| < |J|, \text{ then } I + z \in \mathcal{I} \text{ for some } z \in J \setminus I. \end{aligned}$$

For a matroid $\mathcal{M} = (S, \mathcal{I})$, $I \subseteq S$ is *independent* if $I \in \mathcal{I}$ and *dependent* otherwise. $B \subseteq S$ is called a *base* or *basis* if it is an inclusion maximal independent set. That is, $B \in \mathcal{I}$ and there does not exist $A \in \mathcal{I}$ such that $B \subset A \subseteq S$. We denote the collection of bases of $\mathcal{M}$ as $\mathcal{B}_{\mathcal{M}}$, or sometimes $\mathcal{B}$ when the matroid is clear. An easy consequence of Equation 1.1 is that all bases have the same cardinality: If not, let B_1, B_2 be two bases of $(S, \mathcal{I})$ such that $|B_1| < |B_2|$. Then by (ii) in Equation 1.1 there exists $z \in B_2$ such that $B_1 + z \in \mathcal{I}$. But this contradicts the inclusionwise maximality of B_1 .

Given any $U \subseteq S$ the *rank* of U is given by the *rank function* of $\mathcal{M}$, $\varphi: 2^{[n]} \rightarrow \mathbb{Z}$ where

$$\varphi(A) := \max\{|X| \mid X \subseteq A, X \in \mathcal{I}\}.$$

The common cardinality of the bases of $\mathcal{M}$ is called the *rank* of the matroid. Throughout we will refer to the rank of $\mathcal{M}$ as $\text{rank}(\mathcal{M}) := \text{rank}(S)$.

EXAMPLE 1 (Uniform matroid). Let $[n] := \{1, \dots, n\}$. The *uniform* matroid of rank r is $U^{r,n} := \{A \subseteq [n] \mid |A| \leq r\}$. It is easy to see that $U^{r,n}$ satisfies both (i) and (ii) of [Equation 1.1](#).

$$U^{2,4} = \{\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}\}$$

We state an additional useful axiomatization on the independence sets.

LEMMA 2 (Theorem 39.1 [\[Sch03\]](#)). *Let S be a finite set and let $\mathcal{I}$ be a nonempty collection of subsets satisfying [Equation 1.1\(i\)](#). Then [Equation 1.1\(ii\)](#) is equivalent to:*

$$(1.2) \quad \text{if } I, J \in \mathcal{I} \text{ and } |I \setminus J| = 1, |J \setminus I| = 2, \text{ then } I + z \in \mathcal{I} \text{ for some } z \in J \setminus I.$$

PROOF. Obviously [Equation 1.1\(ii\)](#) implies [Equation 1.2](#). Conversely, [Equation 1.1\(ii\)](#) follows from [Equation 1.2](#) by induction on $|I \setminus J|$, the case $|I \setminus J| = 0$ being trivial. If $|I \setminus J| \geq 1$, choose $i \in I \setminus J$. We apply the induction hypothesis twice: first to $I - i$ and J to find $j \in J \setminus I$ with $I - i + j \in \mathcal{I}$, and then to $I - i + j$ and J to find $j' \in J \setminus (I + j)$ with $I - i + j + j' \in \mathcal{I}$. Then by [Equation 1.2](#) applied to I and $I - i + j + j'$, we have $I + j \in \mathcal{I}$ or $I + j' \in \mathcal{I}$. $\square$

We can now prove one of the most useful axiomatizations for the purposes of this dissertation. Considering the properties of [Equation 1.1](#) it is natural to think that the bases of $\mathcal{M}$ are all that are needed to fully describe a matroid and indeed the following theorem gives an important characterization of matroids in terms of bases.

THEOREM 3 (Theorem 39.6 [\[Sch03\]](#)). *Let S be a set and $\mathcal{B}$ be a nonempty collection of subsets of S . Then the following are equivalent:*

- (i) $\mathcal{B}$ is the collection of bases of a matroid;
- (1.3) (ii) if $B, B' \in \mathcal{B}$ and $x \in B' \setminus B$, then $B' - x + y \in \mathcal{B}$ for some $y \in B \setminus B'$;
- (iii) if $B, B' \in \mathcal{B}$ and $x \in B' \setminus B$, then $B' - y + x \in \mathcal{B}$ for some $y \in B \setminus B'$.

PROOF. (i) $\Rightarrow$ (ii): Let $\mathcal{B}$ be the collection of bases of a matroid $(S, \mathcal{I})$. Then all sets in $\mathcal{B}$ have the same size. Now, let $B, B' \in \mathcal{B}$ and $x \in B' \setminus B$. Since $B' - x \in \mathcal{I}$, there exists

a $y \in B \setminus B'$ with $B'' := B' - x + y \in \mathcal{I}$. Since $|B''| = |B'|$, we know $B'' \in \mathcal{B}$.

(iii) $\Rightarrow$ (i): (iii) directly implies that no set in $\mathcal{B}$ is contained in another. Let $\mathcal{I}$ be the collection of sets I with $I \subseteq B$ for some $B \in \mathcal{B}$. We check Equation 1.2. Let $I, J \in \mathcal{I}$ with $|I \setminus J| = 1$ and $|J \setminus I| = 2$. Let $I \setminus J = \{x\}$.

Consider sets $B, B' \in \mathcal{B}$ with $I \subseteq B, J \subseteq B'$. If $x \in B'$, we are done. So assume $x \notin B'$. Then by (iii), $B' - y + x \in \mathcal{B}$ for some $y \in B' \setminus B$. As $|J \setminus I| = 2$, there is a $z \in J \setminus I$ with $z \neq y$. Then $I + z \subseteq B' - y + x$, and so $I + z \in \mathcal{I}$.

(ii) $\Rightarrow$ (iii): By the foregoing we know that (iii) implies (ii). Now axioms (ii) and (iii) interchange if we replace $\mathcal{B}$ by the collection of complements of sets in $\mathcal{B}$. Hence the implication (ii) $\Rightarrow$ (iii) also holds. $\square$

Parts (ii) and (iii) of Equation 1.3 might seem identical, but (ii) says that we can pick an element to *remove* from B' and there exist some element of B to add to get a basis while (iii) says that we can pick an element to *add* to B' and there exists some element of B to remove to get a basis. Having the flexibility to choose the element to add or remove is a useful property for many proofs, algorithms and heuristics on the bases of matroids.

EXAMPLE 4 (Graphical Matroid). Let $G = (V, E)$ be graph with nodes V and edges E . The collection of spanning forests of G are the bases of the *graphical* matroid on ground set V , denoted $\mathcal{M}_G$. We can verify the basis exchange axiom Equation 1.3(iii) by considering two spanning forests F_1, F_2 . If $e \in F_2 \setminus F_1$, then $F_1 + e$ contains a cycle C in G . Also there exists $f \in C \setminus F_2$ and thus $F_1 - f + e$ is a spanning forest.

Matroids play a prominent role in combinatorial optimization and are intrinsically tied to the greedy algorithm which we state now: If $w : S \rightarrow \mathbb{R}$ is a weight function we use the notation $w(I) := \sum_{i \in I} w(i)$ where $I \subseteq S$. Letting $\mathcal{I} \subset 2^S$ be closed under taking subsets and given a weighting w , the *greedy algorithm* starts by setting $I := \emptyset$, and repeatedly choosing $y \in S \setminus I$ such that $I + y \in \mathcal{I}$ where $w(y)$ is as large as possible. We terminate if no such y exists.




FIGURE 1.1. (a) A graph G and (b) its spanning forests, i.e. bases $\mathcal{B}_{\mathcal{M}_G}$ of $\mathcal{M}_G$.

For arbitrary $\mathcal{I} \subseteq 2^S$ the greedy algorithm does not always terminate at the maximum solution, although for matroids this will always be the case. In fact, the converse is true: If the greedy algorithm always works on $\mathcal{I}$ then it is a matroid.

THEOREM 5 (Theorem 40.1 [Sch03]). *Let $\mathcal{I}$ be a nonempty collection of subsets of a set S , closed under taking subsets. Then the pair $(S, \mathcal{I})$ is a matroid if and only if for each weight function $w : S \rightarrow \mathbb{R}_+$, the greedy algorithm leads to a set $I \in \mathcal{I}$ of maximum weight $w(I)$.*

PROOF. Necessity. Let $(S, \mathcal{I})$ be a matroid and let $w : S \rightarrow \mathbb{R}_+$ be any weight function on S . Call an independent set I good if it is contained in a maximum-weight basis. It suffices to show that if I is good, and y is an element in $S \setminus I$ with $I + y \in \mathcal{I}$ and with $w(y)$ as large as possible, then $I + y$ is good.

As I is good, there exists a maximum-weight basis $B \supseteq I$. If $y \in B$, then $I + y$ is good again. If $y \notin B$, then there exists a base B' containing $I + y$ and contained in $B + y$. So $B' = B - z + y$ for some $z \in B \setminus I$. As $w(y)$ is chosen maximum and as $I + z \in \mathcal{I}$ since $I + z \subseteq B$, we know $w(y) \geq w(z)$. Hence $w(B') \geq w(B)$, and therefore B' is a maximum-weight base. So $I + y$ is good.

Sufficiency. Suppose that the greedy algorithm leads to an independent set of maximum weight for each weight function $w : S \rightarrow \mathbb{R}_+$. We show that $(S, \mathcal{I})$ is a matroid.

Condition Equation 1.1(i) is satisfied by assumption. To see condition Equation 1.1(ii), let $I, J \in \mathcal{I}$ with $|I| < |J|$. Suppose that $I + z \notin \mathcal{I}$ for each $z \in J \setminus I$. Let $k := |I|$. Consider the following weight function w on S :

$$w(S) := \begin{cases} k+2 & \text{if } s \in I, \\ k+1 & \text{if } s \in J \setminus I, \\ 0 & \text{if } s \in S \setminus (I \cup J). \end{cases}$$

Now in the first iterations of the greedy algorithm we find the k elements in I . By assumption, at any further iteration, we cannot choose any element in $J \setminus I$. Hence any further element chosen, has weight 0. So the greedy algorithm yields an independent set of weight $k(k+2)$.

However, J has weight at least $|J|(k+1) \geq (k+1)(k+1) > k(k+2)$. Hence the greedy algorithm does not give a maximum-weight independent set, contradicting our assumption. $\square$

The rank function φ of a matroid $\mathcal{M}$ also determines $\mathcal{M}$ and we can easily see that a set $A \subseteq S$ is independent if and only if $\varphi(A) = |A|$.

THEOREM 6 (Theorem 39.8 [Sch03]). *Let S be a set and $\varphi : 2^S \rightarrow \mathbb{Z}_+$. Then φ is the rank function of a matroid if and only if for all $T, U \subseteq S$:*

$$(1.4) \quad \begin{aligned} (i) \quad & \varphi(T) \leq \varphi(U) \leq |U| \text{ if } T \subseteq U, \\ (ii) \quad & \varphi(T \cap U) + \varphi(T \cup U) \leq \varphi(T) + \varphi(U). \end{aligned}$$

PROOF. Necessity. Let φ the rank function of a matroid $(S, \mathcal{I})$. Choose $T, U \subseteq S$. Clearly Equation 1.4(i) holds. To see (ii), let I be an inclusionwise maximal set in $\mathcal{I}$ with $I \subseteq T \cap U$ and let J be an inclusionwise maximal set in $\mathcal{I}$ with $I \subseteq J \subseteq T \cup U$. Since $(S, \mathcal{I})$ is a matroid, we know that $\varphi(T \cap U) = |I|$ and $\varphi(T \cup U) = |J|$. Then

$$\varphi(T) + \varphi(U) \geq |J \cap T| + |J \cap U| = |J \cap (T \cap U)| + |J \cap (T \cup U)| \geq |I| + |J| = \varphi(T \cap U) + \varphi(T \cup U);$$

that is, we have Equation 1.4(ii).

Sufficiency Let $\mathcal{I}$ be the collection of subsets I of S with $\varphi(I) = |I|$. We show that $(S, \mathcal{I})$ is a matroid, with rank function φ .

Trivially, $\emptyset \in \mathcal{I}$. Moreover, if $I \in \mathcal{I}$ and $J \subset I$, then

$$\varphi(J) \leq \varphi(I) - \varphi(I \setminus J) \geq |I| - |I \setminus J| = |J|.$$

So $J \in \mathcal{I}$.

In order to check [Equation 1.2](#), let $I, J \in \mathcal{I}$ with $|I \setminus J| = 1$ and $|J \setminus I| = 2$. Let $J \setminus I = \{z_1, z_2\}$. If $I + z_1, I + z_2 \notin \mathcal{I}$, we have $\varphi(I + z_1) = \varphi(I + z_2) = |I|$. Then by [Equation 1.4\(ii\)](#),

$$\varphi(J) \leq \varphi(I + z_1 + z_2) \leq \varphi(I + z_1) + \varphi(I + z_2) - \varphi(I) = |I| < |J|,$$

contradicting the fact that $J \in \mathcal{I}$.

So $(S, \mathcal{I})$ is a matroid. Its rank function is φ , since $\varphi(U) = \max\{|I| \mid I \subseteq U, I \in \mathcal{I}\}$ for each $U \in S$. Here $\geq$ follows from [Equation 1.4\(ii\)](#), since if $I \subseteq U$ and $I \in \mathcal{I}$, then $\varphi(U) \geq \varphi(I) = |I|$. Equality can be shown by induction on $|U|$, the case $U = \emptyset$ being trivial. If $U \neq \emptyset$, choose $y \in U$. By induction, there is an $I \subseteq U - y$ with $I \in \mathcal{I}$ and $|I| = \varphi(U - y)$. If $\varphi(U) = \varphi(U - y)$ we are done, so assume $\varphi(U) > \varphi(U - y)$. Then $I + y \in \mathcal{I}$, since $\varphi(I + y) \geq \varphi(I) + \varphi(U) - \varphi(U - y) \geq |I| + 1$. Moreover, $\varphi(U) \leq \varphi(U - y) + \varphi(\{y\}) \leq |I| + 1$. This proves equality for U . $\square$

One of the most natural methods of presenting a matroid in a computational model is by *independence oracle* which given any $A \subseteq S$ asserts whether $A \in \mathcal{I}$ or not. The independence oracle allows us to compute the rank of any $A \subseteq S$. Begin with $I := \emptyset$ and for each $i \in A$ (any order) if $I + i \in \mathcal{I}$ then set $I := I + i$. Repeat and thus, $\text{rank}(A) := |I|$. For an interesting overview of the computational relation of various matroid axiomatization representations see [\[May08\]](#).

In [Theorem 5](#) it was assumed that the weights are non-negative. But even with a weight function $w : S \rightarrow \mathbb{R}$ it can still be shown that for a matroid $\mathcal{M}$ the greedy algorithm finds a maximum-weight basis. Not surprising, one can replace the condition “as large as possible” with “as small as possible” and the greedy algorithm will find a minimum-weight basis. Assuming an independence oracle and [Theorem 5](#) the corollary follows:

COROLLARY 7 (Corollary 40.1a [\[Sch03\]](#)). *A maximum-weight independent set in a matroid can be found in strongly polynomial time.*

EXAMPLE 8 (Linear, Realizable, or Vector Matroids). Let $A := \{a_1 \mid a_2 \mid \cdots \mid a_n\} \in \mathbb{R}^{m \times n}$ be an $n \times m$ matrix. Let $S := \{1, \dots, n\}$ and we say $I \subseteq S$ is independent if the

columns of A indexed by I are linearly independent.

$$A = \begin{array}{c|ccccc} & \mathbf{1} & \mathbf{2} & \mathbf{3} & \mathbf{4} & \mathbf{5} \\ \hline & 1 & 0 & 1 & -1 & 2 \\ & 1 & 1 & 0 & 1 & 2 \end{array} \quad \{\mathbf{1}, \mathbf{2}\}, \{\mathbf{2}, \mathbf{3}\} \text{ are independent while } \{\mathbf{1}, \mathbf{5}\} \text{ is not.}$$

Such a matroid $\mathcal{M}_A$ is called *linear* or *realizable*. If the entries of A are in a field $\mathbb{F}$ then we say $\mathcal{M}_A$ is realizable over $\mathbb{F}$. Graphical matroids are linear and realizable over any field. This can be done by arbitrarily orienting the edges of G and take its incidence matrix. Not all matroids are linear, e.g. see F_8 in the appendix of [Oxl92]. Some matroids can only be realized in certain fields. For example, the Fano matroid can only be realized in a field of characteristic two [Oxl92].

1.2. Matroid Polytopes & Polymatroids

Let $A \subset [n] := \{1, \dots, n\}$ and define the *incidence vector* of A as

$$\mathbf{e}_A := \sum_{i \in A} \mathbf{e}_i$$

where $\mathbf{e}_i \in \mathbb{R}^n$ are the standard unit vectors. For example if $A = \{1, 3, 4\} \subseteq [5]$ then $\mathbf{e}_A = (1, 0, 1, 1, 0)^\top \in \mathbb{R}^5$. We also adopt the notation $\text{Inc}(X) := \{\mathbf{e}_A \mid A \in X\}$ where $X \subseteq 2^S$. Now we introduce the main object of this thesis.

DEFINITION 9. Let $\mathcal{M}$ be a matroid with bases $\mathcal{B}_\mathcal{M}$. The *matroid polytope* of $\mathcal{M}$ is

$$\mathcal{P}_\mathcal{M} := \text{conv}(\mathbf{e}_B \mid B \in \mathcal{B}_\mathcal{M})$$


where $\text{conv}(\cdot)$ denotes the convex hull.

EXAMPLE 10. Recall $U^{2,4} = \{\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}\}$. Thus

$$\mathcal{P}_{U^{2,4}} = \text{conv}((1, 1, 0, 0)^\top, (1, 0, 1, 0)^\top, (1, 0, 0, 1)^\top, (0, 1, 1, 0)^\top, (0, 1, 0, 1)^\top, (0, 0, 1, 1)^\top).$$

The polytope $\mathcal{P}_{U^{2,4}} \in \mathbb{R}^4$ but we can easily see that all vertices lay in a hyperplane since they all sum to two. $\mathcal{P}_{U^{2,4}}$ is in fact isomorphic to the cross polytope shown in [Figure 1.2](#).

The matroid polytope is different than the following well-known polytope.

FIGURE 1.2. The cross polytope $\mathcal{P}_{U^{2,4}}$.

DEFINITION 11. Let $\mathcal{M} = (S, \mathcal{I})$ be a matroid. The *independence matroid polytope* of $\mathcal{M}$ is

$$\mathcal{P}_{\mathcal{M}}^{\mathcal{I}} := \text{conv}(\mathbf{e}_I \mid I \in \mathcal{I}).$$

We can see that $\mathcal{P}(\mathcal{M}) \subseteq \mathcal{P}^{\mathcal{I}}(\mathcal{M})$ and $\mathcal{P}_{\mathcal{M}}$ is a face of $\mathcal{P}_{\mathcal{M}}^{\mathcal{I}}$ in the hyperplane $\sum_{i=1}^n x_i = \text{rank}(\mathcal{M})$.

Polymatroids are closely related to matroid polytopes and independence matroid polytopes. A function $\psi: 2^{[n]} \rightarrow \mathbb{R}$ is *submodular* if $\psi(X \cap Y) + \psi(X \cup Y) \leq \psi(X) + \psi(Y)$ for all $X, Y \subseteq [n]$ and *non-decreasing* if $\psi(X) \leq \psi(Y)$ for all $X \subseteq Y \subseteq [n]$. We say ψ is a *polymatroid rank function* if it is submodular, non-decreasing, and $\psi(\emptyset) = 0$. For example, the rank function of a matroid is a polymatroid rank function. The *polymatroid* determined by a polymatroid rank function ψ is the convex polyhedron in $\mathbb{R}^n$ given by

$$\mathcal{P}_{\psi} := \left\{ \mathbf{x} \in \mathbb{R}^n \mid \sum_{i \in A} x_i \leq \psi(A), \forall A \subseteq [n], \mathbf{x} \geq \mathbf{0} \right\}.$$

Independence matroid polytopes are a special class of polymatroids. Indeed, if φ is a rank function on some matroid $\mathcal{M}$, then $\mathcal{P}_{\mathcal{M}}^{\mathcal{I}} = \mathcal{P}_{\varphi}$ [Edm03]. Moreover, the matroid polytope $\mathcal{P}_{\mathcal{M}}$ is the face of $\mathcal{P}_{\varphi}$ lying in the hyperplane $\sum_{i=1}^n x_i = \varphi([n])$. Matroid polytopes and polymatroids appear in combinatorial optimization [Sch03], algebraic combinatorics [FS05], and algebraic geometry [GGMS87]. Edmonds showed that a linear function can be optimized over a polymatroid $\mathcal{P}_{\psi}$ via an extension of the greedy algorithm [Edm03]. This also provides a technique which easily lists off the vertices of a polymatroid $\mathcal{P}_{\psi}$. One




FIGURE 1.3. An example of a polymatroid

important implication we will see is that if ψ is an integral polymatroid rank function then $\mathcal{P}_\psi$ will be integral, that is, all the vertices of $\mathcal{P}_\psi$ are integral.

First we recall some basic definitions and key theorems from linear programming [Sch86a]. If S is a finite subset then $\mathbb{R}^S$ is the vector space of vectors with coordinates indexed by S . That is, if $\mathbf{x} \in \mathbb{R}^S$ and $s \in S$ then the s th coordinate of $\mathbf{x}$ is given by $x(s) \in \mathbb{R}$. If $A \subseteq S$ then $\mathbf{x}(A) := \sum_{a \in A} x(a)$.

Let S, T be finite sets, $\mathbf{c} \in \mathbb{R}^S$, $\mathbf{b} \in \mathbb{R}^T$ and $A = \{a_{ij} \in \mathbb{R} \mid i \in T, j \in S\}$ be a matrix. A *linear program* is

$$\max\{\mathbf{c} \cdot \mathbf{x} \mid A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq 0, \mathbf{x} \in \mathbb{R}^S\}.$$

The *dual linear program* is

$$\min\{\mathbf{b} \cdot \mathbf{y} \mid A^\top \mathbf{y} \leq \mathbf{c}, \mathbf{y} \geq 0, \mathbf{y} \in \mathbb{R}^T\}.$$

A vector $\mathbf{x} \in \mathbb{R}_+^S$ satisfying $A\mathbf{x} \leq \mathbf{b}$ is a *feasible solution* and a vector $\mathbf{y} \in \mathbb{R}_+^T$ satisfying $A^\top \mathbf{y} \leq \mathbf{c}$ is a *feasible dual solution*.

THEOREM 12 ([Sch03]). *For any linear program maximization problem exactly one of the following is true;*

- (i) *There exists no feasible solution,*
- (ii) *For any $\alpha \in \mathbb{R}$ there is a feasible solution $\mathbf{x}$ such that $\mathbf{c} \cdot \mathbf{x} > \alpha$,*
- (iii) *There is an optimal (feasible) solution.*

A linear program and its dual are related and we state the famous *strong duality theorem*.

THEOREM 13 (Corollary 7.1g [Sch86a]). *Let S, T be finite sets, $A \in \mathbb{R}^S \times \mathbb{R}^T$, $\mathbf{c} \in \mathbb{R}^S$, $\mathbf{b} \in \mathbb{R}^T$. Then*

$$\max\{\mathbf{c} \cdot \mathbf{x} \mid A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq 0, \mathbf{x} \in \mathbb{R}^S\} = \min\{\mathbf{b} \cdot \mathbf{y} \mid A^\top \mathbf{y} \leq \mathbf{c}, \mathbf{y} \geq 0, \mathbf{y} \in \mathbb{R}^T\}$$

if both sets are nonempty.

In order to justify the following approach to finding the vertices of a polymatroid we will use a fact from linear programming and polyhedral theory; every vertex of a polyhedra is the optimal solution to some linear program. Let φ be a polymatroid rank function on the finite set S and $\mathbf{w} : S \rightarrow \mathbb{R}$. We start by ordering the elements of S as $s_1, \dots, s_n$ such that $w(s_1) \geq w(s_2) \geq \dots \geq w(s_n)$ and define $U_i := \{s_1, \dots, s_i\}$ for $i = 0, \dots, n$. Next, define $\hat{\mathbf{x}} \in \mathbb{R}^S$ as

$$(1.5) \quad \hat{x}(s_i) := \varphi(U_i) - \varphi(U_{i-1}) \text{ for } i = 1, \dots, n.$$

The following theorem will show that $\hat{\mathbf{x}}$ maximizes $\mathbf{w} \cdot \mathbf{x}$ over $\mathcal{P}_\varphi$. Consider the linear program and its dual:

$$(1.6) \quad \begin{aligned} & \max\{\mathbf{w} \cdot \mathbf{x} \mid \mathbf{x} \in \mathcal{P}_\varphi\} \\ & = \min \left\{ \sum_{T \subseteq S} \mathbf{y}(T) \varphi(T) \mid \mathbf{y} \in \mathbb{R}_+^{2^S}, \sum_{T \subseteq S} \mathbf{y}(T) \cdot e_T \geq \mathbf{w} \right\} \end{aligned}$$

Also, define:

$$(1.7) \quad \begin{aligned} \hat{\mathbf{y}}(U_i) &= w(s_i) - w(s_{i+1}) & (i = 1, \dots, n-1) \\ \hat{\mathbf{y}}(S) &= w(s_n), \\ \hat{\mathbf{y}}(T) &= 0 & (T \neq U, \text{ for each } i). \end{aligned}$$

THEOREM 14 (Theorem 44.3 [Sch03]). *Let φ be a polymatroid rank function on S and let $\mathbf{w} : S \rightarrow \mathbb{R}^S$. Then $\hat{\mathbf{x}}$ and $\hat{\mathbf{y}}$ are optimum solutions to [Equation 1.6](#).*

PROOF. We first show that $\hat{\mathbf{x}} \in \mathcal{P}_\varphi$; that is, $\hat{\mathbf{x}} \leq \varphi(T)$ for each $T \subseteq S$. This is shown by induction on $|T|$, the case $T = \emptyset$ being trivial. Let $T \neq \emptyset$ and let k be the largest index with $s_k \in T$. Then by induction,

$$\hat{\mathbf{x}}(T - s_k) \leq \varphi(T - s_k).$$

Hence

$$\widehat{\mathbf{x}}(T) \leq \varphi(T - s_k) + \widehat{\mathbf{x}}(x_k) = \varphi(T - s_k) + \varphi(U_k) - \varphi(U_{k-1}) \leq \varphi(T)$$

(the last inequality follows from the submodularity of φ). So $\widehat{\mathbf{x}} \in \mathcal{P}_\varphi$.

Also, $\widehat{\mathbf{y}}$ is feasible for [Equation 1.6](#). Trivially, $\widehat{\mathbf{y}} \geq \mathbf{0}$. Moreover, for any i we have by [Equation 1.7](#):

$$\sum_{T \ni s_i} \widehat{\mathbf{y}}(T) = \sum_{j \geq i} \widehat{\mathbf{y}}(U_j) = \mathbf{w}(s_i).$$

So $\widehat{\mathbf{y}}$ is a feasible solution of [Equation 1.6](#).

Optimality of $\widehat{\mathbf{x}}$ and $\widehat{\mathbf{y}}$ follows from:

$$\begin{aligned} \mathbf{w} \cdot \widehat{\mathbf{x}} &= \sum_{s \in S} w(s) \widehat{x}(s) = \sum_{i=1}^n w(s_i) (\varphi(U_i) - \varphi(U_{i-1})) \\ &= \sum_{i=1}^{n-1} \varphi(U_i) (w(s_i) - w(s_{i+1})) + \varphi(S) w(s_n) = \sum_{T \subseteq S} \widehat{\mathbf{y}}(T) \varphi(T). \end{aligned}$$

The third equality follows from a straightforward reordering of the terms, using that $\varphi(\emptyset) = 0$. □

Thus, [Equation 1.5](#) gives a vertex of $\mathcal{P}_\varphi$. If we vary the weights we will get all the vertices of $\mathcal{P}_\varphi$ and it suffices to choose weights that permute the elements of S . [Equation 1.5](#) and [Theorem 14](#) imply the following important corollary.

COROLLARY 15 (Corollary 44.3d [[Sch03](#)]). *Let φ be an integral polymatroid rank function. Then the vertices of $\mathcal{P}_\varphi$ are integral.*

Given a matroid $\mathcal{M}$ of rank r it is easy to see that $\mathcal{P}_\mathcal{M}$ is contained in the $(n-1)$ -simplex

$$\Delta = \{ (x_1, \dots, x_n) \in \mathbb{R}^n \mid x_i \geq 0, x_1 + \dots + x_n = r. \}$$

Moreover, the following theorem of Gel'fand, Goresky, MacPherson, and Serganova gives an equivalent geometric axiomatization of matroids:

THEOREM 16 ([[GGMS87](#)]). *Let $\mathcal{B}$ be a family of r -element subsets of $[n]$. The following are equivalent:*

- (i) $\mathcal{B}$ are the bases for a matroid,
- (ii) Every edge of $\text{conv}(\mathbf{e}_B \mid B \in \mathcal{B})$ is parallel to an edge of the simplex Δ .

The following is a more useful characterization of adjacency of vertices of $\mathcal{P}_{\mathcal{M}}$, as well as a characterization of adjacency in the independence matroid polytope $\mathcal{P}_{\mathcal{M}}^{\mathcal{I}}$.

LEMMA 17 (See Theorem 4.1 in [GGMS87], Theorem 5.1 and Corollary 5.5 in [Top84]).
Let $\mathcal{M}$ be a matroid.

- (i) *Two vertices $\mathbf{e}_{B_1}$ and $\mathbf{e}_{B_2}$ are adjacent in $\mathcal{P}_{\mathcal{M}}$ if and only if $\mathbf{e}_{B_1} - \mathbf{e}_{B_2} = \mathbf{e}_i - \mathbf{e}_j$ for some i, j .*
- (ii) *If two vertices $\mathbf{e}_{I_1}$ and $\mathbf{e}_{I_2}$ are adjacent in $\mathcal{P}_{\mathcal{M}}^{\mathcal{I}}$ then $\mathbf{e}_{I_1} - \mathbf{e}_{I_2} \in \{\mathbf{e}_i - \mathbf{e}_j, \mathbf{e}_i, -\mathbf{e}_j\}$ for some i, j . Moreover if $\mathbf{v}$ is a vertex of $\mathcal{P}_{\mathcal{M}}^{\mathcal{I}}$ then all adjacent vertices of $\mathbf{v}$ can be computed in polynomial time in n , even if the matroid $\mathcal{M}$ is only presented by an evaluation oracle of its rank function φ .*

We prove one direction of (i), that every edge of $\mathcal{P}_{\mathcal{M}}$ is parallel to $\mathbf{e}_i - \mathbf{e}_j$ for some $i, j \in \mathbb{N}$. The following can be found in the proof of Proposition 2.2 in [Onn03].

PROOF. Consider any pair $A, B \in \mathcal{B}_{\mathcal{M}}$ of bases such that $[\mathbf{e}_A, \mathbf{e}_B]$ is an edge (that is, a 1-face) of $\mathcal{P}_{\mathcal{M}}$, and let $\mathbf{a} \in \mathbb{R}^n$ be a linear functional maximized over $\mathcal{P}_{\mathcal{M}}$ uniquely on that edge. If $A \setminus B = \{i\}$ is a singleton, then $B \setminus A = \{j\}$ is a singleton as well, in which case $\mathbf{e}_A - \mathbf{e}_B = \mathbf{e}_i - \mathbf{e}_j$ and we are done. Suppose then, indirectly, that it is not, and pick an element i in the symmetric difference $(A \setminus B) \cup (B \setminus A)$ of A and B of minimum value a_i . Without loss of generality assume $i \in A \setminus B$. Then there is a $j \in B \setminus A$ such that $C := A - i + j$ is a basis of $\mathcal{M}$. Since $|(A \setminus B) \cup (B \setminus A)| > 2$, C is neither A nor B . By the choice of i , this basis satisfies $\mathbf{a} \cdot \mathbf{e}_C = \mathbf{a} \cdot \mathbf{e}_A - a_i + a_j \geq \mathbf{a} \cdot \mathbf{e}_A$, and hence $\mathbf{e}_C$ is also a maximizer of $\mathbf{a}$ over $\mathcal{P}_{\mathcal{M}}$ and so lies in the 1-face $[\mathbf{e}_A, \mathbf{e}_B]$. But no $\{0, 1\}$ -vector is a convex combination of other, yielding a contradiction. $\square$

A similar property holds for integral polymatroids which we also state but we first recall some needed definitions from [Top84]. Let $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ and define $\Delta(\mathbf{v}, \mathbf{w}) := \{i \in [n] \mid v_i \neq w_i\}$ and $\text{cl}(\mathbf{v}) := \{G \mid G \subseteq [n], \sum_{i \in G} v_i = \psi(G)\}$. Let $F = \{f_1, \dots, f_{|F|}\}$ be an ordered subset of $[n]$ and $F_i := \{f_1, \dots, f_i\}$. If ψ is a polymatroid rank function then we construct $\mathbf{v} \in \mathbb{R}^n$ where $v_i = \psi(F_i) - \psi(F_{i-1})$ where $v_j = 0$ when $j \notin F$ and one says F generates $\mathbf{v}$.

LEMMA 18 (See Theorem 4.1 and Section 2 in [Top84]). *Let ψ be a polymatroid rank function. If $\mathbf{v}$ and $\mathbf{w}$ are vertices of the polymatroid $\mathcal{P}(\psi)$ then either*

- (i) $|\Delta(\mathbf{v}, \mathbf{w})| = 1$ or
- (ii) $\text{cl}(\mathbf{v}) = \text{cl}(\mathbf{w})$ and $\Delta(\mathbf{v}, \mathbf{w}) = \{c, d\}$ for some $c, d \in [n]$ where there exists some ordered set $F = \{f_1, \dots, f_{|F|}\}$ which generates $\mathbf{v}$ with $f_{k+1} = d$ and $f_k = c$ for some integer k , $1 \leq k \leq |F| - 1$; moreover the ordered set $\tilde{F} := \{f_1, \dots, f_{k-1}, f_{k+1}, f_k, f_{k+2}, \dots, f_{|F|}\}$ generates $\mathbf{w}$.

In order to describe the matroid polytope $\mathcal{P}_{\mathcal{M}}$ by a system of linear inequalities we require a few more definitions. A *circuit* $C \subseteq S$ of $\mathcal{M} = (S, \mathcal{I})$ is an inclusionwise minimal dependent set. That is, a dependent set C is a circuit of $\mathcal{M}$ if there does not exist a dependent set A such that $A \subset C \subseteq S$. A *flat* F of $\mathcal{M}$ is a subset of S such that $\varphi(F + s) = \varphi(F) + 1$ for all $s \in S$. The *span* of $A \subseteq S$ is the smallest flat F such that $A \subseteq F$. A matroid is called *connected* if for every $x, y \in S$ there exists a circuit C such that $x, y \in C$. In fact the circuits of a matroid $\mathcal{M}$ define an equivalence class on the ground set S where x, y are equivalent if there exists a circuit C such that $x, y \in C$. The *connected components* of $\mathcal{M}$ are defined as the number of such equivalence classes.

LEMMA 19 ([FS05]). *The matroid polytope $\mathcal{P}_{\mathcal{M}}$ equals the following subset of the simplex Δ :*

$$\mathcal{P}_{\mathcal{M}} = \left\{ (x_1, \dots, x_n) \in \Delta \mid \sum_{i \in F} x_i \leq \text{rank}(F) \text{ for all flats } F \subseteq S \right\}$$

PROOF. Consider any facet of the polytope $\mathcal{P}_{\mathcal{M}}$ and let $\sum_{i=1}^n a_i x_i \leq b$ be an inequality defining this facet. The normal vector $(a_1, a_2, \dots, a_n)$ is perpendicular to the edges of that facet. But each edge of the facet is parallel to some difference of unit vectors $e_i - e_j$ by Lemma 17. Hence the only constraints on the coordinates of the normal vector are of the form $a_i = a_j$. Using the equation $\sum_{i=1}^n x_i = r$ and scaling the right hand side b , we can therefore assume that $(a_1, a_2, \dots, a_n)$ is a vector in $\{0, 1\}^n$. Hence the polytope $\mathcal{P}_{\mathcal{M}}$ is characterized by the inequalities of the form $\sum_{i \in G} x_i \leq b_G$ for some $G \subseteq [n]$. The right hand side b_G equals

$$b_G = \max\{|\sigma| \cap G \mid \sigma \text{ basis of } \mathcal{M}\} = \text{rank}(G).$$

The second equality holds because every independent subset of G can be completed to a basis σ . Let F be the flat spanned by G . Then $G \subseteq F$ and $\text{rank}(G) = \text{rank}(F)$, and hence the inequality $\sum_{i \in F} x_i \leq \text{rank}(F)$ implies the inequality $\sum_{i \in G} x_i \leq \text{rank}(G)$. $\square$

LEMMA 20 ([FS05]). *The dimension of the matroid polytope $\mathcal{P}_{\mathcal{M}}$ equals $n - \#\text{connected components of } \mathcal{M}$.*

PROOF. Two elements i and j are equivalent if and only if there exist bases σ and τ with $i \in \sigma$ and $\tau = \sigma + i - j$. The linear space parallel to the affine span of $\mathcal{P}_{\mathcal{M}}$ is spanned by the vectors $\mathbf{e}_i - \mathbf{e}_j$ arising in this manner. The dimension of this space equals $n - c(M)$. $\square$

1.2.1. Matroid Optimization. Matroids find application throughout combinatorics. The *matroid intersection* problem is defined as finding the largest size common independent set of two matroids. We have seen that the greedy algorithm fully characterizes matroids and Edmonds proved that matroid intersection, can be solved efficiently. This implies that bipartite matchings, common transversals, and tree packing and covering can be computed efficiently too [Sch03]. Interestingly, it has been shown that finding matroid intersection of three or more matroids is NP-complete.

Matroids are particularly important in optimization. The greedy algorithm requires weightings on the ground set S of the matroid and it has many applications to optimization. But in real life, multiple weightings can be applied to the ground set along with a balancing function, or concrete notion of optimality over multiple weightings. Consider d weightings $\mathbf{w}_1, \dots, \mathbf{w}_d \in \mathbb{R}^n$ on the ground set $[n]$. That is, every $\mathbf{w}_i$ assigns a real-value to each element of $[n]$. We let $W \in \mathbb{R}^{d \times n}$ be the matrix with rows $\mathbf{w}_1, \dots, \mathbf{w}_d$. We also define $W\mathcal{P}_{\mathcal{M}} := \{W\mathbf{e}_B \mid B \in \mathcal{B}_{\mathcal{M}}\} \subseteq \mathbb{R}^d$, $\#W\mathcal{P}_{\mathcal{M}}$ to mean the cardinality of $W\mathcal{P}_{\mathcal{M}}$, $\overline{W\mathcal{P}_{\mathcal{M}}} := \text{conv}(W\mathcal{P}_{\mathcal{M}})$ and $\text{vert}(\mathcal{P}) = \{\text{vertices of } \mathcal{P}\}$ for a polytope $\mathcal{P}$. Intuitively one thinks of the rows of W as the set of criteria that (possibly conflicting) parties may bring to a discussion. For example, think of two groups (loggers vs. environmentalists) trying to decide how to choose an optimal spanning tree (a network of logging roads) of a graph. Each group will have different weights.

Hence in [Chapter 4](#) of this dissertation, we will consider four generalizations of the classical single-criterion (linear-objective) matroid optimization problem (i.e. $d = 1$ which is well-known to be solvable via the greedy method).

Non-Linear Matroid Optimization: Given a matroid $\mathcal{M}$ on $[n]$ with bases $\mathcal{B}_{\mathcal{M}}$, $W \in \mathbb{R}^{d \times n}$, and a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, find a base $B \in \mathcal{B}_{\mathcal{M}}$ such that $f(W\mathbf{e}_B) =$

$$\min(f(W\mathbf{e}_{B'}) \mid B' \in \mathcal{B}).$$

Two important special cases of our investigations are the following problem.

Convex Matroid Optimization: Given a matroid $\mathcal{M}$ on $[n]$ with bases $\mathcal{B}$, $W \in \mathbb{R}^{d \times n}$ where d is fixed, and a convex function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, find a base $B \in \mathcal{B}$ such that $f(W\mathbf{e}_B) = \max f(W\mathbf{e}_{B'} \mid B' \in \mathcal{B})$. Similarly we consider the minimization problem $f(W\mathbf{e}_B) = \min f(W\mathbf{e}_{B'} \mid B' \in \mathcal{B})$

Min-Max Multi-criteria Matroid Optimization: Given a matroid $\mathcal{M}$ on $[n]$ with bases $\mathcal{B}_{\mathcal{M}}$, $W \in \mathbb{R}^{d \times n}$, find a base $B \in \mathcal{B}_{\mathcal{M}}$ such that

$$\max_{i=1\dots d} ((W\mathbf{e}_B)_i) = \min(\max_{i=1\dots d} ((W\mathbf{e}_{B'})_i) \mid B' \in \mathcal{B}_{\mathcal{M}}).$$

Also, we investigate the following problem.

Pareto Multi-criteria Matroid Optimization: Given a matroid $\mathcal{M}$ on n -elements with bases $\mathcal{B}_{\mathcal{M}}$, $W \in \mathbb{R}^{d \times n}$, find a base $B \in \mathcal{B}_{\mathcal{M}}$ such that $B = \operatorname{argmin}_{Pareto} (W\mathbf{e}_{B'} \mid B' \in \mathcal{B}_{\mathcal{M}})$.

In the previous statement, $\min_{Pareto}$ is understood in the sense of Pareto optimality for problems with multiple objective functions, namely, we adopt the convention that for vectors $\mathbf{a}, \mathbf{b} \in \mathbb{R}^d$, we have $\mathbf{a} \leq \mathbf{b}$ if and only if $a_i \leq b_i$ for all entries of the vectors. Further, we say that $\mathbf{a} < \mathbf{b}$ if $\mathbf{a} \leq \mathbf{b}$ and $\mathbf{a} \neq \mathbf{b}$. We note that an optimum of the minmax problem will be a Pareto optimum. This is easy to see since if $\mathbf{a} \leq \mathbf{b}$ then $\max_i(a_i) \leq \max_i(b_i)$. The Pareto multi-criteria matroid optimization problem has been studied by several authors before. For example, Ehrgott [Ehr96] investigated two optimization problems for matroids with multiple objective functions, and he pioneered a study of Pareto bases via the base-exchange property of matroids. See [EG00] for a detailed introduction to multicriteria combinatorial optimization.

The matroid optimization problems we consider here have wide applicability. For example, in [BLMA⁺08] the authors consider the “minimum aberration model fitting” problem

in statistics, which can be reduced to a non-linear matroid optimization problem. Multi-criteria problems concerning minimum spanning trees of graphs are common in applications (see [EG00] and [KC02] and references there in).

The Min-Max optimization problem includes the NP-complete partition problem [GJ79], certain multi-processor scheduling problems [GLLR79], and specific worst-case stochastic optimization problems [War85].

EXAMPLE 21 (Multi-criteria Spanning-Tree Optimization). Consider a graph G with several linear criteria on the edges,

1. the first row of W encodes the *fixed installation cost* of each edge of G ;
2. the second row of W encodes the *monthly operating cost* of each edge of G ;
3. assuming that the edge j fails independently with probability $1 - p_j$, then by having the $\log p_j$ as the third row of W (scaled and rounded suitably), $\sum_{j \in T} \log p_j$ captures the *reliability* of the spanning tree T of G .

It can be difficult for a decision maker to balance these three competing objectives. There are many issues to consider such as the time horizon, repairability, fault tolerance, etc. These issues can be built into a concrete function f , for example a weighted norm, or can be thought of as determining a black-box f .

Unfortunately, although useful, the problems we are considering are also very difficult in general. For example, multi-criteria matroid optimization is generally NP-hard even in the case when $d = 2$ for uniform matroids. It is also NP-hard for the case of spanning trees. Nevertheless, recently many theoretical and complexity properties about these problems have been explored, yielding good complexity bounds and algorithms under nice assumptions about W, f and d . For instance, it has been shown that although convex matroid optimization is NP-complete in general, it is polynomial-time solvable under certain restrictions on W and with fixed d . We refer the reader to the recent series of papers on nonlinear matroid optimization [BLOW08, BO08, Onn03, BLMA⁺08]. This dissertation explored the experimental performance of their algorithms and proposed some new heuristics and algorithms which performed well in practice. We will see this in [Chapter 4](#).

CHAPTER 2

Volume and Ehrhart Polynomial Computation

To state our main results recall that given an integer $k > 0$ and a polytope $\mathcal{P} \subseteq \mathbb{R}^n$ we define $k\mathcal{P} := \{k\alpha \mid \alpha \in \mathcal{P}\}$ and the function $i(\mathcal{P}, k) := \#(k\mathcal{P} \cap \mathbb{Z}^n)$, where we define $i(\mathcal{P}, 0) := 1$. It is well known that for integral polytopes, as in the case of matroid polytopes, $i(\mathcal{P}, k)$ is a polynomial, called the *Ehrhart polynomial* of $\mathcal{P}$. Moreover the leading coefficient of the Ehrhart polynomial is the *normalized volume* of $\mathcal{P}$, where a unit is the volume of the fundamental domain of the affine lattice spanned by $\mathcal{P}$ [Sta96]. Our first theorem states:

THEOREM 22. *Let r be a fixed integer. Then there exist algorithms whose input data consists of a number n and an evaluation oracle for*

- (a) *a rank function φ of a matroid M on n elements satisfying $\varphi(A) \leq r$ for all A , or*
- (b) *an integral polymatroid rank function ψ satisfying $\psi(A) \leq r$ for all A ,*

that compute in time polynomial in n the Ehrhart polynomial (in particular, the volume) of the matroid polytope $\mathcal{P}(M)$, the independence matroid polytope $\mathcal{P}^{\mathcal{I}}(M)$, and the polymatroid $\mathcal{P}(\psi)$, respectively.

The computation of volumes is one of the most fundamental geometric operations and it has been investigated by several authors from the algorithmic point of view. While there are a few cases for which the volume can be computed efficiently (e.g., for convex polytopes in fixed dimension), it has been proved that computing the volume of polytopes of varying dimension is $\#P$ -hard [DF88, BW91, Kha93, Law91a]. Moreover it was proved that even approximating the volume is hard [Ele86]. Clearly, computing Ehrhart polynomials is a harder problem still. To our knowledge two previously known families of varying-dimension polytopes for which there is efficient computation of the volume are simplices or simple polytopes for which the number of vertices is polynomially bounded (this follows from Lawrence's volume formula [Law91a]). Already for simplices it is at least NP-hard to compute the whole list of coefficients of the Ehrhart polynomial, while recently [Bar06] presented a polynomial

time algorithm to compute any fixed number of the highest coefficients of the Ehrhart polynomial of a simplex of varying dimension. Theorem 22 provides another interesting family of varying dimension with volume and Ehrhart polynomial that can be computed efficiently. The proof of Theorem 22, presented in Section 2.1, relies on the geometry of tangent cones at vertices of our polytopes as well as a new refined analysis of the evaluation of Todd polynomials in the context of the computational theory of rational generating functions developed by [Bar94, BP99, BW03, Bar06, DLHTY04, DLHH⁺04, Woo04, VW08]. A nice introduction to these topics can be found at [BR07]

2.1. Computing the Ehrhart Polynomials

2.2. Preliminaries on Rational Generating Functions.

Generating functions are crucial to proving our main results. For a good reference for the basic concepts used here see [BR07]. Let $\mathcal{P} \subseteq \mathbb{R}^n$ be a rational polyhedron. The *multivariate generating function* of $\mathcal{P}$ is defined as the formal Laurent series in $\mathbb{Z}[[z_1, \dots, z_n, z_1^{-1}, \dots, z_n^{-1}]]$

$$\tilde{g}_{\mathcal{P}}(\mathbf{z}) = \sum_{\alpha \in \mathcal{P} \cap \mathbb{Z}^n} \mathbf{z}^{\alpha},$$

where we use the multi-exponent notation $\mathbf{z}^{\alpha} = \prod_{i=1}^n z_i^{\alpha_i}$. If $\mathcal{P}$ is bounded, $\tilde{g}_{\mathcal{P}}$ is a Laurent polynomial, which we consider as a rational function $g_{\mathcal{P}}$. If $\mathcal{P}$ is not bounded but is pointed (i.e., $\mathcal{P}$ does not contain a straight line), there is a non-empty open subset $U \subseteq \mathbb{C}^n$ such that the series converges absolutely and uniformly on every compact subset of U to a rational function $g_{\mathcal{P}}$ (see [BP99] and references therein). If $\mathcal{P}$ contains a straight line, we set $g_{\mathcal{P}} = 0$. The rational function $g_{\mathcal{P}} \in \mathbb{Q}(z_1, \dots, z_n)$ defined in this way is called the *multivariate rational generating function* of $\mathcal{P} \cap \mathbb{Z}^n$. Barvinok [Bar94] proved that in polynomial time, when the dimension of a polyhedron is fixed, $g_{\mathcal{P}}$ can be represented as a short sum of rational functions

$$g_{\mathcal{P}}(\mathbf{z}) = \sum_{i \in I} \varepsilon_i \frac{\mathbf{z}^{\mathbf{a}_i}}{\prod_{j=1}^n (1 - \mathbf{z}^{\mathbf{b}_{ij}})},$$

where $\varepsilon_i \in \{-1, 1\}$.

Our first contribution is to show that in the case of matroid polytopes of fixed rank, this still holds even when their dimension grows. Let $\mathbf{v}$ be a vertex of $\mathcal{P}$. Define the *tangent*

cone or *supporting cone* of $\mathbf{v}$ to be

$$\mathcal{C}_{\mathcal{P}}(\mathbf{v}) := \{ \mathbf{v} + \mathbf{w} \mid \mathbf{v} + \varepsilon \mathbf{w} \in \mathcal{P} \text{ for some } \varepsilon > 0 \}.$$

We rely on the following result, which connects the rational generating function of a rational polyhedron to those of the tangent cones of its vertices. This result was independently discovered by Brion [Bri88] and Lawrence [Law91b]. A proof can also be found in [BP99] and [BHS09].

LEMMA 23 (Brion–Lawrence’s Theorem). *Let $\mathcal{P}$ be a rational polyhedron and $V(\mathcal{P})$ be the set of vertices of $\mathcal{P}$. Then,*

$$g_{\mathcal{P}}(\mathbf{z}) = \sum_{\mathbf{v} \in V(\mathcal{P})} g_{\mathcal{C}_{\mathcal{P}}(\mathbf{v})}(\mathbf{z}),$$

where $\mathcal{C}_{\mathcal{P}}(\mathbf{v})$ is the tangent cone of $\mathbf{v}$.

Thus, we can write the multivariate generating function of $\mathcal{P}$ by writing all multivariate generating functions of all the tangent cones of the vertices of $\mathcal{P}$. Moreover, the map assigning to a rational polyhedron $\mathcal{P}$ its multivariate rational generating function $g_{\mathcal{P}}(\mathbf{z})$ is a *valuation*, i.e., a finitely additive measure, so it satisfies the equation

$$g_{\mathcal{P}_1 \cup \mathcal{P}_2}(\mathbf{z}) = g_{\mathcal{P}_1}(\mathbf{z}) + g_{\mathcal{P}_2}(\mathbf{z}) - g_{\mathcal{P}_1 \cap \mathcal{P}_2}(\mathbf{z}),$$

for arbitrary rational polytopes $\mathcal{P}_1$ and $\mathcal{P}_2$, the so-called *inclusion–exclusion principle*. This allows to break a polyhedron $\mathcal{P}$ into pieces $\mathcal{P}_1$ and $\mathcal{P}_2$ and to compute the multivariate rational generating functions for the pieces (and their intersection) separately in order to get the generating function $g_{\mathcal{P}}$. More generally, let us denote by $[\mathcal{P}]$ the *indicator function* of $\mathcal{P}$, i.e., the function

$$[\mathcal{P}]: \mathbb{R}^n \rightarrow \mathbb{R}, \quad [\mathcal{P}](\mathbf{x}) = \begin{cases} 1 & \text{if } \mathbf{x} \in \mathcal{P} \\ 0 & \text{otherwise.} \end{cases}$$

Let $\sum_{i \in I} \varepsilon_i [\mathcal{P}_i] = 0$ be an arbitrary linear identity of indicator functions of rational polyhedra (with rational coefficients ε_i); the valuation property now implies that it carries over to a linear identity $\sum_{i \in I} \varepsilon_i g_{\mathcal{P}_i}(\mathbf{z}) = 0$ of rational generating functions.

Now let $\mathcal{C}$ be one of the tangent cones of $\mathcal{P}$, and let $\mathcal{T}$ be a triangulation of $\mathcal{C}$, given by its simplicial cones of maximal dimension. Let $\hat{\mathcal{T}}$ denote the set of all (lower-dimensional) intersecting proper faces of the cones $\mathcal{C}_i \in \mathcal{T}$. Then we can assign an integer coefficient ε_i to every cone $\mathcal{C}_i \in \hat{\mathcal{T}}$, such that the following identity holds:

$$[\mathcal{C}] = \sum_{\mathcal{C}_i \in \mathcal{T}} [\mathcal{C}_i] + \sum_{\mathcal{C}_i \in \hat{\mathcal{T}}} \varepsilon_i [\mathcal{C}_i].$$

This identity immediately carries over to an identity of multivariate rational generating functions,

$$(2.1) \quad g_{\mathcal{C}}(\mathbf{z}) = \sum_{\mathcal{C}_i \in \mathcal{T}} g_{\mathcal{C}_i}(\mathbf{z}) + \sum_{\mathcal{C}_i \in \hat{\mathcal{T}}} \varepsilon_i g_{\mathcal{C}_i}(\mathbf{z}).$$

Hence, the problem of computing rational generating functions of a polyhedron is reduced to the case of simplicial cones.

In the following (section 2.3, section 2.4), we study the tangent cones of the matroid polytopes and polymatroids and introduce algorithms that construct triangulations for them. Then, in section 2.5, we construct a short multivariate generating function using an *efficient* variant of identity (2.1) and Brion–Lawrence’s Theorem. Finally, in section 2.6, we compute the Ehrhart polynomial.

2.3. On the Tangent Cones of Matroid Polytopes

Our goal is to compute the multivariate generating function of matroid polytopes and independence matroid polytopes with fixed rank (later, in section 2.4, we will deal with the case of polymatroids), and to do this we will use a crucial property of adjacent vertices. To illustrate our techniques we will use a running example throughout this section.

EXAMPLE 24 (Matroid on K_4). Let K_4 be the complete graph on four vertices. Label the $\binom{4}{2} = 6$ edges with $\{1, \dots, 6\}$ as in Figure 2.1. Every graph induces a matroid on its edges where the bases are all spanning trees (spanning forests for disconnected graphs) [Wel76]. Let $M(K_4)$ be the matroid on the elements $\{1, \dots, 6\}$ with bases as all spanning trees of K_4 . The rank of $M(K_4)$ is the size of any spanning tree of K_4 , thus the rank of $M(K_4)$ is 3. The 16 bases of $M(K_4)$ are: $\{3, 5, 6\}$, $\{3, 4, 6\}$, $\{3, 4, 5\}$, $\{2, 5, 6\}$, $\{2, 4, 6\}$,

$\{2, 4, 5\}, \{2, 3, 5\}, \{2, 3, 4\}, \{1, 5, 6\}, \{1, 4, 6\}, \{1, 4, 5\}, \{1, 3, 6\}, \{1, 3, 4\}, \{1, 2, 6\}, \{1, 2, 5\}, \{1, 2, 3\}.$

Let M be a matroid on n elements with fixed rank r . Then the number of vertices of $\mathcal{P}(M)$ is a polynomial in n of degree r . We can see this since the number of vertices is equal to the number of bases of M , and the number of bases is bounded by $\binom{n}{r}$, a polynomial in n of degree r . Clearly the number of vertices of $\mathcal{P}^{\mathcal{I}}(M)$ is also polynomial in n . It is also clear that, when the rank r is fixed, all vertices of either polytope can be enumerated in polynomial time in n , even when the matroid is only presented by an evaluation oracle for its rank function φ .

Throughout this section we shall discuss polyhedral cones $\mathcal{C}$ with extremal rays $\{\mathbf{r}_1, \dots, \mathbf{r}_l\}$ such that

$$\mathbf{r}_k \in R_A := \{ \mathbf{e}_i - \mathbf{e}_j, \mathbf{e}_i, -\mathbf{e}_j \mid i \in [n], j \in A \} \quad \text{for } k = 1, \dots, l$$

for some $A \subseteq [n]$. We will refer to R_A as the *elementary set of A* . Note that by [Lemma 17](#) the rays of a tangent cone at a vertex $\mathbf{e}_A$ (corresponding to a set $A \subseteq [n]$) of a matroid polytope or an independence matroid polytope form an elementary set of A . Due to convexity and the assumption that $\mathbf{r}_k$ are extremal, for each $i \in [n]$ and $j \in A$ at most two of the three vectors $\mathbf{e}_i - \mathbf{e}_j, \mathbf{e}_i, -\mathbf{e}_j$ are extremal rays $\mathbf{r}_k$ of $\mathcal{C}$. This implies by construction, that considering all pairs $\mathbf{e}_i - \mathbf{e}_j$ and $\mathbf{e}_i$ or $-\mathbf{e}_j$, the number of generators $\mathbf{r}_k$ of $\mathcal{C}$ is bounded by

$$(2.2) \quad n|A| + n + |A|.$$

Recall a cone is *simple* if it is generated by linearly independent vectors and it is *unimodular* if its fundamental parallelepiped contains only one lattice point [[GO97](#)]. A triangulation of $\mathcal{C}$ is *unimodular* if it is a polyhedral subdivision such that each sub-cone is unimodular.

EXAMPLE 25 (Matroid on K_4). The vertices $\mathbf{e}_{\{2,3,5\}}, \mathbf{e}_{\{2,3,4\}}, \mathbf{e}_{\{1,3,6\}}, \mathbf{e}_{\{1,3,4\}}, \mathbf{e}_{\{1,2,6\}}$ and $\mathbf{e}_{\{1,2,5\}}$ are all adjacent to the vertex $\mathbf{e}_{\{1,2,3\}}$, see [Figure 2.1](#). Moreover, the tangent




FIGURE 2.1. $\{2, 3, 5\}$, $\{2, 3, 4\}$, $\{1, 3, 6\}$, $\{1, 3, 4\}$, $\{1, 2, 6\}$ and $\{1, 2, 5\}$ are spanning trees of K_4 that differ from $\{1, 2, 3\}$ by adding one edge and removing one edge.

cone $\mathcal{C}_{\mathcal{P}(M(\mathbf{K}_4))}(\mathbf{e}_{\{1,2,3\}})$ is generated by the differences of these vertices with $\mathbf{e}_{\{1,2,3\}}$:

$$\begin{aligned} \mathcal{C}_{\mathcal{P}(M(\mathbf{K}_4))}(\mathbf{e}_{\{1,2,3\}}) = & \mathbf{e}_{\{1,2,3\}} + \text{cone} \{ \mathbf{e}_{\{2,3,5\}} - \mathbf{e}_{\{1,2,3\}}, \mathbf{e}_{\{2,3,4\}} - \mathbf{e}_{\{1,2,3\}}, \\ & \mathbf{e}_{\{1,3,4\}} - \mathbf{e}_{\{1,2,3\}}, \mathbf{e}_{\{1,2,6\}} - \mathbf{e}_{\{1,2,3\}}, \mathbf{e}_{\{1,2,5\}} - \mathbf{e}_{\{1,2,3\}} \} \end{aligned}$$

LEMMA 26. Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a cone generated by p extremal rays $\{\mathbf{r}_1, \dots, \mathbf{r}_p\} \subseteq R_A$ where R_A is an elementary set of some $A \subseteq [n]$. Every triangulation of $\mathcal{C}$ is unimodular.

PROOF. Without loss of generality, we can assume $\{\mathbf{r}_1, \dots, \mathbf{r}_l\}$ are generators of the form $\mathbf{e}_i - \mathbf{e}_j$ and $\{\mathbf{r}_{l+1}, \dots, \mathbf{r}_p\}$ are generators of the form $\mathbf{e}_i$ or $-\mathbf{e}_j$ for the cone $\mathcal{C}$.

It is easy to see that the matrix $\tilde{T}_{\mathcal{C}} := [\mathbf{r}_1, \dots, \mathbf{r}_l]$ is totally unimodular. Let $G_{\mathcal{C}}$ be a directed graph with vertex set $[n]$ and an edge from vertex i to j if $\mathbf{r}_k = \mathbf{e}_i - \mathbf{e}_j$ is an extremal ray of $\mathcal{C}$. We can see that $G_{\mathcal{C}}$ is a subgraph of the complete directed graph K_n with two arcs between each pair of vertices; one for each direction. Since $\tilde{T}_{\mathcal{C}} := [\mathbf{r}_1, \dots, \mathbf{r}_l]$ is the incidence matrix of the graph $G_{\mathcal{C}}$, it is totally unimodular [Sch86b, Ch. 19, Ex. 2], i.e., every subdeterminant is 0, 1 or -1 [Sch86b, Ch. 19, Thm. 9]. Therefore $T_{\mathcal{C}} := [\mathbf{r}_1, \dots, \mathbf{r}_l, \mathbf{r}_{l+1}, \dots, \mathbf{r}_p]$ is totally unimodular since augmenting $\tilde{T}_{\mathcal{C}}$ by a vector $\mathbf{e}_i$ or $-\mathbf{e}_j$ preserves this subdeterminant property: for any submatrix containing part of a vector $\mathbf{e}_i$ or $-\mathbf{e}_j$ perform the cofactor expansion down the vector $\mathbf{e}_i$ or $-\mathbf{e}_j$ when calculating the determinant.

Since $T_{\mathcal{C}}$ is totally unimodular, each basis of $T_{\mathcal{C}}$ generates the entire integer lattice $\mathbb{Z}^n \cap \text{lin}(\mathcal{C})$ and hence every simplicial cone of a triangulation has normalized volume 1. $\square$

LEMMA 27. *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a cone generated by l extremal rays $\{\mathbf{r}_1, \dots, \mathbf{r}_l\} \subseteq R_A$ where R_A is an elementary set of some $A \subseteq [n]$, where $\dim(\mathcal{C}) < n$. The extremal rays $\{\mathbf{r}_1, \dots, \mathbf{r}_l\}$ can be augmented by a vector $\tilde{\mathbf{r}}$ such that $\dim(\text{cone}\{\mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}\}) = \dim(\mathcal{C}) + 1$, the vectors $\mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}$ are all extremal, and $\tilde{\mathbf{r}} \in R_A$.*

PROOF. It follows from convexity that at most two of $\mathbf{e}_i - \mathbf{e}_j$, $\mathbf{e}_i$ or $-\mathbf{e}_j$ are extremal generators of $\mathcal{C}$ for $i \in [n]$ and $j \in A$. There are at least n possible extremal ray generators, considering two of $\mathbf{e}_i - \mathbf{e}_j$, $\mathbf{e}_i$ or $-\mathbf{e}_j$ for each $i \in [n]$ and $j \in A$. Moreover, all these pairs span $\mathbb{R}^n$. Thus by the basis augmentation theorem of linear algebra, there exists a vector $\tilde{\mathbf{r}}$ such that $\dim(\text{cone}\{\mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}\}) = \dim(\mathcal{C}) + 1$ and $\mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}$ are all extremal. $\square$

LEMMA 28. *Let r be a fixed integer, n be an integer, $A \subseteq [n]$ with $|A| \leq r$ and let $\mathcal{C} \subseteq \mathbb{R}^n$ be a cone generated by l extremal rays $\{\mathbf{r}_1, \dots, \mathbf{r}_l\} \subseteq R_A$ where R_A is an elementary set of A . Then any triangulation of $\text{conv}(\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l\})$ has at most a polynomial in n number of top-dimensional simplices.*

PROOF. Assume $\dim(\mathcal{C}) = n$. Later, we will show how to remove this restriction. We can see that $\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l\} \subseteq [-1, 1]^A \times \tilde{\Delta}_{[n] \setminus A}$ where

$$[-1, 1]^A := \{\mathbf{x} \in \mathbb{R}^A \mid |\mathbf{x}_j| \leq 1 \ j \in A\} \subseteq \mathbb{R}^A$$

$$\tilde{\Delta}_{[n] \setminus A} := \text{conv}(\{\mathbf{e}_i \mid i \in [n] \setminus A\} \cup \{\mathbf{0}\}) \subseteq \mathbb{R}^{[n] \setminus A}.$$

The volume of a d -simplex $\text{conv}\{\mathbf{v}_0, \dots, \mathbf{v}_d\}$ is [GO97]

$$(2.3) \quad \frac{1}{d!} \left| \det \begin{pmatrix} \mathbf{v}_0 & \cdots & \mathbf{v}_d \\ \mathbf{1} & \cdots & \mathbf{1} \end{pmatrix} \right|.$$

Thus the $(n - |A|)$ -volume of $\tilde{\Delta}_{[n] \setminus A}$ is $\frac{1}{(n - |A|)!}$ and the $|A|$ -volume of $[-1, 1]^A$ is $2^{|A|}$. Therefore

$$\text{vol}([-1, 1]^A \times \tilde{\Delta}_{[n] \setminus A}) = 2^{|A|} \frac{1}{(n - |A|)!} = \frac{1}{n!} 2^{|A|} n(n - 1) \cdots (n - |A| + 1).$$

It is also a fact that any integral n -simplex has n -volume bounded below by $\frac{1}{n!}$, using the simplex volume equation (2.3). Therefore any triangulation of $\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l\}$ has at

most

$$2^{|A|} n(n-1) \cdots (n-|A|+1) \leq 2^r n(n-1) \cdots (n-r+1)$$

full-dimensional simplices, a polynomial function in n of degree r .

Let $d_{\mathcal{C}} := n - \dim(\mathcal{C})$. If $\dim(\mathcal{C}) < n$, then by Lemma 27, $\{\mathbf{r}_1, \dots, \mathbf{r}_l\}$ can be augmented with vectors $\{\tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{d_{\mathcal{C}}}\}$ where $\tilde{\mathbf{r}}_k \in R_A$ for A above, such that $\dim(\text{cone}\{\mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{d_{\mathcal{C}}}\}) = n$ and $\{\mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{d_{\mathcal{C}}}\}$ are extremal. Moreover,

$$\begin{aligned} \dim(\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l\}) &< \dim(\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1\}) < \cdots \\ &< \dim(\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{d_{\mathcal{C}}-1}\}) < \dim(\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{d_{\mathcal{C}}}\}), \end{aligned}$$

that is, $\tilde{\mathbf{r}}_k \notin \text{aff}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{k-1}\}$ for $1 \leq k \leq d_{\mathcal{C}}$.




FIGURE 2.2. If $\tilde{\mathbf{r}}_k$ is not contained in the affine span of $\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{k-1}\}$ then every full-dimensional simplex must contain $\tilde{\mathbf{r}}_k$.

Since $\tilde{\mathbf{r}}_k \notin \text{aff}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{k-1}\}$, any full-dimensional simplex in a triangulation of $\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_k\}$ must contain $\tilde{\mathbf{r}}_k$, see Figure 2.2. If not, then there exists a top-dimensional simplex using the points $\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{k-1}\}$, but we know all these points lie in a subspace of one less dimension, a contradiction. Therefore, a bound on the number of simplices in a triangulation of $\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_k\}$ is a bound on that of $\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{k-1}\}$.

Thus, if $\dim(\mathcal{C}) < n$ we can augment $\mathcal{C}$ by vectors $\tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{d_{\mathcal{C}}}$ so that the cone $\tilde{\mathcal{C}} := \text{cone}\{\mathbf{r}_1, \dots, \mathbf{r}_p, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{d_{\mathcal{C}}}\}$ is of dimension n and $\mathbf{r}_l, \tilde{\mathbf{r}}_k \in R_A$ for A above. We proved any triangulation of $\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_p, \tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_{d_{\mathcal{C}}}\}$ has at most polynomially many full-dimensional n -simplices, which implies that any triangulation of $\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_p\}$ has at most polynomially many top-dimensional simplices due to the construction of the generators $\tilde{\mathbf{r}}_k$. $\square$

We have shown that for a cone $\mathcal{C}$ generated by an elementary set of extremal rays $\{\mathbf{r}_1, \dots, \mathbf{r}_l\} \subseteq R_A$ for some $A \subseteq [n]$, any triangulation of $\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l\}$ has at most polynomially many simplices. What we need next is an efficient method to compute some triangulation of $\text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l\}$. We will show that the placing triangulation is a suitable candidate.

Let $\mathcal{P} \subseteq \mathbb{R}^n$ be a polytope of dimension n and Δ be a facet of $\mathcal{P}$ and $\mathbf{v} \in \mathbb{R}^n$. There exists a unique hyperplane H containing Δ and $\mathcal{P}$ is contained in one of the closed sides of H , call it H^+ . If $\mathbf{v}$ is contained in the interior of H^- , the other closed halfspace defined by H , then Δ is *visible* from $\mathbf{v}$ (see chapter 14.2 in [GO97]). The well-known *placing triangulation* is given by an algorithm where a point is iteratively added to an intermediate triangulation by determining which facets are visible to the new point [GO97, DLRS09]. We recall now how to determine if a facet is visible to a vertex in polynomial time.




FIGURE 2.3. Figure (a) shows Δ visible to $\mathbf{v}$. Figure (b) shows Δ **not** visible to $\mathbf{v}$.

LEMMA 29. Let $\mathcal{P} \subseteq \mathbb{R}^n$ be a polytope given by t vertices $\{\mathbf{v}^1, \dots, \mathbf{v}^t\} \subseteq \mathbb{R}^n$ and $\Delta \subseteq \mathcal{P}$ be a facet of $\mathcal{P}$ given by q vertices $\{\tilde{\mathbf{v}}^1, \dots, \tilde{\mathbf{v}}^q\} \subseteq \{\mathbf{v}^1, \dots, \mathbf{v}^t\}$. If $\mathbf{v} \in \mathbb{R}^n$ where $\mathbf{v} \notin \mathcal{P}$ then deciding if Δ is visible to $\mathbf{v}$ can be done in polynomial time in the input $\{\tilde{\mathbf{v}}^1, \dots, \tilde{\mathbf{v}}^q\}$, $\{\mathbf{v}^1, \dots, \mathbf{v}^t\}$ and $\mathbf{v}$.

PROOF. Let $\mathbf{z} := \frac{1}{q} \sum_{i=1}^q \tilde{\mathbf{v}}^i$ so that $\mathbf{z} \in \text{relint}(\Delta)$. We consider the linear program:

$$(2.4) \quad \left\{ \begin{pmatrix} \mathbf{x} \\ \mathbf{y} \\ \lambda \end{pmatrix} \in \mathbb{R}^{n+t+1} \mid \mathbf{x} = \sum_{i=1}^t \mathbf{v}^i y_i, \mathbf{y} \geq \mathbf{0}, \sum_{i=1}^t y_i = 1, \right. \\ \left. 0 \leq \lambda < 1, \lambda \mathbf{v} + (1 - \lambda)\mathbf{z} = \mathbf{x} \right\}.$$

If (2.4) has a solution then there exists a point $\bar{\mathbf{x}} \in \mathcal{P}$ between the facet Δ and $\mathbf{v}$, hence Δ is not visible from $\mathbf{v}$. If (2.4) does not have a solution, then there are no points of $\mathcal{P}$ between $\mathbf{v}$ and Δ , hence Δ is visible from $\mathbf{v}$ (see Lemma 4.2.1 in [DLRS09]). It is well known that a strict inequality, such as the one in (2.4), can be handled by an equivalent linear program which has only one additional variable. Determining if (2.4) has a solution can be done in polynomial time in the input [Sch86b]. $\square$

The placing triangulation is obtained by incrementally adding one point at a time, connecting the new point to the current triangulation. More precisely:

ALGORITHM 30 (The Placing Triangulation [GO97, DLRS09]).

Input: A set of ordered points $\{\mathbf{v}_1, \dots, \mathbf{v}_t\} \in \mathbb{R}^n$.

Output: A triangulation $\mathcal{T}$ of $\{\mathbf{v}_1, \dots, \mathbf{v}_t\}$

```

1:  $\mathcal{T} := \{\{\mathbf{v}_1\}\}$ .
2: for each  $\mathbf{v}_i \in \{\mathbf{v}_2, \dots, \mathbf{v}_t\}$  do
3:   Let  $B \in \mathcal{T}$ .
4:    $P_i := \{\mathbf{v}_1, \dots, \mathbf{v}_{i-1}\}$ 
5:   if  $\mathbf{v}_i \notin \text{aff}(P_i)$  then
6:      $\mathcal{T}' := \emptyset$ 
7:     for each  $D \in \mathcal{T}$  do
8:        $\mathcal{T}' := \mathcal{T}' \cup \{D \cup \{\mathbf{v}_i\}\}$ .
9:   else
10:    for each  $B \in \mathcal{T}$  and each  $(|B| - 1)$ -subset  $C$  of  $B$  do
11:      Create and solve the linear program (2.4) with  $(P_i, C, \mathbf{v}_i)$  to decide
        visibility of  $C$  to  $\mathbf{v}_i$ .
12:      if  $C$  is visible to  $\mathbf{v}_i$  then
13:         $\mathcal{T}' := \mathcal{T}' \cup \{C \cup \{\mathbf{v}_i\}\}$ 
14:    $\mathcal{T} := \mathcal{T}'$ 
15: return  $\mathcal{T}$ 

```

Indeed, Algorithm 30 returns a triangulation [GO97]. We will show that for certain input (a point set corresponding to a vertex cone of a matroid polytope or independence matroid polytope), it runs in polynomial time. We remark that there are exponentially, in n , many lower dimensional simplices in any given triangulation. But, it is important to note that only the highest dimensional simplices are listed in an intermediate triangulation (and thus the final triangulation) in the placing triangulation algorithm.

THEOREM 31. *Let r be a fixed integer, n be an integer, $A \subseteq [n]$ with $|A| \leq r$, and let $\{\mathbf{r}_1, \dots, \mathbf{r}_l\} \subseteq R_A$. Then the placing triangulation (Algorithm 30) with input $\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l\}$ runs in polynomial time.*

PROOF. By Equation (2.2) there is only a polynomial, in n , number of extremal rays $\{\mathbf{r}_1, \dots, \mathbf{r}_l\}$. Thus, the **for** statement on line 2 repeats a polynomial number of times. Step 5 can be done in polynomial time by solving the linear equation $[\mathbf{r}_1, \dots, \mathbf{r}_{i-1}]\mathbf{x} = \mathbf{v}_i$.

The **for** statement on line 7 repeats for every simplex D in the triangulation $\mathcal{T}$, and the number of simplices in $\mathcal{T}$ is bounded by the number of simplices in the final triangulation. By Lemma 28 any triangulation of extremal cone generators in R_A with the origin will use at most polynomially many top-dimensional simplices. Hence the number of top-dimensional simplices of any partial triangulation $\mathcal{T}$ will be polynomially bounded since it is a subset of the final triangulation.

The **for** statement on line 10 repeats for every simplex B and every $(|B| - 1)$ -simplex of B . As before, the number of simplices B is polynomially bounded, and there are at most n $(|B| - 1)$ -simplices of B . Thus the **for** statement will repeat a polynomial number of times.

Finally, by Lemma 29, determining if C is visible to $\mathbf{v}_i$ can be done in polynomial time. Therefore Algorithm 30 runs in a polynomial time. $\square$

COROLLARY 32. *Let r be a fixed integer, n be an integer, $A \subseteq [n]$ with $|A| \leq r$, and let $\mathcal{C} \subseteq \mathbb{R}^n$ be a cone generated by extremal rays $\{\mathbf{r}_1, \dots, \mathbf{r}_l\} \subseteq R_A$. A triangulation of $\mathcal{C}$ can be computed in polynomial time in the input of the extremal ray generators $\{\mathbf{r}_1, \dots, \mathbf{r}_l\}$.*

PROOF. Let $\mathcal{P}_{\mathcal{C}} := \text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l\}$. We give an algorithm which produces a triangulation of $\mathcal{P}_{\mathcal{C}} := \text{conv}\{\mathbf{0}, \mathbf{r}_1, \dots, \mathbf{r}_l\}$ such that each full-dimensional simplex has $\mathbf{0}$ as a vertex. Such a triangulation would extend to a triangulation of the cone $\mathcal{C}$. This can be accomplished by applying two placing triangulations: one to triangulate the boundary of $\mathcal{P}_{\mathcal{C}}$ not incident to $\mathbf{0}$, and another to attach the triangulated boundary faces to $\mathbf{0}$. The algorithm goes as follows:

- 1) **Triangulate $\mathcal{P}_{\mathcal{C}}$ using the placing triangulation algorithm. Call it $\mathcal{T}'$.**
- 2) **Triangulate $\mathcal{P}_{\mathcal{C}}$ using the boundary faces of $\mathcal{T}'$ which do not contain v .**

ALGORITHM 33 (Triangulation joining $\mathbf{0}$ to boundary faces).

Input: A triangulation $\mathcal{T}'$ of $\mathcal{P}_{\mathcal{C}}$, given by its vertices.

Output: A triangulation $\mathcal{T}$ of $\mathcal{P}_{\mathcal{C}}$ such that every highest dimension simplex of $\mathcal{T}$ is incident to $\mathbf{0}$.

```

1:  $\mathcal{T} := \emptyset$ 
2: for each  $C$  where  $C$  is a  $(|B| - 1)$ -simplex of  $B \in \mathcal{T}'$  do
3:   if  $C$  is not a  $(|A| - 1)$ -simplex of  $A \in \mathcal{T}'$  where  $A \neq B$  then
4:      $\mathcal{T} := \mathcal{T} \cup \{C \cup \{\mathbf{0}\}\}$ 
5: return  $\mathcal{T}$ 

```




FIGURE 2.4. A triangulation $\mathcal{T}'$ of $\mathcal{P}_{\mathcal{C}}$ can be used to extend to a triangulation $\mathcal{T}$ such that $\mathbf{0}$ is incident to every highest dimensional simplex

By Theorem 31, triangulating $\mathcal{P}_{\mathcal{C}}$ using Algorithm 30 can be done in polynomial time. Algorithm 33 indeed produces a triangulation of $\mathcal{P}_{\mathcal{C}}$. It covers $\mathcal{P}_{\mathcal{C}}$ since every extremal ray generator $\mathbf{r}_k$ of $\mathcal{C}$ is on some $(\dim(\mathcal{C}) - 1)$ -simplex. Moreover, $\mathcal{T}$ by construction has the property that the intersection of any two simplices of $\mathcal{T}$ is a simplex. Step 3 checks if C is on the boundary, since if C is on the boundary it will not be on the intersection of two higher-dimensional simplices.

Step 2 repeats a polynomial number of times since any triangulation of $\mathcal{P}_{\mathcal{C}}$ has at most a polynomial number of simplices, and each simplex B has at most $n(|B| - 1)$ -simplices. Step 3 can be computed in polynomial time since again there are only polynomially many simplices B in the triangulation $\mathcal{T}'$ and at most $n(|B| - 1)$ -simplices to check if they are equal to C . Hence, Algorithm 33 runs in polynomial time. $\square$

EXAMPLE 34. The tangent cone at the vertex $\mathbf{e}_B := \mathbf{e}_{\{1,2,3\}}$ on the polytope $\mathcal{P}(M(K_4))$ can be triangulated as:

$$\begin{aligned} & \left\{ \left\{ \mathbf{e}_{\{2,3,5\}} - \mathbf{e}_B, \mathbf{e}_{\{2,3,4\}} - \mathbf{e}_B, \mathbf{e}_{\{1,3,6\}} - \mathbf{e}_B, \mathbf{e}_{\{1,3,4\}} - \mathbf{e}_B, \mathbf{e}_{\{1,2,6\}} - \mathbf{e}_B \right\}, \right. \\ & \quad \left\{ \mathbf{e}_{\{2,3,5\}} - \mathbf{e}_B, \mathbf{e}_{\{1,3,6\}} - \mathbf{e}_B, \mathbf{e}_{\{1,3,4\}} - \mathbf{e}_B, \mathbf{e}_{\{1,2,6\}} - \mathbf{e}_B, \mathbf{e}_{\{1,2,5\}} - \mathbf{e}_B \right\}, \\ & \quad \left. \left\{ \mathbf{e}_{\{2,3,5\}} - \mathbf{e}_B, \mathbf{e}_{\{2,3,4\}} - \mathbf{e}_B, \mathbf{e}_{\{1,3,4\}} - \mathbf{e}_B, \mathbf{e}_{\{1,2,6\}} - \mathbf{e}_B, \mathbf{e}_{\{1,2,5\}} - \mathbf{e}_B \right\} \right\}. \end{aligned}$$

2.4. Polymatroids

We will show that certain lemmas from Subsection 2.3 also hold for certain polymatroids. Recall that the rank of the matroid M is the size of any basis of M which equals $\varphi([n])$. Our lemmas from Subsection 2.3 rely on the fact that M has fixed rank, that is, for some $r \in \mathbb{Z}$, $r \geq 0$, $\varphi(A) \leq r$ for all $A \subseteq [n]$. We will show that a similar condition on a polymatroid rank function is sufficient for the lemmas of Subsection 2.3 to hold.

LEMMA 35. *Let $\psi: 2^{[n]} \rightarrow \mathbb{N}$ be an integral polymatroid rank function where $\psi(A) \leq r$ for all $A \subseteq [n]$, where r is a fixed integer. Then the number of vertices of $\mathcal{P}(\psi)$ is bounded by a polynomial in n of degree r .*

PROOF. It is known that if ψ is integral then all vertices of $\mathcal{P}(\psi)$ are integral [Wel76]. The number of vertices of $\mathcal{P}(\psi)$ can be bounded by the number of non-negative integral solutions to $x_1 + \dots + x_n \leq r$, which has $\binom{n+r}{r}$ solutions, a polynomial in n of degree r [Sta97]. $\square$

LEMMA 36. *Let $\psi: 2^{[n]} \rightarrow \mathbb{N}$ be an integral polymatroid rank function. If $\mathbf{v}$ is a vertex of $\mathcal{P}(\psi)$ then all adjacent vertices of $\mathbf{v}$ can be enumerated in polynomial time. Moreover if $\psi(A) \leq r$ for all $A \subseteq [n]$, where r is a fixed integer, then the vertices of $\mathcal{P}(\psi)$ can be enumerated in polynomial time.*

PROOF. If $\mathbf{v}$ is a vertex of $\mathcal{P}(\psi)$ then generating and listing all adjacent vertices to $\mathbf{v}$ can be done in polynomial time by Lemma 17. If $\psi(A) \leq r$ for all $A \subseteq [n]$, where r is a fixed integer, then, by Lemma 35, there is a polynomial number of vertices for $\mathcal{P}(\psi)$. We know that $\mathbf{0} \in \mathbb{R}^n$ is a vertex of any polymatroid. Therefore, beginning with $\mathbf{0}$, we can perform a breadth-first search, which is output-sensitive polynomial time, on the graph of $\mathcal{P}(\psi)$, enumerating all vertices of $\mathcal{P}(\psi)$. $\square$

What remains to be shown is that these polymatroids have cones like the ones in Subsection 2.3.

LEMMA 37. *Let ψ be an integral polymatroid rank function and $\mathcal{C}$ the tangent cone of a vertex $\mathbf{v}$ of the polymatroid $\mathcal{P}(\psi)$, translated to the origin. Then $\mathcal{C}$ is generated by extremal ray generators $\{\mathbf{r}_1, \dots, \mathbf{r}_l\} \subseteq R_{\text{supp}(\mathbf{v})}$, where $R_{\text{supp}(\mathbf{v})}$ is an elementary set of $\text{supp}(\mathbf{v})$.*

PROOF. Let $\psi: 2^{[n]} \rightarrow \mathbb{Z}$ be a integral polymatroid rank function. Let $\mathbf{v}$ and $\mathbf{w}$ be adjacent vertices of the polymatroid $\mathcal{P}(\psi)$. Using Lemma 18, if $|\Delta(\mathbf{v}, \mathbf{w})| = 1$ then $\mathbf{w} - \mathbf{v} = h\mathbf{e}_i$ where h is some integer and $\mathbf{e}_i$ is the standard i th elementary vector for some $i \in [n]$. If $h < 0$ then certainly $i \in \text{supp}(\mathbf{v})$, else $i \in [n]$. Thus $\mathbf{w} - \mathbf{v}$, a generator of $\mathcal{C}$, is parallel to a vector in $R_{\text{supp}(\mathbf{v})}$.

Let $\mathbf{v}$ and $\mathbf{w}$ be adjacent and satisfy (ii) in Lemma 18, where $\Delta(\mathbf{v}, \mathbf{w}) = \{c, d\}$. Hence there exists an $F = \{f_1, \dots, f_{|F|}\}$ which generates $\mathbf{v}$ with $f_{k+1} = d$ and $f_k = c$ for some integer k , $1 \leq k \leq |F| - 1$; moreover the ordered set $\tilde{F} := \{f_1, \dots, f_{k-1}, f_{k+1}, f_k, f_{k+2}, \dots, f_{|F|}\}$ generates $\mathbf{w}$. First we note that $\psi(F_{k-1}) = \psi(\tilde{F}_{k-1})$ and $\psi(F_{k+1}) = \psi(\tilde{F}_{k+1})$. By assumption, we know $v_c \neq w_c$, $v_d \neq w_d$ and $v_l = w_l$ for all $l \in [n] \setminus \{c, d\}$. Thus

$$\begin{aligned} (\mathbf{v} - \mathbf{w})_c &= v_c - w_c &= \psi(F_{k+1}) - \psi(F_k) - (\psi(\tilde{F}_k) - \psi(\tilde{F}_{k-1})) \\ &= \psi(F_{k+1}) - \psi(F_k) - \psi(\tilde{F}_k) + \psi(\tilde{F}_{k-1}) \end{aligned}$$

and

$$\begin{aligned} (\mathbf{v} - \mathbf{w})_d &= v_d - w_d &= \psi(F_k) - \psi(F_{k-1}) - (\psi(\tilde{F}_{k+1}) - \psi(\tilde{F}_k)) \\ &= \psi(F_k) - \psi(\tilde{F}_{k-1}) - (\psi(F_{k+1}) - \psi(\tilde{F}_k)) \\ &= -\psi(F_{k+1}) + \psi(F_k) + \psi(\tilde{F}_k) - \psi(\tilde{F}_{k-1}). \end{aligned}$$

Therefore $(\mathbf{v} - \mathbf{w})_c = -(\mathbf{v} - \mathbf{w})_d$ and $\mathbf{w} - \mathbf{v}$ is parallel to $\mathbf{e}_d - \mathbf{e}_c$. Moreover, $c \in \text{supp}(\mathbf{v})$ since $\mathbf{w}, \mathbf{v} \geq \mathbf{0}$ by assumption that $\mathbf{v}, \mathbf{w} \in \mathcal{P}(\psi)$. Thus $\mathbf{w} - \mathbf{v}$, a generator of $\mathcal{C}$, is parallel to a vector in $R_{\text{supp}(\mathbf{v})}$. $\square$

2.5. The construction of a short multivariate rational generating function

From the knowledge of triangulations of tangent cones of matroid polytopes, independence matroid polytopes, and polymatroids we will now recover short multivariate generating functions.

REMARK 38. Notice that formula (2.1) is of exponential size, even when the triangulation $\mathcal{T}$ only has polynomially many simplicial cones of maximal dimension. The reason is that, when the dimension n is allowed to vary, there are exponentially many intersecting proper faces in the set $\hat{\mathcal{T}}$. Therefore, we cannot use (2.1) to compute the multivariate rational generating function of $\mathcal{C}$ in polynomial time for varying dimension.

To obtain a shorter formula, we use the technique of *half-open exact decompositions* [KV08], which is a refinement of the method of “irrational” perturbations [BS07b, Köp07b]. We use the following result; see also Figure 2.5 and Figure 2.6.

LEMMA 39.

(a) *Let*

$$(2.5) \quad \sum_{i \in I_1} \varepsilon_i[\mathcal{C}_i] + \sum_{i \in I_2} \varepsilon_i[\mathcal{C}_i] = 0$$

be a linear identity (with rational coefficients ε_i) of indicator functions of cones $\mathcal{C}_i \subseteq \mathbb{R}^n$, where the cones $\mathcal{C}_i$ are full-dimensional for $i \in I_1$ and lower-dimensional for $i \in I_2$. Let each cone be given as

$$(2.6) \quad \mathcal{C}_i = \{ \mathbf{x} \in \mathbb{R}^n : \langle \mathbf{b}_{i,j}^*, \mathbf{x} \rangle \leq 0 \text{ for } j \in J_i \}.$$

Let $\mathbf{y} \in \mathbb{R}^n$ be a vector such that $\langle \mathbf{b}_{i,j}^*, \mathbf{y} \rangle \neq 0$ for all $i \in I_1 \cup I_2$, $j \in J_i$. For $i \in I_1$, we define the “half-open cone”

$$(2.7) \quad \tilde{\mathcal{C}}_i = \left\{ \mathbf{x} \in \mathbb{R}^d : \langle \mathbf{b}_{i,j}^*, \mathbf{x} \rangle \leq 0 \text{ for } j \in J_i \text{ with } \langle \mathbf{b}_{i,j}^*, \mathbf{y} \rangle < 0, \right. \\ \left. \langle \mathbf{b}_{i,j}^*, \mathbf{x} \rangle < 0 \text{ for } j \in J_i \text{ with } \langle \mathbf{b}_{i,j}^*, \mathbf{y} \rangle > 0 \right\}.$$

Then

$$(2.8) \quad \sum_{i \in I_1} \varepsilon_i[\tilde{\mathcal{C}}_i] = 0.$$




FIGURE 2.5. An identity, valid modulo lower-dimensional cones, corresponding to a polyhedral subdivision of a cone




FIGURE 2.6. The technique of half-open exact decomposition. The relative location of the vector $\mathbf{y}$ (represented by a dot) determines which defining inequalities are strict (broken lines) and which are weak (solid lines).

(b) *In particular, let*

$$(2.9) \quad [\mathcal{C}] = \sum_{\mathcal{C}_i \in \mathcal{T}} [\mathcal{C}_i] + \sum_{\mathcal{C}_i \in \hat{\mathcal{T}}} \varepsilon_i [\mathcal{C}_i]$$

be the identity corresponding to a triangulation of the cone $\mathcal{C}$, where $\mathcal{T}$ is the set of simplicial cones of maximal dimension and $\hat{\mathcal{T}}$ is the set of intersecting proper faces. Then there exists a polynomial-time algorithm to construct a vector $\mathbf{y} \in \mathbb{Q}^n$ such that the above construction yields the identity

$$(2.10) \quad [\mathcal{C}] = \sum_{\mathcal{C}_i \in \mathcal{T}} [\tilde{\mathcal{C}}_i],$$

which describes a partition of $\mathcal{C}$ into half-open cones of maximal dimension.

PROOF. Part (a) is a slightly less general form of Theorem 3 in [KV08]. Part (b) follows from the discussion in section 2 of [KV08]. $\square$

Since the cones in a triangulation $\mathcal{T}$ of all tangent cones $\mathcal{C}_{\mathcal{P}}(\mathbf{v})$ of our polytopes are unimodular by [Lemma 26](#), we can efficiently write the multivariate generating functions of their half-open counterparts.

LEMMA 40 (Lemma 9 in [\[KV08\]](#)). *Let $\tilde{\mathcal{C}} \subseteq \mathbb{R}^n$ be an N -dimensional half-open pointed simplicial affine cone with an integral apex $\mathbf{v} \in \mathbb{Z}^n$ and the ray description*

$$(2.11) \quad \tilde{\mathcal{C}} = \left\{ \mathbf{v} + \sum_{j=1}^N \lambda_j \mathbf{b}_j : \lambda_j \geq 0 \text{ for } j \in J_{\leq} \text{ and } \lambda_j > 0 \text{ for } j \in J_{<} \right\}$$

where $J_{\leq} \cup J_{<} = \{1, \dots, N\}$ and $\mathbf{b}_j \in \mathbb{Z}^n \setminus \{\mathbf{0}\}$. We further assume that $\tilde{\mathcal{C}}$ is unimodular, i.e., the vectors $\mathbf{b}_j$ form a basis of the lattice $(\mathbb{R}\mathbf{b}_1 + \dots + \mathbb{R}\mathbf{b}_N) \cap \mathbb{Z}^n$. Then the unique point in the fundamental parallelepiped of the half-open cone $\tilde{\mathcal{C}}$ is

$$(2.12) \quad \mathbf{a} = \mathbf{v} + \sum_{j \in J_{<}} \mathbf{b}_j,$$

and the generating function of $\mathcal{C}$ is given by

$$(2.13) \quad g_{\mathcal{C}}(\mathbf{z}) = \frac{\mathbf{z}^{\mathbf{a}}}{\prod_{j=1}^N (1 - \mathbf{z}^{\mathbf{b}_j})}.$$

Taking all results together, we obtain:

COROLLARY 41. *Let r be a fixed integer. There exist algorithms that, given*

- (a) *a matroid M on n elements, presented by an evaluation oracle for its rank function φ , which is bounded above by r , or*
- (b) *an evaluation oracle for an integral polymatroid rank function $\psi: 2^{[n]} \rightarrow \mathbb{N}$, which is bounded above by r ,*

compute in time polynomial in n vectors $\mathbf{a}_i \in \mathbb{Z}^n$, $\mathbf{b}_{i,j} \in \mathbb{Z}^n \setminus \{\mathbf{0}\}$, and $\mathbf{v}_i \in \mathbb{Z}^n$ for $i \in I$ (a polynomial-size index set) and $j = 1, \dots, N$, where $N \leq n$, such that the multivariate generating function of $\mathcal{P}(M)$, $\mathcal{P}^{\mathcal{I}}(M)$ and $\mathcal{P}(\psi)$, respectively, is the sum of rational functions

$$(2.14) \quad g_{\mathcal{P}}(\mathbf{z}) = \sum_{i \in I} \frac{\mathbf{z}^{\mathbf{a}_i}}{\prod_{j=1}^N (1 - \mathbf{z}^{\mathbf{b}_{i,j}})}$$

and the k -th dilation of the polytope has the multivariate rational generating function

$$(2.15) \quad g_{k\mathcal{P}}(\mathbf{z}) = \sum_{i \in I} \frac{\mathbf{z}^{\mathbf{a}_i + (k-1)\mathbf{v}_i}}{\prod_{j=1}^N (1 - \mathbf{z}^{\mathbf{b}_{i,j}})}.$$

PROOF. Lemma 23 implies that finding the multivariate generating function of $\mathcal{P}(M)$, $\mathcal{P}^{\mathcal{I}}(M)$ or $\mathcal{P}(\psi)$ can be reduced to finding the multivariate generating functions of their tangent cones. Moreover, $\mathcal{P}(M)$, $\mathcal{P}^{\mathcal{I}}(M)$ and $\mathcal{P}(\psi)$ have only polynomially in n many vertices as described in subsection 2.3 or Lemma 35. Enumerating their vertices can be done in polynomial time by Lemma 17 or 36.

Given a vertex $\mathbf{v}$ of $\mathcal{P}(M)$, $\mathcal{P}^{\mathcal{I}}(M)$ or $\mathcal{P}(\psi)$, its neighbors can be computed in polynomial time by Lemma 17 or 36. The tangent cone $\mathcal{C}(\mathbf{v})$ at $\mathbf{v}$ is generated by elements in R_A , where R_A is an elementary set for some $A \subseteq [n]$. See subsection 2.3 or Lemma 37. We also proved in Lemma 26 that every triangulation of $\mathcal{C}(\mathbf{v})$ generated by elements in R_A is unimodular and Lemma 28 states that any triangulation of $\mathcal{C}(\mathbf{v})$ has at most a polynomial in n number of top-dimensional simplices. Moreover, a triangulation of the cone $\mathcal{C}(\mathbf{v})$ can be computed in polynomial time by Lemma 31. Finally, using Lemmas 39 and 40 we can write the polynomial sized multivariate generating function of $\mathcal{C}(\mathbf{v})$ in polynomial time. Therefore we can write the multivariate generating function of $\mathcal{P}(M)$, $\mathcal{P}^{\mathcal{I}}(M)$ or $\mathcal{P}(\psi)$ in polynomial time. □

2.6. Polynomial-time specialization of rational generating functions in varying dimension

We now compute the Ehrhart polynomial $i(\mathcal{P}, k) = \#(k\mathcal{P} \cap \mathbb{Z}^n)$ from the multivariate rational generating function $g_{k\mathcal{P}}(\mathbf{z})$ of Corollary 41. This amounts to the problem of evaluating or *specializing* a rational generating function $g_{k\mathcal{P}}(\mathbf{z})$, depending on a parameter k , at the point $\mathbf{z} = \mathbf{1}$. This is a pole of each of its summands but a regular point (removable singularity) of the function itself. From now on we call this the *specialization problem*. We explain a very general procedure to solve it which we hope will allow future applications.

To this end, let the generating function of a polytope $\mathcal{P} \subseteq \mathbb{R}^n$ be given in the form

$$(2.16) \quad g_{\mathcal{P}}(\mathbf{z}) = \sum_{i \in I} \varepsilon_i \frac{\mathbf{z}^{\mathbf{a}_i}}{\prod_{j=1}^{s_i} (1 - \mathbf{z}^{\mathbf{b}_{i,j}})}$$

where $\varepsilon_i \in \{\pm 1\}$, $\mathbf{a}_i \in \mathbb{Z}^n$, and $\mathbf{b}_{ij} \in \mathbb{Z}^n \setminus \{\mathbf{0}\}$. Let $s = \max_{i \in I} s_i$ be the maximum number of binomials in the denominators. In general, if s is allowed to grow, more poles need to be considered for each summand, so the evaluation will need more computational effort.

In previous literature, the specialization problem has been considered, but not in sufficient generality for our purpose. In the original paper by (author?) [Bar94, Lemma 4.3], the dimension n is fixed, and each summand has exactly $s_i = n$ binomials in the denominator. The same restriction can be found in the survey by (author?) [BP99]. In the more general algorithmic theory of monomial substitutions developed by (author?) [BW03, Woo04], there is no assumption on the dimension n , but the number s of binomials in the denominators is fixed. The same restriction appears in the paper by (author?) [VW08, Lemma 2.15]. In a recent paper, (author?) [Bar06, section 5] gives a polynomial-time algorithm for the specialization problem for rational functions of the form

$$(2.17) \quad g(\mathbf{z}) = \sum_{i \in I} \varepsilon_i \frac{\mathbf{z}^{\mathbf{a}_i}}{\prod_{j=1}^s (1 - \mathbf{z}^{\mathbf{b}_{ij}})^{\gamma_{ij}}}$$

where the dimension n is fixed, the number s of different binomials in each denominator equals n , but the multiplicity γ_{ij} is varying.

We will show that the technique from (author?) [Bar06, section 5] can be implemented in a way such that we obtain a polynomial-time algorithm even for the case of a general formula (2.16), when the dimension and the number of binomials are allowed to grow.

THEOREM 42 (Polynomial-time specialization [DLHK08]). (a) *There exists an algorithm for computing the specialization of a rational function of the form*

$$(2.18) \quad g_{\mathcal{P}}(\mathbf{z}) = \sum_{i \in I} \varepsilon_i \frac{\mathbf{z}^{\mathbf{a}_i}}{\prod_{j=1}^{s_i} (1 - \mathbf{z}^{\mathbf{b}_{ij}})}$$

at its removable singularity $\mathbf{z} = \mathbf{1}$, which runs in time polynomial in the encoding size of its data $\varepsilon_i \in \mathbb{Q}$, $\mathbf{a}_i \in \mathbb{Z}^n$ for $i \in I$ and $\mathbf{b}_{ij} \in \mathbb{Z}^n$ for $i \in I, j = 1, \dots, s_i$, even when the dimension n and the numbers s_i of terms in the denominators are not fixed.

(b) *In particular, there exists a polynomial-time algorithm that, given data $\varepsilon_i \in \mathbb{Q}$, $\mathbf{a}_i \in \mathbb{Z}^n$ for $i \in I$ and $\mathbf{b}_{ij} \in \mathbb{Z}^n$ for $i \in I, j = 1, \dots, s_i$ describing a rational function in the form (2.18), computes a vector $\boldsymbol{\lambda} \in \mathbb{Q}^n$ with $\langle \boldsymbol{\lambda}, \mathbf{b}_{ij} \rangle \neq 0$ for all i, j and rational weights*

$w_{i,l}$ for $i \in I$ and $l = 0, \dots, s_i$. Then the number of integer points is given by

$$(2.19) \quad \#(\mathcal{P} \cap \mathbb{Z}^n) = \sum_{i \in I} \varepsilon_i \sum_{l=0}^{s_i} w_{i,l} \langle \boldsymbol{\lambda}, \mathbf{a}_i \rangle^l.$$

(c) Likewise, given a parametric rational function for the dilations of an integral polytope $\mathcal{P}$,

$$(2.20) \quad g_{k\mathcal{P}}(\mathbf{z}) = \sum_{i \in I} \varepsilon_i \frac{\mathbf{z}^{\mathbf{a}_i + (k-1)\mathbf{v}_i}}{\prod_{j=1}^d (1 - \mathbf{z}^{\mathbf{b}_{i,j}})},$$

the Ehrhart polynomial $i(\mathcal{P}, k) = \#(k\mathcal{P} \cap \mathbb{Z}^n)$ is given by the explicit formula

$$(2.21) \quad i(\mathcal{P}, k) = \sum_{m=0}^M \left(\sum_{i \in I} \varepsilon_i \langle \boldsymbol{\lambda}, \mathbf{v}_i \rangle^m \sum_{l=m}^{s_i} \binom{l}{m} w_{i,l} \langle \boldsymbol{\lambda}, \mathbf{a}_i - \mathbf{v}_i \rangle^{l-m} \right) k^m,$$

where $M = \min\{s, \dim \mathcal{P}\}$.

PROOF OF THEOREM 22. Corollary 41 and Theorem 42 imply Theorem (Theorem 22) directly. □

The remainder of this section contains the proof of Theorem 42. We follow [BP99] and recall the definition of Todd polynomials. We will prove that they can be efficiently evaluated in rational arithmetic.

DEFINITION 43. We consider the function

$$H(x, \xi_1, \dots, \xi_s) = \prod_{i=1}^s \frac{x \xi_i}{1 - \exp\{-x \xi_i\}},$$

a function that is analytic in a neighborhood of $\mathbf{0}$. The m -th (s -variate) *Todd polynomial* is the coefficient of x^m in the Taylor expansion

$$H(x, \xi_1, \dots, \xi_s) = \sum_{m=0}^{\infty} \text{td}_m(\xi_1, \dots, \xi_s) x^m.$$

We remark that, when the numbers s and m are allowed to vary, the Todd polynomials have an exponential number of monomials.

THEOREM 44 ([DLHK08]). The Todd polynomial $\text{td}_m(\xi_1, \dots, \xi_s)$ can be evaluated for given rational data $\xi_1, \dots, \xi_s$ in time polynomial in s , m , and the encoding length of $\xi_1, \dots, \xi_s$.

The proof makes use of the following lemma.

LEMMA 45 ([DLHK08]). *The function $h(x) = x/(1 - e^{-x})$ is a function that is analytic in a neighborhood of 0. Its Taylor series about $x = 0$ is of the form*

$$(2.22) \quad h(x) = \sum_{n=0}^{\infty} b_n x^n \quad \text{where} \quad b_n = \frac{1}{n! (n+1)!} c_n$$

with integer coefficients c_n that have a binary encoding length of $O(n^2 \log n)$. The coefficients c_n can be computed from the recursion

$$(2.23) \quad \begin{aligned} c_0 &= 1 \\ c_n &= \sum_{j=1}^n (-1)^{j+1} \binom{n+1}{j+1} \frac{n!}{(n-j+1)!} c_{n-j} \quad \text{for } n = 1, 2, \dots \end{aligned}$$

PROOF. The reciprocal function $h^{-1}(x) = (1 - e^{-x})/x$ has the Taylor series

$$h^{-1}(x) = \sum_{i=0}^{\infty} a_n x^n \quad \text{with} \quad a_n = \frac{(-1)^n}{(n+1)!}.$$

Using the identity $h^{-1}(x)h(x) = (\sum_{n=0}^{\infty} a_n x^n)(\sum_{n=0}^{\infty} b_n x^n) = 1$, we obtain the recursion

$$(2.24) \quad \begin{aligned} b_0 &= \frac{1}{a_0} = 1 \\ b_n &= -(a_1 b_{n-1} + a_2 b_{n-2} + \dots + a_n b_0) \quad \text{for } n = 1, 2, \dots \end{aligned}$$

We prove (2.22) inductively. Clearly $b_0 = c_0 = 1$. For $n = 1, 2, \dots$, we have

$$\begin{aligned} c_n &= n! (n+1)! b_n \\ &= -n! (n+1)! (a_1 b_{n-1} + a_2 b_{n-2} + \dots + a_n b_0) \\ &= n! (n+1)! \sum_{j=1}^n \frac{(-1)^{j+1}}{(j+1)!} \cdot \frac{1}{(n-j)! (n-j+1)!} c_{n-j} \\ &= \sum_{j=1}^n (-1)^{j+1} \frac{(n+1)!}{(j+1)! (n-j)!} \cdot \frac{n!}{(n-j+1)!} c_{n-j}. \end{aligned}$$

Thus we obtain the recursion formula (2.23), which also shows that all c_n are integers. A rough estimate shows that

$$|c_n| \leq n(n+1)! n! \quad |c_{n-1}| \leq ((n+1)!)^2 |c_{n-1}|,$$

thus $|c_n| \leq ((n+1)!)^{2n}$, so c_n has a binary encoding length of $O(n^2 \log n)$. □

PROOF OF THEOREM 44. By definition, we have

$$H(x, \xi_1, \dots, \xi_s) = \sum_{m=0}^{\infty} \text{td}_m(\xi_1, \dots, \xi_s) x^m = \prod_{j=1}^s h(x\xi_j).$$

From Lemma 45 we have

$$(2.25) \quad h(x\xi_j) = \sum_{n=0}^m \beta_{j,n} x^n + o(x^m) \quad \text{where} \quad \beta_{j,n} = \frac{\xi_j^n}{n!(n+1)!} c_n$$

with integers c_n given by the recursion (2.23). Thus we can evaluate $\text{td}_m(\xi_1, \dots, \xi_s)$ by summing over all the possible compositions $n_1 + \dots + n_s = m$ of the order m from the orders n_j of the factors:

$$(2.26) \quad \text{td}_m(\xi_1, \dots, \xi_s) = \sum_{\substack{(n_1, \dots, n_s) \in \mathbb{Z}_+^s \\ n_1 + \dots + n_s = m}} \beta_{1,n_1} \dots \beta_{s,n_s}$$

We remark that the length of the above sum is equal to the number of compositions of m into s non-negative parts,

$$\begin{aligned} C'_s(m) &= \binom{m+s-1}{s-1} \\ &= \frac{(m+s-1)(m+s-2) \dots (m+s-(s-1))}{(s-1)(s-2) \dots 2 \cdot 1} \\ &= \Omega \left(\left(1 + \frac{m}{s-1}\right)^s \right), \end{aligned}$$

which is *exponential* in s (whenever $m \geq s$). Thus we cannot evaluate the formula (2.26) efficiently when s is allowed to grow.

However, we show that we can evaluate $\text{td}_m(\xi_1, \dots, \xi_s)$ more efficiently. To this end, we multiply up the s truncated Taylor series (2.25), one factor at a time, truncating after order m . Let us denote

$$\begin{aligned} H_1(x) &= h(x\xi_1) \\ H_2(x) &= H_1(x) \cdot h(x\xi_2) \\ &\vdots \\ H_s(x) &= H_{s-1}(x) \cdot h(x\xi_s) = H(x, \xi_1, \dots, \xi_s). \end{aligned}$$

Each multiplication can be implemented in $O(m^2)$ elementary rational operations. We finally show that all numbers appearing in the calculations have polynomial encoding size. Let Ξ be the largest binary encoding size of any of the rational numbers $\xi_1, \dots, \xi_s$. Then every $\beta_{j,n}$ given by (2.25) has a binary encoding size $O(\Xi n^5 \log^3 n)$. Let $H_j(x)$ have the truncated Taylor series $\sum_{n=0}^m \alpha_{j,n} x^n + o(x^m)$ and let A_j denote the largest binary encoding length of any $\alpha_{j,n}$ for $n \leq m$. Then

$$H_{j+1}(x) = \sum_{n=0}^m \alpha_{j+1,n} x^n + o(x^m) \quad \text{with} \quad \alpha_{j+1,n} = \sum_{l=0}^n \alpha_{j,l} \beta_{j,n-l}.$$

Thus the binary encoding size of $\alpha_{j+1,n}$ (for $n \leq m$) is bounded by $A_j + O(\Xi m^5 \log^3 m)$. Thus, after s multiplication steps, the encoding size of the coefficients is bounded by $O(s \Xi m^5 \log^3 m)$, a polynomial quantity. $\square$

PROOF OF THEOREM 42. *Parts (a) and (b).* We recall the technique of (author?) [Bar94, Lemma 4.3], refined by (author?) [Bar06, section 5].

We first construct a rational vector $\lambda \in \mathbb{Z}^n$ such that $\langle \lambda, \mathbf{b}_{ij} \rangle \neq 0$ for all i, j . One such construction is to consider the *moment curve* $\lambda(\xi) = (1, \xi, \xi^2, \dots, \xi^{n-1}) \in \mathbb{R}^n$. For each exponent vector $\mathbf{b}_{ij}$ occurring in a denominator of (2.16), the function $f_{ij}: \xi \mapsto \langle \lambda(\xi), \mathbf{b}_{ij} \rangle$ is a polynomial function of degree at most $n - 1$. Since $\mathbf{b}_{ij} \neq \mathbf{0}$, the function f_{ij} is not identically zero. Hence f_{ij} has at most $n - 1$ zeros. By evaluating all functions f_{ij} for $i \in I$ and $j = 1, \dots, s_i$ at $M = (n - 1)s|I| + 1$ different values for ξ , for instance at the integers $\xi = 0, \dots, M$, we can find one $\xi = \bar{\xi}$ that is not a zero of any f_{ij} . Clearly this search can be implemented in polynomial time, even when the dimension n and the number s of terms in the denominators are not fixed. We set $\lambda = \lambda(\bar{\xi})$.

For $\tau > 0$, let us consider the points $\mathbf{z}_\tau = \mathbf{e}^{\tau \lambda} = (\exp\{\tau \lambda_1\}, \dots, \exp\{\tau \lambda_n\})$. We have

$$\mathbf{z}_\tau^{\mathbf{b}_{ij}} = \prod_{l=1}^n \exp\{\tau \lambda_l b_{ijl}\} = \exp\{\tau \langle \lambda, \mathbf{b}_{ij} \rangle\};$$

since $\langle \lambda, \mathbf{b}_{ij} \rangle \neq 0$ for all i, j , all the denominators $1 - \mathbf{z}_\tau^{\mathbf{b}_{ij}}$ are nonzero. Hence for every $\tau > 0$, the point $\mathbf{z}_\tau$ is a regular point not only of $g(\mathbf{z})$ but also of the individual summands of (2.16). We have

$$g(\mathbf{1}) = \lim_{\tau \rightarrow 0^+} \sum_{i \in I} \varepsilon_i \frac{\mathbf{z}_\tau^{\mathbf{a}_i}}{\prod_{j=1}^{s_i} (1 - \mathbf{z}_\tau^{\mathbf{b}_{ij}})}$$

$$\begin{aligned}
 &= \lim_{\tau \rightarrow 0^+} \sum_{i \in I} \varepsilon_i \frac{\exp\{\tau \langle \boldsymbol{\lambda}, \mathbf{a}_i \rangle\}}{\prod_{j=1}^{s_i} (1 - \exp\{\tau \langle \boldsymbol{\lambda}, \mathbf{b}_{ij} \rangle\})} \\
 &= \lim_{\tau \rightarrow 0^+} \sum_{i \in I} \varepsilon_i \tau^{-s_i} \exp\{\tau \langle \boldsymbol{\lambda}, \mathbf{a}_i \rangle\} \prod_{j=1}^{s_i} \frac{\tau}{1 - \exp\{\tau \langle \boldsymbol{\lambda}, \mathbf{b}_{ij} \rangle\}} \\
 &= \lim_{\tau \rightarrow 0^+} \sum_{i \in I} \varepsilon_i \tau^{-s_i} \exp\{\tau \langle \boldsymbol{\lambda}, \mathbf{a}_i \rangle\} \prod_{j=1}^{s_i} \frac{-1}{\langle \boldsymbol{\lambda}, \mathbf{b}_{ij} \rangle} h(-\tau \langle \boldsymbol{\lambda}, \mathbf{b}_{ij} \rangle) \\
 &= \lim_{\tau \rightarrow 0^+} \sum_{i \in I} \varepsilon_i \frac{(-1)^{s_i}}{\prod_{j=1}^{s_i} \langle \boldsymbol{\lambda}, \mathbf{b}_{ij} \rangle} \tau^{-s_i} \exp\{\tau \langle \boldsymbol{\lambda}, \mathbf{a}_i \rangle\} H(\tau, -\langle \boldsymbol{\lambda}, \mathbf{b}_{i1} \rangle, \dots, -\langle \boldsymbol{\lambda}, \mathbf{b}_{is_i} \rangle)
 \end{aligned}$$

where $H(x, \xi_1, \dots, \xi_{s_i})$ is the function from [Definition 43](#). We will compute the limit by finding the constant term of the Laurent expansion of each summand about $\tau = 0$. Now the function $\tau \mapsto \exp\{\tau \langle \boldsymbol{\lambda}, \mathbf{a}_i \rangle\}$ is holomorphic and has the Taylor series

$$(2.27) \quad \exp\{\tau \langle \boldsymbol{\lambda}, \mathbf{a}_i \rangle\} = \sum_{l=0}^{s_i} \alpha_{i,l} \tau^l + o(\tau^{s_i}) \quad \text{where} \quad \alpha_{i,l} = \frac{\langle \boldsymbol{\lambda}, \mathbf{a}_i \rangle^l}{l!},$$

and $H(\tau, \xi_1, \dots, \xi_{s_i})$ has the Taylor series

$$H(\tau, \xi_1, \dots, \xi_s) = \sum_{m=0}^{s_i} \text{td}_m(\xi_1, \dots, \xi_s) \tau^m + o(\tau^{s_i}).$$

Because of the factor τ^{-s_i} , which gives rise to a pole of order s_i in the summand, we can compute the constant term of the Laurent expansion by summing over all the possible compositions $s_i = l + (s_i - l)$ of the order s_i :

$$(2.28) \quad g(\mathbf{1}) = \sum_{i \in I} \varepsilon_i \frac{(-1)^{s_i}}{\prod_{j=1}^{s_i} \langle \boldsymbol{\lambda}, \mathbf{b}_{ij} \rangle} \sum_{l=0}^{s_i} \frac{\langle \boldsymbol{\lambda}, \mathbf{a}_i \rangle^l}{l!} \text{td}_{s_i-l}(-\langle \boldsymbol{\lambda}, \mathbf{b}_{i1} \rangle, \dots, -\langle \boldsymbol{\lambda}, \mathbf{b}_{is_i} \rangle).$$

We use the notation

$$w_{i,l} = (-1)^{s_i} \frac{\text{td}_{s_i-l}(-\langle \boldsymbol{\lambda}, \mathbf{b}_{i1} \rangle, \dots, -\langle \boldsymbol{\lambda}, \mathbf{b}_{is_i} \rangle)}{l! \cdot \langle \boldsymbol{\lambda}, \mathbf{b}_{i1} \rangle \cdots \langle \boldsymbol{\lambda}, \mathbf{b}_{is_i} \rangle} \quad \text{for } i \in I \text{ and } l = 0, \dots, s_i;$$

these rational numbers can be computed in polynomial time using [Theorem 44](#). We now obtain the formula of the claim,

$$g(\mathbf{1}) = \sum_{i \in I} \varepsilon_i \sum_{l=0}^{s_i} w_{i,l} \langle \boldsymbol{\lambda}, \mathbf{a}_i \rangle^l.$$

Part (c). Applying the same technique to the parametric rational function (2.20), we obtain

$$\begin{aligned}
 \#(k\mathcal{P} \cap \mathbb{Z}^n) &= g_{k\mathcal{P}}(\mathbf{1}) \\
 &= \sum_{i \in I} \varepsilon_i \sum_{l=0}^{s_i} w_{i,l} \langle \boldsymbol{\lambda}, \mathbf{a}_i + (k-1)\mathbf{v}_i \rangle^l, \\
 &= \sum_{i \in I} \varepsilon_i \sum_{l=0}^{s_i} w_{i,l} \sum_{m=0}^l \binom{l}{m} \langle \boldsymbol{\lambda}, \mathbf{a}_i - \mathbf{v}_i \rangle^{l-m} k^m \langle \boldsymbol{\lambda}, \mathbf{v}_i \rangle^m \\
 &= \sum_{m=0}^s \left(\sum_{i \in I} \varepsilon_i \langle \boldsymbol{\lambda}, \mathbf{v}_i \rangle^m \sum_{l=m}^{s_i} \binom{l}{m} w_{i,l} \langle \boldsymbol{\lambda}, \mathbf{a}_i - \mathbf{v}_i \rangle^{l-m} \right) k^m,
 \end{aligned}$$

an explicit formula for the Ehrhart polynomial. We remark that, since the Ehrhart polynomial is of degree equal to the dimension of $\mathcal{P}$, all coefficients of k^m for $m > \dim \mathcal{P}$ must vanish. Thus we obtain the formula of the claim, where we sum only up to $\min\{s, \dim \mathcal{P}\}$ instead of s . □

CHAPTER 3

Results on the Algebraic Combinatorics of Matroid Polytopes

3.1. Algebraic Properties of h^* -vectors and Ehrhart polynomials of Matroid Polytopes

Here we investigate algebraic properties of the Ehrhart functions of matroid polytopes: The *Ehrhart series* of a polytope $\mathcal{P}$ is the infinite series $\sum_{k=0}^{\infty} i(\mathcal{P}, k)t^k$. We recall the following classic result about Ehrhart series (see e.g., [Hib92, Sta96]). Let $\mathcal{P} \subseteq \mathbb{R}^n$ be an integral convex polytope of dimension d . Then it is known that its Ehrhart series is a rational function of the form

$$(3.1) \quad \sum_{k=0}^{\infty} i(\mathcal{P}, k)t^k = \frac{h_0^* + h_1^*t + \cdots + h_{d-1}^*t^{d-1} + h_d^*t^d}{(1-t)^{d+1}}.$$

The numerator is often called the h^* -*polynomial* of $\mathcal{P}$ (some authors also call it the *Ehrhart h -polynomial*), and we define the coefficients of the polynomial in the numerator of Lemma 3.1, $h_0^* + h_1^*t + \cdots + h_{d-1}^*t^{d-1} + h_d^*t^d$, as the h^* -*vector* of $\mathcal{P}$, which we write as $\mathbf{h}^*(\mathcal{P}) := (h_0^*, h_1^*, \dots, h_{d-1}^*, h_d^*)$.

A vector $(c_0, \dots, c_d)$ is *unimodal* if there exists an index p , $0 \leq p \leq d$, such that $c_{i-1} \leq c_i$ for $i \leq p$ and $c_j \geq c_{j+1}$ for $j \geq p$. Due to its algebraic implications, several authors have studied the unimodality of h^* -vectors (see [Hib92] and [Sta96] and references therein).

Suppose, as before, that $\mathcal{P} \subseteq \mathbb{R}^n$, and each vertex of $\mathcal{P}$ has integral (or rational) vertices. Let $Y_1, Y_2, \dots, Y_n$ and T be indeterminates over a field K . Letting $q \geq 1$ we define $A(\mathcal{P})_q$ as the vector space over K which is spanned by the monomials $Y_1^{\alpha_1}, Y_2^{\alpha_2}, \dots, Y_n^{\alpha_n} T^q$ such that $(\alpha_1, \alpha_2, \dots, \alpha_n) \in q\mathcal{P} \cap \mathbb{Z}^n$. Since $\mathcal{P}$ is convex it follows that $A(\mathcal{P})_q A(\mathcal{P})_p \subseteq A_{q+p}(\mathcal{P})$ for all p, q , and thus the *Ehrhart ring* of $\mathcal{P}$, $A(\mathcal{P}) = \bigoplus_{q=0}^{\infty} A(\mathcal{P})_q$, is a graded algebra [Hib92, Sta96].

It is well-known that if the Ehrhart ring of an integral polytope $\mathcal{P}$, $A(\mathcal{P})$, is Gorenstein, then $\mathbf{h}^*(\mathcal{P})$ is unimodal, and symmetric [Hib92, Sta96]. Nevertheless, the vector $\mathbf{h}^*(\mathcal{P})$ can be unimodal even when the Ehrhart ring $A(\mathcal{P})$ is not Gorenstein. For matroid polytopes, their Ehrhart ring is indeed often not Gorenstein. For instance, De Negri and Hibi [DNH97] prove explicitly when the Ehrhart ring of a uniform matroid polytope is Gorenstein or not. Two fascinating facts, uncovered through experimentation, are that all h^* -vectors seen thus far are unimodal, even for the cases when their Ehrhart rings are not Gorenstein. In addition, when we computed the explicit Ehrhart polynomials of matroid polytopes we observe their coefficients are always positive. We conjecture:

CONJECTURE 46. *Let $\mathcal{P}(M)$ be the matroid polytope of a matroid M .*

- (A) *The h^* -vector of $\mathcal{P}(M)$ is unimodal.*
- (B) *The coefficients of the Ehrhart polynomial of $\mathcal{P}(M)$ are positive.*

We have proved both parts of this conjecture in many instances. A class of matroids that we considered are the uniform matroids; recall that the *uniform matroid* on n elements of rank r is the collection of all r -subsets of n . Using computers, we were able to verify Conjecture 46 for all uniform matroids up to 75 elements as well as for a wide variety of non-uniform matroids which are collected at [Haw09]. We include here this information just for the 28 famous matroids presented in [Oxl92]. Results in [Kat05], with some additional careful calculations, imply that Conjecture 46 is true for all rank 2 uniform matroids. Regarding part (A) of the conjecture we were also able to prove *partial unimodality* for uniform matroids of rank 3. Concretely we obtained:

THEOREM 47.

- (1) *Conjecture 46 is true for all uniform matroids up to 75 elements and all uniform matroids of rank 2. It is also true for all matroids listed in [Haw09].*
- (2) *Let $\mathcal{P}(U^{3,n})$ be the matroid polytope of a uniform matroid of rank 3 on n elements, and let I be a non-negative integer. Then there exists $n(I) \in \mathbb{N}$ such that for all $n \geq n(I)$ the h^* -vector of $\mathcal{P}(U^{3,n})$, $(h_0^*, \dots, h_n^*)$, is non-decreasing from index 0 to I . That is, $h_0^* \leq h_1^* \leq \dots \leq h_I^*$.*

Using the programs `cdd+` [Fuk06], `LattE` [DLHH⁺05] and `LattE macchiato` [Köp07a] we explored patterns for the Ehrhart polynomials of matroid polytopes. Since previous authors proposed other invariants of a matroid (e.g., the Tutte polynomials and the invariants of [Spe06, BJR06]) we wished to know how well does the Ehrhart polynomial distinguish non-isomorphic matroids. It is natural to compare it with other known invariants. Some straightforward properties are immediately evident. For example, the Ehrhart polynomial of a matroid and that of its dual are equal. Also the Ehrhart polynomial of a direct sum of matroids is the product of their Ehrhart polynomials.

TABLE 3.1. Coefficients of the Ehrhart polynomials of selected matroids in [BJR06, Spe06]

	Ehrhart Polynomial
Speyer1	$1, \frac{21}{5}, \frac{343}{45}, \frac{63}{8}, \frac{91}{18}, \frac{77}{40}, \frac{29}{90}$
Speyer2	$1, \frac{135}{28}, \frac{3691}{360}, \frac{1511}{120}, \frac{88}{9}, \frac{39}{8}, \frac{529}{360}, \frac{89}{420}$
BJR1	$1, \frac{109}{30}, \frac{23}{4}, \frac{59}{12}, \frac{9}{4}, \frac{9}{20}$
BJR2	$1, \frac{211}{60}, \frac{125}{24}, \frac{33}{8}, \frac{43}{24}, \frac{43}{120}$
BJR3	$1, \frac{83}{20}, \frac{2783}{360}, \frac{199}{24}, \frac{391}{72}, \frac{247}{120}, \frac{61}{180}$
BJR4	$1, \frac{25}{6}, \frac{353}{45}, \frac{101}{12}, \frac{193}{36}, \frac{23}{12}, \frac{53}{180}$

We call the last two matroids in Figure 2 in [Spe06] *Speyer1* and *Speyer2* and the matroids of Figure 2 in [BJR06] *BJR1*, *BJR2*, *BJR3*, and *BJR4*, and list their Ehrhart polynomials in Table 3.1. We note that *BJR3* and *BJR4* have the same Tutte polynomial, yet their Ehrhart polynomials are different. This proves that the Ehrhart polynomial cannot be computed using deletion and contractions, as is the case for the Tutte polynomial. Examples *BJR1* and *BJR2* show that the Ehrhart polynomial of a matroid may help to distinguish non-isomorphic matroids: These two matroids are not isomorphic yet they have the same Tutte polynomials and the same quasi-symmetric function studied in [BJR06]. Although they share some properties, there does not seem to be an obvious relation to Speyer's univariate polynomials introduced in [Spe06]; examples *Speyer1* and *Speyer2* show they are relatively prime with their corresponding Ehrhart polynomials.

Our experiments included, among others, many examples coming from small graphical matroids, random realizable matroids over fields of small positive characteristic, and the classical examples listed in the Appendix of [Oxl92] for which we list the results in Table

3.2. For a comprehensive list of all our calculations visit [Haw09]. Soon it became evident that all instances verified both parts of our **Conjecture 46**.

TABLE 3.2. Coefficients of the Ehrhart polynomials and h^* -vectors of selected matroids in [Oxl92]

	n	r	h^* -vector	Ehrhart Polynomial
K_4	6	3	1, 10, 20, 10, 1	$1, \frac{107}{30}, \frac{21}{4}, \frac{49}{12}, \frac{7}{4}, \frac{7}{20}$
W^3	6	3	1, 11, 24, 11, 10	$1, \frac{18}{5}, \frac{11}{2}, \frac{9}{2}, 2, \frac{2}{5}$
Q_6	6	3	1, 12, 28, 12, 1	$1, \frac{109}{30}, \frac{23}{4}, \frac{59}{12}, \frac{9}{4}, \frac{9}{20}$
P_6	6	3	1, 13, 32, 13, 1	$1, \frac{11}{3}, 6, \frac{16}{3}, \frac{5}{2}, \frac{1}{2}$
R_6	6	3	1, 12, 28, 12, 1	$1, \frac{109}{30}, \frac{23}{4}, \frac{59}{12}, \frac{9}{4}, \frac{9}{20}$
F_7	7	3	21, 98, 91, 21, 1	$1, \frac{21}{5}, \frac{343}{45}, \frac{63}{8}, \frac{91}{18}, \frac{77}{40}, \frac{29}{90}$
F_7^-	7	3	21, 101, 97, 22, 1	$1, \frac{253}{60}, \frac{2809}{360}, \frac{33}{4}, \frac{193}{36}, \frac{61}{30}, \frac{121}{360}$
P^7	7	3	21, 104, 103, 23, 1	$1, \frac{127}{30}, \frac{479}{60}, \frac{69}{8}, \frac{17}{3}, \frac{257}{120}, \frac{7}{20}$
$AG(3, 2)$	8	4	1, 62, 561, 1014, 449, 48, 1	$1, \frac{209}{42}, \frac{1981}{180}, \frac{881}{60}, \frac{119}{9}, \frac{95}{12}, \frac{499}{180}, \frac{89}{210}$
$AG'(3, 2)$	8	4	1, 62, 562, 1023, 458, 49, 1	$1, \frac{299}{60}, \frac{4007}{360}, \frac{5401}{360}, \frac{122}{9}, \frac{2911}{360}, \frac{1013}{360}, \frac{77}{420}$
R_8	8	4	1, 62, 563, 1032, 467, 50, 1	$1, \frac{524}{105}, \frac{1013}{90}, \frac{1379}{90}, \frac{125}{9}, \frac{743}{90}, \frac{257}{90}, \frac{136}{315}$
F_8	8	4	1, 62, 563, 1032, 467, 50, 1	$1, \frac{524}{105}, \frac{1013}{90}, \frac{1379}{90}, \frac{125}{9}, \frac{743}{90}, \frac{257}{90}, \frac{136}{315}$
Q_8	8	4	1, 62, 564, 1041, 476, 51, 1	$1, \frac{2099}{420}, \frac{4097}{360}, \frac{1877}{120}, \frac{128}{9}, \frac{337}{40}, \frac{1043}{360}, \frac{61}{140}$
S_8	8	4	1, 44, 337, 612, 305, 40, 1	$1, \frac{1021}{210}, \frac{377}{36}, \frac{475}{36}, \frac{193}{18}, \frac{511}{90}, \frac{65}{36}, \frac{67}{252}$
V_8	8	4	1, 62, 570, 1095, 530, 57, 1	$1, \frac{2117}{420}, \frac{4367}{360}, \frac{2107}{120}, \frac{146}{9}, \frac{1133}{120}, \frac{1133}{360}, \frac{193}{420}$
T_8	8	4	1, 62, 564, 1041, 476, 51, 1	$1, \frac{2099}{420}, \frac{4097}{360}, \frac{1877}{120}, \frac{128}{9}, \frac{337}{40}, \frac{1043}{360}, \frac{61}{140}$
V_8^+	8	4	1, 62, 569, 1086, 521, 56, 1	$1, \frac{151}{30}, \frac{2161}{180}, \frac{3103}{180}, \frac{143}{9}, \frac{1669}{180}, \frac{559}{180}, \frac{41}{90}$
L_8	8	4	1, 62, 567, 1068, 503, 54, 1	$1, \frac{527}{105}, \frac{529}{45}, \frac{83}{5}, \frac{137}{9}, \frac{134}{15}, \frac{136}{45}, \frac{47}{105}$
J	8	4	1, 44, 339, 630, 323, 42, 1	$1, \frac{512}{105}, \frac{193}{18}, \frac{83}{6}, \frac{205}{18}, \frac{361}{60}, \frac{17}{9}, \frac{23}{84}$
P_8	8	4	1, 62, 565, 1050, 485, 52, 1	$1, \frac{1051}{210}, \frac{2071}{180}, \frac{2873}{180}, \frac{131}{9}, \frac{1547}{180}, \frac{529}{180}, \frac{277}{630}$
W_4	8	4	1, 38, 262, 475, 254, 37, 1	$1, \frac{135}{28}, \frac{3691}{360}, \frac{1511}{120}, \frac{88}{9}, \frac{39}{8}, \frac{529}{360}, \frac{89}{420}$
W^4	8	4	1, 38, 263, 484, 263, 38, 1	$1, \frac{169}{35}, \frac{467}{45}, \frac{581}{45}, \frac{91}{9}, \frac{227}{45}, \frac{68}{45}, \frac{68}{315}$
$K_{3,3}$	9	5	78, 1116, 3492, 3237, 927, 72, 1	$1, \frac{307}{56}, \frac{137141}{10080}, \frac{3223}{160}, \frac{37807}{1920}, \frac{211}{16}, \frac{5743}{960}, \frac{1889}{1120}, \frac{8923}{40320}$
$AG(2, 3)$	9	3	1, 147, 1230, 1885, 714, 63, 1	$1, \frac{1453}{280}, \frac{41749}{3360}, \frac{581}{32}, \frac{34069}{1920}, \frac{927}{80}, \frac{4541}{960}, \frac{239}{224}, \frac{449}{4480}$
Pappus	9	3	1, 147, 1230, 1915, 744, 66, 1	$1, \frac{729}{140}, \frac{3573}{280}, \frac{381}{20}, \frac{1499}{80}, \frac{243}{20}, \frac{49}{10}, \frac{153}{140}, \frac{57}{560}$
Non-Pappus	9	3	1, 147, 1230, 1925, 754, 67, 1	$1, \frac{4379}{840}, \frac{25951}{2016}, \frac{9287}{480}, \frac{21967}{1152}, \frac{987}{80}, \frac{2855}{576}, \frac{3701}{3360}, \frac{275}{2688}$
$Q_3(GF(3)^*)$	9	3	1, 147, 1098, 1638, 632, 59, 1	$1, \frac{433}{84}, \frac{3079}{252}, \frac{4193}{240}, \frac{5947}{360}, \frac{167}{16}, \frac{601}{144}, \frac{787}{840}, \frac{149}{1680}$
R_9	9	3	1, 147, 1142, 1717, 656, 60, 1	$1, \frac{723}{140}, \frac{49}{4}, \frac{88}{5}, \frac{24217}{1440}, \frac{1291}{120}, \frac{625}{144}, \frac{821}{840}, \frac{133}{1440}$

By far the most comprehensive study we made was for the family of uniform matroids. In this case we based our computations on the theory of Veronese algebras as developed by M. Katzman in [Kat05]. There, Katzman gives an explicit equation for the h^* -vector of uniform matroid polytopes (again, using the language of Veronese algebras). For this family we were able to verify computationally the conjecture is true for all uniform matroids up to 75 elements and to prove partial unimodality as explained in the introduction.

LEMMA 48. *The coefficients of the Ehrhart polynomial of the matroid polytope of the uniform matroid $U^{2,n}$ are positive.*

PROOF. We begin with the expression in Corollary 2.2 in [Kat05] which explicitly gives the Ehrhart polynomial of $\mathcal{P}(U^{r,n})$ as

$$(3.2) \quad i(\mathcal{P}(U^{r,n}), k) = \sum_{s=0}^{r-1} (-1)^s \binom{n}{s} \binom{k(r-s) - s + n - 1}{n-1}.$$

Letting $r = 2$, Equation (3.2) becomes

$$\frac{(2k+n-1)(2k+n-2)\cdots(2k+1)}{(n-1)!} - n \frac{(k+n-2)(k+n-3)\cdots(k)}{(n-1)!}$$

We next consider the coefficient of k^{n-p-1} for $0 \leq p \leq n-1$, which can be written as

$$\begin{aligned} & 2^{n-p-1} \sum_{\substack{1 \leq j_1 < \cdots < j_p \leq n-1 \\ j_q \in \mathbb{Z}}} \frac{1}{(n-j_1)\cdots(n-j_p)} \\ & - \frac{n}{n-1} \sum_{\substack{1 \leq j_1 < \cdots < j_{p-1} \leq n-2 \\ j_q \in \mathbb{Z}}} \frac{1}{(n-j_1)\cdots(n-j_{p-1})} \\ & = 2^{n-p-1} \sum_{\substack{1 \leq j_1 < \cdots < j_{p-1} \leq n-2 \\ j_q \in \mathbb{Z}}} \left(\frac{1}{(n-j_1)\cdots(n-j_{p-1})} \sum_{j_p=1+j_{p-1}}^{n-1} \frac{1}{n-j_p} \right) \\ & - \frac{n}{n-1} \sum_{\substack{1 \leq j_1 < \cdots < j_{p-1} \leq n-2 \\ j_q \in \mathbb{Z}}} \frac{1}{(n-j_1)\cdots(n-j_{p-1})} \\ (3.3) \quad & = \sum_{\substack{1 \leq j_1 < \cdots < j_{p-1} \leq n-2 \\ j_q \in \mathbb{Z}}} \left(\frac{1}{(n-j_1)\cdots(n-j_{p-1})} \left[2^{n-p-1} \sum_{j_p=1+j_{p-1}}^{n-1} \frac{1}{n-j_p} - \frac{n}{n-1} \right] \right). \end{aligned}$$

It is known that the constant in any Ehrhart polynomial is 1 [Sta96], thus we only need to show that Equation (3.3) is positive for $0 \leq p \leq n-2$. It is sufficient to show that the square-bracketed term of (3.3),

$$(3.4) \quad 2^{n-p-1} \sum_{j_p=1+j_{p-1}}^{n-1} \frac{1}{n-j_p} - \left(1 + \frac{1}{n-1} \right),$$

is positive for $0 \leq p \leq n-2$ and all $1 \leq j_1 \leq \cdots \leq j_{p-1} \leq n-2$. We can see that $\sum_{j_p=1+j_{p-1}}^{n-1} \frac{1}{n-j_p} \geq 1$. Moreover, since $0 \leq p \leq n-2$ then $2^{n-p-1} \geq 2$ and $1 + \frac{1}{n-1} \leq 2$ since $n \geq 2$, proving the result. $\square$

To present our results about the h^* -vector we begin explaining the details with the following numbers introduced in [Kat05], which we refer to as the *Katzman coefficients*:

DEFINITION 49. For any positive integers n and r define the coefficients $A_i^{n,r}$ by

$$\sum_{i=0}^{n(r-1)} A_i^{n,r} T^i = (1 + T + \cdots + T^{r-1})^n.$$

We also define the vector $\mathbf{A}^{n,r}$ as $(A_0^{n,r}, A_1^{n,r}, \dots, A_{n(r-1)}^{n,r})$.

Looking at the definition of the Katzman coefficient, we see that $A_j^{n,2} = \binom{n}{j}$ and $A_j^{n,1} = 0$ unless $j = 0$, in which case we have $A_0^{n,1} = 1$.

Below we derive some new and useful equalities and prove symmetry and unimodality for the Katzman coefficients. Katzman [Kat05] gave an explicit equation for the h^* -vector of uniform matroid polytopes and the coefficients of their Ehrhart polynomials, although he did not use the same language. We restate it here for our purposes:

LEMMA 50 (See Corollary 2.9 in [Kat05]). *Let $\mathcal{P}(U^{r,n})$ be the matroid polytope of the uniform matroid of rank r on n elements. Then the h^* -polynomial of $\mathcal{P}(U^{r,n})$ is given by*

$$(3.5) \quad \sum_{s=0}^{r-1} \sum_{j=0}^s \sum_{k=0}^j \sum_{l \geq k} (-1)^{s+j+k} \left[\binom{n}{s} \binom{s}{j} \binom{j}{k} A_{(l-k)(r-s)}^{n-j, r-s} \right] T^l.$$

That is, for $\mathbf{h}^*(\mathcal{P}(U^{n,r})) = (h_0^*, \dots, h_r^*)$,

$$h_l^* = \sum_{s=0}^{r-1} \sum_{j=0}^s \sum_{k=0}^j (-1)^{s+j+k} \binom{n}{s} \binom{s}{j} \binom{j}{k} A_{(l-k)(r-s)}^{n-j, r-s}.$$

For $r = 2$ the h^* -polynomial of $\mathcal{P}(U)^{n,2}$ is

$$(3.6) \quad \left(\sum_{l \geq 0} \binom{n}{2l} T^l \right) - nT.$$

The following lemma is a direct consequence of Corollary 2.9 in [Kat05]:

LEMMA 51. *Let $\mathcal{P}(U^{2,n})$ be the matroid polytope of the uniform matroid of rank 2 on n elements. The h^* -vector of $\mathcal{P}(U^{2,n})$ is unimodal.*

The rank 2 case is an interesting example already. Although the h^* -vector is unimodal, it is not always symmetric. Next we present some useful lemmas, the first a combinatorial description of the Katzman coefficients.

LEMMA 52. *For $i = 0, \dots, n(r-1)$ we have*

$$(3.7) \quad A_i^{n,r} = \sum_{\substack{0a_0+1a_1+\dots+(r-1)a_{r-1}=i \\ a_0+a_1+\dots+a_{r-1}=n}} \binom{n}{a_0, a_1, \dots, a_{r-1}}$$

where $a_0, \dots, a_{r-1}$ run through non-negative integers.

PROOF. Using the multinomial formula [Sta97] we have

$$\begin{aligned} \sum_{i=0}^{n(r-1)} A_i^{n,r} T^i &= (1 + T + \dots + T^{r-1})^n \\ &= \sum_{a_0+a_1+\dots+a_{r-1}=n} \binom{n}{a_0, a_1, \dots, a_{r-1}} 1^{a_0} T^{a_1} T^{2a_2} \dots T^{(r-1)a_{r-1}} \\ &= \sum_{a_0+a_1+\dots+a_{r-1}=n} \binom{n}{a_0, a_1, \dots, a_{r-1}} T^{0a_0+1a_1+\dots+(r-1)a_{r-1}}. \end{aligned}$$

By grouping the powers of T we get equation (3.7). □

Next we present a generalization of a property of the binomial coefficients. The following lemma relates the Katzman coefficients to Katzman coefficients with one less element.

LEMMA 53.

$$(3.8) \quad A_i^{n,r} = \sum_{k=i-r+1}^i A_k^{n-1,r}$$

where we define $A_p^{n-1,r} := 0$ when $p < 0$ or $p > (n-1)(r-1)$.

PROOF.

$$\begin{aligned}
& \sum_{i=0}^{n(r-1)} A_i^{n,r} T^i \\
&= (1 + T + \cdots + T^{r-1})^n = (1 + T + \cdots + T^{r-1})^{n-1} (1 + T + \cdots + T^{r-1}) \\
&= \left(\sum_{i=0}^{(n-1)(r-1)} A_i^{n-1,r} T^i \right) (1 + T + \cdots + T^{r-1}) \\
&= \sum_{i=0}^{(n-1)(r-1)} A_i^{n-1,r} T^i + \sum_{i=0}^{(n-1)(r-1)} A_i^{n-1,r} T^{i+1} + \cdots + \sum_{i=0}^{(n-1)(r-1)} A_i^{n-1,r} T^{i+r-1} \\
&= \sum_{i=0}^{(n-1)(r-1)} A_i^{n-1,r} T^i + \sum_{i=1}^{(n-1)(r-1)+1} A_{i-1}^{n-1,r} T^i + \cdots + \sum_{i=r-1}^{(n-1)(r-1)+r-1} A_{i-r+1}^{n-1,r} T^i \\
&= \sum_{i=0}^{(n-1)(r-1)+r-1} \left(A_i^{n-1,r} + A_{i-1}^{n-1,r} + \cdots + A_{i-r+1}^{n-1,r} \right) T^i.
\end{aligned}$$

Thus we get equation (3.8). $\square$

The following lemma relates the Katzman coefficients of rank r with those of rank $r-1$.

LEMMA 54.

$$\sum A_i^{n,r} T^i = \sum_{k=0}^n \binom{n}{k} T^k \left(\sum_{l=0}^{k(r-2)} A_l^{k,r-1} T^l \right)$$

or in other words

$$A_i^{n,r} = \sum_{\substack{k+l=i \\ 0 \leq k \leq n \\ 0 \leq l \leq k(r-2)}} \binom{n}{k} A_l^{k,r-1}.$$

PROOF. From Definition 49

$$\begin{aligned}
& \sum_{i=0}^{n(r-1)} A_i^{n,r} T^i = (1 + T + \cdots + T^{r-1})^n = (1 + [T + \cdots + T^{r-1}])^n \\
&= \sum_{k=0}^n \binom{n}{k} [T + \cdots + T^{r-1}]^k = \sum_{k=0}^n \binom{n}{k} T^k [1 + \cdots + T^{r-2}]^k \\
&= \sum_{k=0}^n \binom{n}{k} T^k \left(\sum_{l=0}^{k(r-2)} A_l^{k,r-1} T^l \right).
\end{aligned}$$

$\square$

LEMMA 55. *The Katzman coefficients are unimodal and symmetric in the index i . That is, the vector $(A_0^{n,r}, A_1^{n,r}, \dots, A_{n(r-1)}^{n,r})$ is unimodal and symmetric.*

PROOF. We first prove symmetry. Considering equation (3.7) we assume that

$$0a_0 + 1a_1 + \dots + (r-1)a_{r-1} = i \quad \text{and} \quad a_0 + a_1 + \dots + a_{r-1} = n.$$

These two assumptions imply that

$$\begin{aligned} & ((r-1) - 0)a_0 + ((r-1) - 1)a_1 + \dots + ((r-1) - (r-1))a_{r-1} \\ &= (r-1)a_0 + (r-1)a_1 + \dots + (r-1)a_{r-1} - 0a_0 - 1a_1 - \dots - (r-1)a_{r-1} \\ &= (r-1)n - i. \end{aligned}$$

Therefore

$$\begin{aligned} A_i^{n,r} &= \sum_{\substack{0a_0 + 1a_1 + \dots + (r-1)a_{r-1} = i \\ a_0 + a_1 + \dots + a_{r-1} = n}} \binom{n}{a_0, a_1, \dots, a_{r-1}} \\ &= \sum_{\substack{(r-1)a_0 + (r-2)a_1 + \dots + 0a_{r-1} = i \\ a_0 + a_1 + \dots + a_{r-1} = n}} \binom{n}{a_0, a_1, \dots, a_{r-1}} = A_{n(r-1)-i}^{n,r}. \end{aligned}$$

To prove unimodality we proceed by induction on n , where r is fixed. First, $\mathbf{A}^{1,r}$ is unimodal. Assume for $n-1$ that $\mathbf{A}^{n-1,r} = (A_0^{n-1,r}, A_1^{n-1,r}, \dots, A_{(n-1)(r-1)}^{n-1,r})$ is unimodal. Using equation (3.8) and the fact that $\mathbf{A}^{n-1,r}$ is symmetric we get that $\mathbf{A}^{n,r}$ is unimodal. To help see this, one can view equation (3.8) as a sliding window over r elements of the vector $\mathbf{A}^{n-1,r}$, that is, $A_i^{n,r}$ is equal to the sum of the r elements in a window over the vector $\mathbf{A}^{n-1,r}$. As the window slides up the vector $\mathbf{A}^{n-1,r}$, the sum will increase. When the window is on the center of $\mathbf{A}^{n-1,r}$ symmetry and unimodality of $\mathbf{A}^{n-1,r}$ will imply unimodality of $\mathbf{A}^{n,r}$. $\square$

Now we use the explicit equation for the h^* -vector of uniform matroid polytopes to prove partial unimodality of rank 3 uniform matroids. First we note that the coefficient of T^l , h_l^* , in equation (3.5) is

$$h_l^* = \sum_{s=0}^{r-1} \sum_{j=0}^s \sum_{k=0}^j (-1)^{s+j+k} \binom{n}{s} \binom{s}{j} \binom{j}{k} A_{(p-k)(r-s)}^{n-j, r-s}.$$

Letting the rank $r = 3$, and using equation (3.5), we get the h^* -polynomial (which is grouped by values of s from (3.5)),

$$\sum_{l \geq 0} \left[\binom{n}{0} A_{3l}^{n,3} + \binom{n}{1} \left(-A_{2l}^{n,2} + A_{2l}^{n-1,2} - A_{2(l-1)}^{n-1,2} \right) + \right. \\ \left. + \binom{n}{2} \left(A_l^{n,1} - 2A_l^{n-1,1} + 2A_{l-1}^{n-1,1} + A_l^{n-2,1} - 2A_{l-1}^{n-2,1} + A_{l-2}^{n-2,1} \right) \right] T^l.$$

Now using that $A_i^{n,2} = \binom{n}{i}$ and $A_i^{n,1} = \delta_0(i)$, where $\delta_j(p) = \begin{cases} 1 & \text{if } p = j \\ 0 & \text{else} \end{cases}$, we get

$$\sum_{l \geq 0} \left[A_{3l}^{n,3} + n \left(-\binom{n}{2l} + \binom{n-1}{2l} - \binom{n-1}{2l-2} \right) + \right. \\ \left. + \binom{n}{2} (\delta_0(l) - 2\delta_0(l) + 2\delta_1(l) + \delta_0(l) - 2\delta_1(l) + \delta_2(l)) \right] T^l \\ = \sum_{l \geq 0} \left[A_{3l}^{n,3} + n \left(-\binom{n}{2l} + \binom{n-1}{2l} - \binom{n-1}{2l-2} \right) + \delta_2(l) \binom{n}{2} \right] T^l.$$

Using properties of the binomial coefficients, we see that

$$\left[-\binom{n}{2l} \right] + \binom{n-1}{2l} - \binom{n-1}{2l-2} = \left[-\binom{n-1}{2l} - \binom{n-1}{2l-1} \right] + \binom{n-1}{2l} - \binom{n-1}{2l-2} \\ = -\binom{n-1}{2l-1} - \binom{n-1}{2l-2} \\ = -\binom{n}{2l-1}.$$

So the h^* -polynomial of rank three uniform matroid polytopes is

$$(3.9) \quad \sum_{l \geq 0} \left[A_{3l}^{n,3} - n \binom{n}{2l-1} + \delta_2(l) \binom{n}{2} \right] T^l.$$

Using Lemma 3.8, the coefficient of T^l , if $3l \leq n$, is

$$(3.10) \quad h_l^* = \sum_{\substack{k+p=3l \\ 0 \leq p \leq k \leq n}} \binom{n}{k} \binom{k}{p} - n \binom{n}{2l-1} + \delta_2(l) \binom{n}{2}$$

$$= \left[\binom{n}{3l} \binom{3l}{0} + \binom{n}{3l-1} \binom{3l-1}{1} + \cdots + \binom{n}{3l-\lfloor 3l/2 \rfloor} \binom{3l-\lfloor 3l/2 \rfloor}{\lfloor 3l/2 \rfloor} \right] + \\ - n \binom{n}{2l-1} + \delta_2(l) \binom{n}{2}.$$

Next we show that when g is fixed $A_g^{n,3}$ is a polynomial of degree g in the indeterminate n , with positive leading coefficient. Assume $g \leq n$. Considering Lemma 54 and when $g \leq n$,

$$(3.11) \quad A_g^{n,3} = \sum_{\substack{k+p=g \\ 0 \leq p \leq k \leq n}} \binom{n}{k} \binom{k}{p}$$

where $\binom{n}{q}$ is a polynomial of degree q with positive leading coefficient. The highest degree polynomial in the sum is $\binom{n}{g}$, a degree g polynomial. Hence $A_g^{n,3}$ is a polynomial of degree g in the indeterminate n , with positive leading coefficient. If $g \geq n$, then $A_g^{n,3} = A_{n-g}^{n,3}$ since the Katzman coefficients are symmetric by Lemma 55.

PROOF OF THEOREM 47 PART (2). Let I be a non-negative integer. From above we see that $A_g^{n,3}$ is a degree g polynomial in the indeterminate n , with positive leading coefficient. Equation (3.9) is the h^* -polynomial of $U^{3,n}$, which is a sum of polynomials in n , the highest degree polynomial being $A_{3l}^{n,3}$, a polynomial of degree $3l$. So, $h_l^* - h_{l-1}^*$ is the difference of a degree $3l$ and $3(l-1)$ polynomial, hence $h_l^* - h_{l-1}^*$ is a degree $3l$ polynomial with positive leading coefficient. For sufficiently large n , call it $n(I)$, $h_l^* - h_{l-1}^*$ is positive for $0 \leq l \leq I$. Hence, the h^* -vector of $U^{3,n}$ is non-decreasing up to the index I for $n \geq n(I)$. $\square$

One might ask if equation (3.11) has a simpler form. We ran the WZ algorithm on our expression, which proved that equation (3.11) can not be written as a linear combination of a fixed number of hypergeometric terms (*closed form*) [PWZ96]. There is still the possibility that this expression has a simpler form, though not a closed form as described above.

3.2. Unimodular Simplices of Matroid Polytopes

A k -simplex $S = \{s_1, \dots, s_{k+1}\}$ is the collection of $k+1$ affinely independent points in $\mathbb{R}^n$ and we say $\text{conv}(S) := \text{conv}(s_1, \dots, s_{k+1})$. When we refer to a simplex S we consider it both as a collection of points and as a convex polytope when the context is clear. Let $\mathcal{P} \subseteq \mathbb{R}^n$ be a polytope with vertices $\mathbf{v}_1, \dots, \mathbf{v}_p$. A *triangulation* $\mathcal{T}$ of $\mathcal{P}$ is a collection of simplices on the vertices $\mathbf{v}_1, \dots, \mathbf{v}_p$ of $\mathcal{P}$ such that

- (i) If $A \in \mathcal{T}$ then all faces of A are in $\mathcal{T}$,
- (ii) If $A, B \in \mathcal{T}$ then $A \cap B$ is a face of both A and B ,
- (iii) $\bigcup_{A \in \mathcal{T}} \text{conv}(A) = \mathcal{P}$.

It is sufficient to list only the highest dimensional simplices of a triangulation. A triangulation $\mathcal{T}$ is *unimodular* if the volume of every highest dimensional simplex of $\mathcal{T}$ are the same. We also say a simplex is unimodular if its normalized volume is one [DLRS09]. A vector in R^n is a ± 1 -vector if it contains $+1$ and -1 and the rest are zeros.

Recall that two elements $s, t \in S$ of a matroid $\mathcal{M} = (S, \mathcal{I})$ are *connected* if there exists a circuit that contains s and t . This defines an equivalence relation on S and moreover $\mathcal{M}$ can be written as a direct sum of restrictions to the connected components of $\mathcal{M}$ [Oxl92]. This implies a matroid polytope $\mathcal{P}_{\mathcal{M}}$ can be written as a direct product of connected matroid polytopes. We note that if a polytope $\mathcal{P}$ is a direct product of polytopes $\mathcal{Q}_1, \dots, \mathcal{Q}_k$ which each have a unimodular triangulation, then $\mathcal{P}$ has a unimodular triangulation [DLRS09]. Hence, we need only consider connected matroids when investigating unimodular triangulations.

We wish to find sufficient conditions such that a $(n-1)$ -simplex of a connected matroid polytope $\mathcal{P}_{\mathcal{M}}$ is unimodular. We will see later that a simplex of $\mathcal{P}_{\mathcal{M}}$ is unimodular if and only if the determinant of the vertices is $\text{rank}(\mathcal{M})$.

Neil White proposed an algebraic conjecture on the toric ideal determined by a matroid [Whi80]. Let $\mathcal{F}$ be a field and define the polynomial ring $S_{\mathcal{M}} := \mathcal{F}[\mathbf{y}_B \mid B \in \mathcal{B}_{\mathcal{M}}]$. Let the toric ideal $I_{\mathcal{M}}$ be defined as the kernel of the homomorphism $S_{\mathcal{M}} \rightarrow \mathcal{F}[x_1, \dots, x_n]$ given by $\mathbf{y}_B \mapsto \prod_{i \in B} x_i$. Given any pair of bases $B_1, B_2 \in \mathcal{B}$ the *symmetric exchange property* states that for every $x \in B_1$ there exists $y \in B_2$ such that $B_1 - x + y$ and $B_2 - y + x$ are bases and it is said that $x \in B_1$ *double swaps* into B_2 . If $B_1 - x + y$ and $B_2 - y + x$ are bases they are called a *double swap*. White proposed the following conjecture:

CONJECTURE 56 ([Whi80]). *For any matroid $\mathcal{M}$, the toric ideal $I_{\mathcal{M}}$ is generated by the quadratic binomials $\mathbf{y}_{B_1}\mathbf{y}_{B_2} - \mathbf{y}_{D_1}\mathbf{y}_{D_2}$ such that the pair of bases D_1, D_2 can be obtained from the pair B_1, B_2 by a double swap.*

A stronger conjecture is that the previous quadratic binomials are in fact a Gröbner bases for some term ordering. This implies that there exists a regular unimodular triangulation of any matroid polytope. The following is a geometric variation of White's famous conjecture:

CONJECTURE 57. *Let $\mathcal{M}$ be a matroid of rank r on n elements. There exists a unimodular triangulation of $\mathcal{P}_{\mathcal{M}}$.*

In [Stu96] it was shown that all uniform matroids have a regular unimodular triangulation. A partial proof of White's original conjecture was given in [Bla08] for graphical matroids, but Conjecture 57 has yet to be proved in general.

A *covering* of $\mathcal{P}$ is a triangulation without condition (ii). That is, the simplices of a covering can intersect in any fashion, as long as the polytope is covered. See Figure 3.1. If asking for a unimodular triangulation is too much, we propose a weaker conjecture:




FIGURE 3.1. A covering of a hexagon.

CONJECTURE 58. *Let $\mathcal{M}$ be a matroid of rank r on n elements. There exists a unimodular covering of $\mathcal{P}_{\mathcal{M}}$.*

Coverings are important to matroid polytopes. If a unimodular covering exists for a connected matroid polytope $\mathcal{P}_{\mathcal{M}}$ it would imply its Caratheodory rank is n . In other words, every integer point in the cone of $\mathcal{P}_{\mathcal{M}}$ can be written as an integral linear combination of vertices of $\mathcal{P}_{\mathcal{M}}$. That is, the vertices of $\mathcal{P}_{\mathcal{M}}$ are a Hilbert basis for the cone of $\mathcal{P}_{\mathcal{M}}$. See [CS86, Seb90, dPS03].

We now focus on sufficient conditions for the existence unimodular simplices of matroid polytopes and begin by defining two graphs on any collection of incidence vectors of a matroid polytope.

DEFINITION 59. Let $\mathcal{M}$ be a matroid on n elements. Given a subset $X \subseteq \text{Inc}(\mathcal{B}_{\mathcal{M}}) \subseteq \mathbb{R}^n$ we define two graphs G and g on n nodes. The graph $G(X)$ has an edge (i, j) for every pair of bases $\mathbf{X}_i, \mathbf{X}_j \in X$ that are adjacent, i.e. $\mathbf{X}_i - \mathbf{X}_j = \mathbf{e}_s - \mathbf{e}_t$ for some s, t . The graph $g(X)$ has an edge (i, j) every pair of adjacent bases $\mathbf{X}_s, \mathbf{X}_t \in X$ such that they differ in position i and j , i.e. $\mathbf{X}_s - \mathbf{X}_t = \mathbf{e}_i - \mathbf{e}_j$. Succinctly we define $G(X)$ and $g(X)$ as:

$$G(X) := ([n], \{ (i, j) \mid \mathbf{X}_i, \mathbf{X}_j \in X, \mathbf{X}_i - \mathbf{X}_j = \mathbf{e}_s - \mathbf{e}_t \})$$

$$g(X) := ([n], \{ (i, j) \mid \mathbf{X}_s, \mathbf{X}_t \in X, \mathbf{X}_s - \mathbf{X}_t = \mathbf{e}_i - \mathbf{e}_j \})$$

where $\mathbf{e}_i, \mathbf{e}_j, \mathbf{e}_s, \mathbf{e}_t \in \mathbb{R}^n$ are standard unit vectors.

Throughout this section we will take X to mean a set of incidence vectors as well as a 0/1 matrix with rows as the incidence vectors. The following proposition is easy to see using elementary row sums on X and cofactor expansion on the resulting last row.

PROPOSITION 60. *Let $\mathcal{M}$ be a connected matroid of rank r on n elements and $X \subset \text{Inc}(\mathcal{B}_{\mathcal{M}})$ be n linearly independent incidence vectors of $\mathcal{P}_{\mathcal{M}}$. Then $|\det(X)| = kr$, $k \in \mathbb{N}$.*

EXAMPLE 61. Consider the following incidence vectors X in row form. We label the rows $\mathbf{1}, \dots, \mathbf{6}$ and columns $\mathbf{a}, \dots, \mathbf{f}$.

$$\begin{array}{c} \mathbf{a} \quad \mathbf{b} \quad \mathbf{c} \quad \mathbf{d} \quad \mathbf{e} \quad \mathbf{f} \\ \mathbf{1} \left(\begin{array}{cccccc} 1 & 1 & 0 & 0 & 1 & 0 \end{array} \right) \\ \mathbf{2} \left(\begin{array}{cccccc} 1 & 1 & 0 & 0 & 0 & 1 \end{array} \right) \\ \mathbf{3} \left(\begin{array}{cccccc} 1 & 0 & 1 & 1 & 0 & 0 \end{array} \right) \\ \mathbf{4} \left(\begin{array}{cccccc} 0 & 1 & 1 & 1 & 0 & 0 \end{array} \right) \\ \mathbf{5} \left(\begin{array}{cccccc} 0 & 0 & 1 & 0 & 1 & 1 \end{array} \right) \\ \mathbf{6} \left(\begin{array}{cccccc} 0 & 0 & 0 & 1 & 1 & 1 \end{array} \right) \end{array} \quad \begin{array}{l} G(X) = ([6], \{(\mathbf{1}, \mathbf{2}), (\mathbf{3}, \mathbf{4}), (\mathbf{5}, \mathbf{6})\}) \\ g(X) = ([6], \{(\mathbf{a}, \mathbf{b}), (\mathbf{c}, \mathbf{d}), (\mathbf{e}, \mathbf{f})\}) \end{array}$$

We can equivalently define $g(X)$ as the graph whose incidence matrix is given by

$$\{ \mathbf{X}_s - \mathbf{X}_t \mid \mathbf{X}_s, \mathbf{X}_t \in X, \mathbf{X}_s - \mathbf{X}_t = \mathbf{e}_i - \mathbf{e}_j \}$$

where $\mathbf{e}_i, \mathbf{e}_j \in \mathbb{R}^n$ are standard unit vectors. We will use this fact for the following theorem.

THEOREM 62. *Let $\mathcal{M}$ be a connected matroid of rank r on n elements and $X \subset \text{Inc}(\mathcal{B}_{\mathcal{M}})$ be n linearly independent incidence vectors of $B_{\mathcal{M}}$. Then*

- (i) $G(X)$ and $g(X)$ have the same number of connected components c ,
- (ii) Let $X = \{\mathbf{v}^1, \dots, \mathbf{v}^n\}$ and without loss of generality assume $\{1, \dots, l_1-1\}, \{l_1, \dots, l_2-1\}, \dots, \{l_{c-1}, \dots, n\}$, are the connected components of $g(X)$. Then

$$(3.12) \quad |\det(X)| = \det \begin{pmatrix} \sum_{i=1}^{l_1-1} v_i^{m_1} & \sum_{i=l_1}^{l_2-1} v_i^{m_1} & \cdots & \sum_{i=l_{c-1}}^n v_i^{m_1} \\ \sum_{i=1}^{l_1-1} v_i^{m_2} & \sum_{i=l_1}^{l_2-1} v_i^{m_2} & \cdots & \sum_{i=l_{c-1}}^n v_i^{m_2} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{i=1}^{l_1-1} v_i^{m_c} & \sum_{i=l_1}^{l_2-1} v_i^{m_c} & \cdots & \sum_{i=l_{c-1}}^n v_i^{m_c} \end{pmatrix}$$

where $\mathbf{v}^{m_1}, \dots, \mathbf{v}^{m_c}$ are representative incidence vectors of the connected components of $G(X)$.

PROOF. We offer a constructive proof of the theorem. Assume $G(X)$ has p connected component and $g(X)$ has c connected components. First let $\mathbf{v}^{m_1}, \dots, \mathbf{v}^{m_p}$ be representative incidence vectors of the connected components of $G(X)$. Performing elementary row operations such as row addition, difference or reordering will only change the sign of the determinant. First we organize the rows of X by the p connected components of $G(X)$.

$$X = \begin{pmatrix} v_1^{m_1} & \cdots & v_n^{m_1} \\ \vdots & \ddots & \vdots \\ \hline v_1^{m_2} & \cdots & v_n^{m_2} \\ \vdots & \ddots & \vdots \\ \hline \vdots & \ddots & \vdots \\ \hline v_1^{m_p} & \cdots & v_n^{m_p} \\ \vdots & \ddots & \vdots \end{pmatrix}$$

Consider a connected component of $G(X)$ with representative incidence vector $\mathbf{v}^{m_i}$. Let $\mathbf{v}^{m_i}$ be the root of a spanning tree T of the corresponding component of $G(X)$. For all non-root vectors in the component determined by $\mathbf{v}^{m_i}$ perform the row operation $\mathbf{v}^g := \mathbf{v}^g - \mathbf{v}^h$ where h is the parent of g . Now all the rows except $\mathbf{v}^{m_i}$ of the component of $G(X)$ are

± 1 -vectors. Repeat for each connected component of $G(X)$. (By $\dots \pm 1 \dots$ we mean some ± 1 vector).

$$\begin{pmatrix} v_1^{m_1} & \dots & v_n^{m_1} \\ & \dots \pm 1 \dots & \\ & \vdots & \\ & \dots \pm 1 \dots & \\ \hline v_1^{m_2} & \dots & v_n^{m_2} \\ & \vdots & \vdots \\ & \ddots & \vdots \\ \hline v_1^{m_p} & \dots & v_n^{m_p} \\ & \vdots & \vdots \end{pmatrix} \rightarrow \begin{pmatrix} v_1^{m_1} & \dots & v_n^{m_1} \\ & \dots \pm 1 \dots & \\ & \vdots & \\ & \dots \pm 1 \dots & \\ \hline v_1^{m_2} & \dots & v_n^{m_2} \\ & \dots \pm 1 \dots & \\ & \vdots & \\ & \dots \pm 1 \dots & \\ \hline \vdots & \ddots & \vdots \\ \hline v_1^{m_p} & \dots & v_n^{m_p} \\ & \dots \pm 1 \dots & \\ & \vdots & \\ & \dots \pm 1 \dots & \end{pmatrix}$$

Next, due to the assumption on the ordering of the connected components of $g(X)$ we partition the columns of matrix X by the components of $g(X)$. We also reorder the rows with the representatives vectors $\mathbf{v}^{m_1}, \dots, \mathbf{v}^{m_p}$ on top and the ± 1 -vectors by their corresponding components of $g(X)$.

$$\begin{pmatrix} v_1^{m_1} \dots v_{l_1-1}^{m_1} & v_{l_1}^{m_1} \dots v_{l_2-1}^{m_1} & \dots & v_{l_{c-1}}^{m_1} \dots v_n^{m_1} \\ v_1^{m_2} \dots v_{l_1-1}^{m_2} & v_{l_1}^{m_2} \dots v_{l_2-1}^{m_2} & \dots & v_{l_{c-1}}^{m_2} \dots v_n^{m_2} \\ \vdots & \vdots & \ddots & \vdots \\ v_1^{m_p} \dots v_{l_1-1}^{m_p} & v_{l_1}^{m_p} \dots v_{l_2-1}^{m_p} & \dots & v_{l_{c-1}}^{m_p} \dots v_n^{m_p} \\ \hline \dots \pm 1 \dots & 0 \dots 0 & \dots & 0 \dots 0 \\ \vdots & \vdots & \ddots & \vdots \\ \dots \pm 1 \dots & 0 \dots 0 & \dots & 0 \dots 0 \\ \hline 0 \dots 0 & \dots \pm 1 \dots & \dots & 0 \dots 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 \dots 0 & \dots \pm 1 \dots & \dots & 0 \dots 0 \\ \hline \dots & \dots & \ddots & \dots \\ \hline 0 \dots 0 & 0 \dots 0 & \dots & \dots \pm 1 \dots \\ \vdots & \vdots & \ddots & \vdots \\ 0 \dots 0 & 0 \dots 0 & \dots & \dots \pm 1 \dots \end{pmatrix}$$

We note that for isolated nodes of $g(X)$ there will be no corresponding ± 1 -vectors, but this will not interfere with this proof. Consider any submatrix of ± 1 -vectors determined by a connected component C of $g(X)$. First, if there are k many nodes in C , then there will be $k - 1$ ± 1 -vectors since X is linearly independent and these vectors are a spanning

tree for an orientation of the component C . Using the connectedness of the component C , elementary row addition and the fact that ± 1 -vectors sum to 0 we can rewrite the parts of each representative incidence vector $\mathbf{v}_{m_1}, \dots, \mathbf{v}_{m_p}$ contained in the columns corresponding to C . We sum the elements of each representative incidence vector over the component C and place it in the first position, with 0's in all other position of C .

$$\begin{pmatrix}
 \sum_{i=1}^{l_1-1} v_i^{m_1} 0 \dots 0 & v_{l_1}^{m_1} \dots v_{l_2-1}^{m_1} & \dots & v_{l_{c-1}}^{m_1} \dots v_n^{m_1} \\
 \sum_{i=1}^{l_1-1} v_i^{m_2} 0 \dots 0 & v_{l_1}^{m_2} \dots v_{l_2-1}^{m_2} & \dots & v_{l_{c-1}}^{m_2} \dots v_n^{m_2} \\
 \vdots & \vdots & \ddots & \vdots \\
 \sum_{i=1}^{l_1-1} v_i^{m_p} 0 \dots 0 & v_{l_1}^{m_p} \dots v_{l_2-1}^{m_p} & \dots & v_{l_{c-1}}^{m_p} \dots v_n^{m_p} \\
 \hline
 \dots \pm 1 \dots & 0 \dots 0 & \dots & 0 \dots 0 \\
 \vdots & \vdots & \ddots & \vdots \\
 \dots \pm 1 \dots & 0 \dots 0 & \dots & 0 \dots 0 \\
 \hline
 0 \dots 0 & \dots \pm 1 \dots & \dots & 0 \dots 0 \\
 \vdots & \vdots & \ddots & \vdots \\
 0 \dots 0 & \dots \pm 1 \dots & \dots & 0 \dots 0 \\
 \hline
 \dots & \dots & \ddots & \dots \\
 \hline
 0 \dots 0 & 0 \dots 0 & \dots & \dots \pm 1 \dots \\
 \vdots & \vdots & \ddots & \vdots \\
 0 \dots 0 & 0 \dots 0 & \dots & \dots \pm 1 \dots
 \end{pmatrix} \rightarrow
 \begin{pmatrix}
 \sum_{i=1}^{l_1-1} v_i^{m_1} 0 \dots 0 & \sum_{i=l_1}^{l_2-1} v_i^{m_1} 0 \dots 0 & \dots & \sum_{i=l_{c-1}}^n v_i^{m_1} 0 \dots 0 \\
 \sum_{i=1}^{l_1-1} v_i^{m_2} 0 \dots 0 & \sum_{i=l_1}^{l_2-1} v_i^{m_2} 0 \dots 0 & \dots & \sum_{i=l_{c-1}}^n v_i^{m_2} 0 \dots 0 \\
 \vdots & \vdots & \ddots & \vdots \\
 \sum_{i=1}^{l_1-1} v_i^{m_p} 0 \dots 0 & \sum_{i=l_1}^{l_2-1} v_i^{m_p} 0 \dots 0 & \dots & \sum_{i=l_{c-1}}^n v_i^{m_p} 0 \dots 0 \\
 \hline
 \dots \pm 1 \dots & 0 \dots 0 & \dots & 0 \dots 0 \\
 \vdots & \vdots & \ddots & \vdots \\
 \dots \pm 1 \dots & 0 \dots 0 & \dots & 0 \dots 0 \\
 \hline
 0 \dots 0 & \dots \pm 1 \dots & \dots & 0 \dots 0 \\
 \vdots & \vdots & \ddots & \vdots \\
 0 \dots 0 & \dots \pm 1 \dots & \dots & 0 \dots 0 \\
 \hline
 \dots & \dots & \ddots & \dots \\
 \hline
 0 \dots 0 & 0 \dots 0 & \dots & \dots \pm 1 \dots \\
 \vdots & \vdots & \ddots & \vdots \\
 0 \dots 0 & 0 \dots 0 & \dots & \dots \pm 1 \dots
 \end{pmatrix}$$

For each component C of $g(X)$ there exists a column corresponding to a leaf of the spanning tree determined by the ± 1 vectors in C , rooted at the first column. This column contains all 0's and one 1. Performing cofactor expansion down this column eliminates a row and column. This can be repeated until all ± 1 -vectors of C are eliminated, preserving the

summed parts of the incidence vectors. Repeating for all connected components, all ± 1 -vectors can be eliminated while preserving the determinant of X (under a sign change). We get the desired form and since each operation preserves the matrix being square we see that $c = p$.

$$\begin{pmatrix} \sum_{i=1}^{l_1-1} v_i^{m_1} & \sum_{i=l_1}^{l_2-1} v_i^{m_1} & \cdots & \sum_{i=l_{c-1}}^n v_i^{m_1} \\ \sum_{i=1}^{l_1-1} v_i^{m_2} & \sum_{i=l_1}^{l_2-1} v_i^{m_2} & \cdots & \sum_{i=l_{c-1}}^n v_i^{m_2} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{i=1}^{l_1-1} v_i^{m_c} & \sum_{i=l_1}^{l_2-1} v_i^{m_c} & \cdots & \sum_{i=l_{c-1}}^n v_i^{m_c} \end{pmatrix} \in \mathbb{R}^{c \times c}$$

□

COROLLARY 63. *Let $\mathcal{M}$ be a connected matroid of rank r on n elements and $X \subset \text{Inc}(\mathcal{B}_{\mathcal{M}})$ be n linearly independent incidence vectors of $\mathcal{B}_{\mathcal{M}}$. If $G(X)$ is connected then $|\det(X)| = r$.*

EXAMPLE 64. We offer an example of [Theorem 62](#) in action. Consider the uniform matroid $U^{2,6}$ and the following collection of incidence vectors of bases in row form. Partition into components of $G(X)$ and $g(X)$.

$$\left| \begin{array}{cccccc} 1 & 1 & 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{array} \right| \Rightarrow \left| \begin{array}{cc|cc|cc} 1 & 1 & 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 & 1 \\ \hline 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{array} \right|$$

Find rooted spanning trees of each component of $G(X)$ and take differences to get ± 1 -vectors and the representative vectors.

$$\left| \begin{array}{cc|cc|cc} 1 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 \\ \hline 1 & 0 & 1 & 1 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & -1 & 1 & 0 & 0 \end{array} \right| \Rightarrow \left| \begin{array}{cc|cc|cc} 2 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 \\ \hline 1 & 0 & 2 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 & 2 & 0 \\ 0 & 0 & -1 & 1 & 0 & 0 \end{array} \right|$$

For each component of $g(X)$ eliminate use ± 1 -vectors to eliminate rows and columns using cofactor expansion.

$$\begin{array}{c}
 \left| \begin{array}{c|c|c|c|c|c} 2 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 \\ 1 & 0 & 2 & 0 & 0 & 0 \\ \hline -1 & 1 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 & 2 & 0 \\ 0 & 0 & -1 & 1 & 0 & 0 \end{array} \right| \Rightarrow \left| \begin{array}{cccccc} 2 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 & 1 \\ 1 & 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 2 & 0 \\ 0 & -1 & 1 & 0 & 0 \end{array} \right| \\
 \\
 \left| \begin{array}{c|c|c|c|c} 2 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 & 1 \\ 1 & 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 2 & 0 \\ \hline 0 & -1 & 1 & 0 & 0 \end{array} \right| \Rightarrow \left| \begin{array}{cccc} 2 & 0 & 1 & 0 \\ 0 & 0 & -1 & 1 \\ 1 & 2 & 0 & 0 \\ 0 & 1 & 2 & 0 \end{array} \right| \\
 \\
 \left| \begin{array}{c|c|c|c} 2 & 0 & 1 & 0 \\ 0 & 0 & -1 & 1 \\ \hline 1 & 2 & 0 & 0 \\ 0 & 1 & 2 & 0 \end{array} \right| \Rightarrow \left| \begin{array}{ccc} 2 & 0 & 1 \\ 1 & 2 & 0 \\ 0 & 1 & 2 \end{array} \right|
 \end{array}$$

We explore more connections between the graphs $G(X)$ and $g(X)$. The following is an alternate proof of part of [Theorem 62](#)

LEMMA 65. *Let $\mathcal{M}$ be a connected matroid of rank r on n elements and $X \subset \text{Inc}(\mathcal{B}_{\mathcal{M}})$ be n linearly independent incidence vectors of $\mathcal{B}_{\mathcal{M}}$. If $G(X)$ is connected then $g(X)$ is connected.*

PROOF. Consider the matrix X and, as in the proof of [Theorem 62](#), we pick a row as a root of the spanning tree of $G(X)$ and perform row operations so that the resulting matrix is our root and the rest are ± 1 -vectors. If $g(X)$ is not connected then let S be one connected component and $T = [n] \setminus S$. Also, let $s = |S|$ and $t = |T|$. Then the following non-zero vector is orthogonal to all the elements of X , contradicting its full-dimensionality.

$$t \sum_{i \in S} \mathbf{e}_i - s \sum_{j \in T} \mathbf{e}_j$$

□

More generally we can show the following:

LEMMA 66. *Let $\mathcal{M}$ be a connected matroid of rank r on n elements and $X \subset \text{Inc}(\mathcal{B}_{\mathcal{M}})$ be n incidence vectors of $\mathcal{M}$. If $G(X)$ is connected, then $\text{rank}(X) = n + 1 - \# \text{connected components of } g(X)$.*

PROOF. We will show that the space of functionals constant on X has dimension equal to the number k of connected components of $g(X)$ minus one. Considering X as a matrix with incidence row vectors we first observe that all functionals in the kernel of X must give the same value to all entries ($\mathbf{e}_i$'s) in the same connected component of $g(X)$. That is, they are characterized by k parameters. Thus, the dimension of the space of functionals constant on X is at most $k - 1$.

Consider the matrix X and, as in the proof of [Theorem 62](#), we pick a row as a root of the spanning tree of $G(X)$ and perform row operations so that the resulting matrix is our root row and the rest are ± 1 -vectors. Omitting the root row of X , the orthogonal complement of the remaining rows of X is exactly the k -dimensional linear space described above. Reinserting the root row decreases the dimension of the orthogonal complement by at most one. $\square$

LEMMA 67. *Let $\mathcal{M}$ be a connected matroid of rank r on n elements and $X \subset \text{Inc}(\mathcal{B}_{\mathcal{M}})$ be n incidence vectors of $\mathcal{B}_{\mathcal{M}}$. If $g(X)$ is connected, $\text{rank}(X) = n$. Moreover $G(X)$ is connected and thus $|\det(X)| = r$.*

PROOF. Consider the matrix X and, as in the proof of [Theorem 62](#), we pick representative vectors $\mathbf{v}^{m_1}, \dots, \mathbf{v}^{m_p}$ for each of the components of $G(X)$. Next perform row operations so that the resulting matrix consists of the representative vectors and ± 1 -vectors. Moreover since $g(X)$ is connected these ± 1 -vectors must be a spanning tree of an orientation of $g(X)$, hence there must be $(n - 1)$ ± 1 -vectors. This implies that we have only one representative vector and thus $G(X)$ is connected. We also see that these ± 1 -vectors must be linearly independent. It follows that X must be linearly independent. Then $|\det(X)| = r$ follows from [Corollary 63](#). $\square$

In order to prove a unimodular covering exists we propose another conjecture which implies [Conjecture 58](#):

CONJECTURE 68. *Let $\mathcal{M}$ be a matroid of rank r on n elements. For every $x \in \mathcal{P}_{\mathcal{M}}$ there exists n linearly independent vectors $\mathbf{v}_1, \dots, \mathbf{v}_n \in \text{Inc}(\mathcal{B}_{\mathcal{M}})$ such that $x \in \text{conv}(\mathbf{v}_1, \dots, \mathbf{v}_n)$ and $G(X)$ connected. Hence, $\text{conv}(\mathbf{v}_1, \dots, \mathbf{v}_n)$ is unimodular.*

In support of [Conjecture 57](#), [Conjecture 58](#) and [Conjecture 68](#) we now present some experimental results. We obtained bases description for all matroids of nine elements or less from [\[MRar\]](#). The number of matroids by rank and elements less than or equal to nine is given by the table:

Elements	0	1	2	3	4	5	6	7	8	9
Rank										
0	1	1	1	1	1	1	1	1	1	1
1		1	2	3	4	5	6	7	8	9
2			1	3	7	13	23	37	58	87
3				1	4	13	38	108	325	1275
4					1	5	23	108	940	190214
5						1	6	37	325	190214
6							1	7	58	1275
7								1	8	87
8									1	9
9										1
Total Matroids	1	2	4	8	17	38	98	306	1724	383172

With this data in hand we used TOPCOM [\[Ram02\]](#) to compute triangulations of all the connected matroid polytopes. From these computational experiments we can justify the following theorem:

THEOREM 69. *All 1317 matroid polytopes of connected matroids with eight or less elements have a unimodular triangulation. Moreover, the placing triangulation, determined by the order of bases as presented in [\[MRar\]](#), is a unimodular triangulation.*

Whether the ordering is important for the placing triangulation to be unimodular for these experiments is unclear. As a note, there certainly are non-unimodular triangulations of matroid polytopes. As a natural follow up experiment, we attempted to find as many matroids in the test set that were unimodular as well as triangulated by simplices that had connected graphs $G(X)$.

THEOREM 70. *All 48 matroid polytopes of connected matroids with six or less elements have a triangulation where for each simplex X , $G(X)$ is connected. Hence there exists a unimodular triangulation of each matroid polytope.*

For matroids with seven or more elements, it becomes especially time consuming to find a triangulation with G -connected simplices. We were able to find 36 matroids of rank seven that had G -connected simplices. All matroids and triangulations for [Theorem 69](#) and [Theorem 70](#) can be found at [\[Haw09\]](#).

3.3. Two Dimensional Faces of Matroid Polytopes

In the pursuit of the proof of [Conjecture 57](#), [Conjecture 58](#), and [Conjecture 68](#) related to matroid polytope triangulations, coverings and decompositions, we studied the two dimensional faces of $\mathcal{P}_{\mathcal{M}}$ in order to understand how such subdivisions interact with the low dimensional faces. We obtained a few results.

LEMMA 71 (See Theorem 4.8 in [\[BGW07\]](#)). *Let $\mathcal{M}$ be a matroid and $\mathcal{P}_{\mathcal{M}}$ be its matroid polytope. The 2-dimensional faces of $\mathcal{P}_{\mathcal{M}}$ are equilateral triangles or squares.*




FIGURE 3.2. Square 2-face adjacency structure. $\mathbf{e}_s \neq \mathbf{e}_m$, $\mathbf{e}_t \neq \mathbf{e}_l$, $\mathbf{e}_s \neq \mathbf{e}_t$, $\mathbf{e}_m \neq \mathbf{e}_l$.

LEMMA 72. *Let $\mathcal{M}$ be a matroid of rank r on n elements and F be a 2-dimensional square face of $\mathcal{P}_{\mathcal{M}}$ with vertices $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{w}_4$ such that $(\mathbf{w}_1, \mathbf{w}_2)$, $(\mathbf{w}_2, \mathbf{w}_4)$, $(\mathbf{w}_1, \mathbf{w}_3)$, and $(\mathbf{w}_3, \mathbf{w}_4)$ are the edges. There exists standard unit vectors in $\mathbf{e}_s, \mathbf{e}_t, \mathbf{e}_m, \mathbf{e}_l \in \mathbb{R}^n$ such that $\mathbf{w}_2 = \mathbf{w}_1 + \mathbf{e}_s - \mathbf{e}_t$, $\mathbf{w}_3 = \mathbf{w}_1 + \mathbf{e}_m - \mathbf{e}_l$, and $\mathbf{w}_4 = \mathbf{w}_1 + \mathbf{e}_s - \mathbf{e}_t + \mathbf{e}_m - \mathbf{e}_l$. Moreover, and $\mathbf{e}_s \neq \mathbf{e}_m$, $\mathbf{e}_t \neq \mathbf{e}_l$, $\mathbf{e}_s \neq \mathbf{e}_t$ and $\mathbf{e}_m \neq \mathbf{e}_l$.*

PROOF. Let $\mathcal{M}$ be a matroid and F a 2-face of $\mathcal{P}_{\mathcal{M}}$ with $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{w}_4$ vertices of F such that $(\mathbf{w}_1, \mathbf{w}_2)$, $(\mathbf{w}_2, \mathbf{w}_4)$, $(\mathbf{w}_1, \mathbf{w}_3)$, and $(\mathbf{w}_3, \mathbf{w}_4)$ are the only adjacencies. Due to [Lemma 17](#) we know if $\mathbf{w}_2 = \mathbf{w}_1 + \mathbf{e}_s - \mathbf{e}_t$ and $\mathbf{w}_4 = \mathbf{w}_1 + \mathbf{e}_s - \mathbf{e}_t + \mathbf{e}_m - \mathbf{e}_l$, then $\mathbf{w}_3 = \mathbf{w}_1 + \mathbf{e}_m - \mathbf{e}_l$. Otherwise an additional adjacency (edge) is created among $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{w}_4$, a contradiction to F a 2-face. E.g., if $\mathbf{w}_3 = \mathbf{w}_1 + \mathbf{e}_s - \mathbf{e}_l$ then $\mathbf{w}_3$ and $\mathbf{w}_2$ are adjacent, a contradiction. Then by vector addition $\mathbf{w}_4 = \mathbf{w}_3 + \mathbf{e}_m - \mathbf{e}_l$. Additionally, $\mathbf{e}_s \neq \mathbf{e}_m$, $\mathbf{e}_t \neq \mathbf{e}_l$, $\mathbf{e}_s \neq \mathbf{e}_t$ and $\mathbf{e}_m \neq \mathbf{e}_l$. Otherwise it would create an additional adjacency, hence edge. See [Figure 3.2](#).

□




FIGURE 3.3. Six adjacent vertices restrict any four from being a 2-face.

LEMMA 73. Let $\mathcal{M}$ be a matroid of rank r on n elements, $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{w}_4 \in \mathcal{P}_{\mathcal{M}}$. Let $\mathbf{w}_1$ and $\mathbf{w}_4$ be adjacent to $\mathbf{w}_2$ and $\mathbf{w}_3$ in $\mathcal{P}_{\mathcal{M}}$ where $\mathbf{w}_2 = \mathbf{w}_1 + (\mathbf{e}_s - \mathbf{e}_t)$, $\mathbf{w}_3 = \mathbf{w}_1 + (\mathbf{e}_m - \mathbf{e}_l)$ and $\mathbf{w}_4 = \mathbf{w}_1 + (\mathbf{e}_s - \mathbf{e}_t) + (\mathbf{e}_m - \mathbf{e}_l)$, $\mathbf{e}_s \neq \mathbf{e}_m$, $\mathbf{e}_t \neq \mathbf{e}_l$, $\mathbf{e}_s \neq \mathbf{e}_t$ and $\mathbf{e}_m \neq \mathbf{e}_l$. Then $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3$ and $\mathbf{w}_4$ is a square 2-face if and only if $\mathbf{w}_5 := \mathbf{w}_1 + (\mathbf{e}_s - \mathbf{e}_l)$ or $\mathbf{w}_6 := \mathbf{w}_1 + (\mathbf{e}_m - \mathbf{e}_t)$ are not vertices of $\mathcal{P}_{\mathcal{M}}$.

PROOF. Assume that both $\mathbf{w}_5$ and $\mathbf{w}_6$ are vertices of $\mathcal{P}_{\mathcal{M}}$ and F is a square 2-face with vertices $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3$ and $\mathbf{w}_4$. Let

$$(3.13) \quad f(\mathbf{x}) := a_1x_1 + a_2x_2 + \cdots + a_nx_n = c$$

be a supporting hyperplane of F , i.e. $\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \mathbf{w}_4$ satisfy (3.14). Moreover, for all $z \in \mathcal{P}_{\mathcal{M}} \setminus F$, $f(z) < c$. Since Equation (3.14) is a supporting hyperplane and $\mathbf{w}_5, \mathbf{w}_6 \notin F$, $f(\mathbf{w}_5) < c$ and $f(\mathbf{w}_6) < c$. But $2c = f(\mathbf{w}_2 + \mathbf{w}_3) = f(\mathbf{w}_5 + \mathbf{w}_6) < 2c$, a contradiction.

Assume, without loss of generality, that $\mathbf{w}_5$ is a vertex of $\mathcal{P}_{\mathcal{M}}$, while $\mathbf{w}_6$ is not. We will construct a supporting hyperplane of F . Let

$$(3.14) \quad f(\mathbf{x}) := a_1x_1 + a_2x_2 + \cdots + a_nx_n.$$

Let $a_s = a_t = 1$, $a_m = a_l = 2$, $a_i = 3$ if $i \in \text{supp}(\mathbf{w}_1) \setminus \{s, t, m, l\}$ and $a_j = -1$ if $j \notin \text{supp}(\mathbf{w}_1) \cup \{s, t, m, l\}$. We see that $f(\mathbf{w}_1) = f(\mathbf{w}_2) = f(\mathbf{w}_3) = f(\mathbf{w}_4) = 3(r-2) + 3$, yet $f(\mathbf{w}_5) = 3(r-1) + 2$ and for all vertices $\mathbf{z} \in \mathcal{P}_{\mathcal{M}} \setminus F$, $f(\mathbf{z}) < 3(r-1) + 3$. Hence f is a supporting hyperplane and F is a square 2-face. If $\mathbf{w}_5$ is also not a vertex, then f is still a supporting hyperplane of F . $\square$

COROLLARY 74. *Let $\mathcal{M}$ be a uniform matroid of rank r on n elements. Then the 2-dimensional faces of $\mathcal{P}_{\mathcal{M}}$ are all equilateral triangles.*

PROOF. This follows from Lemma 73 and that if $\mathbf{w}_2 = \mathbf{w}_1 + (\mathbf{e}_s - \mathbf{e}_t)$ and $\mathbf{w}_3 = \mathbf{w}_1 + (\mathbf{e}_m - \mathbf{e}_l)$ are vertices of $\mathcal{P}_{\mathcal{M}}$ then so are $\mathbf{w}_5 := v + (\mathbf{e}_s - \mathbf{e}_l)$ and $\mathbf{w}_6 := v + (\mathbf{e}_m - \mathbf{e}_t)$ since $\mathcal{M}$ is a uniform matroid. $\square$

PROPOSITION 75. *Let $\mathcal{M}$ be a connected matroid of rank r on n elements and v be a vertex of $\mathcal{P}_{\mathcal{M}}$. Then all adjacent vertices of v lie on a $(n-2)$ -hyperplane*

PROOF. Without loss of generality assume $\mathbf{v} = (\underbrace{1, \dots, 1}_r, \underbrace{0, \dots, 0}_{n-r})^\top$. Then by Lemma 17 all vertices $\mathbf{w}$ adjacent to $\mathbf{v}$ lie in the hyperplane $x_1 + \cdots + x_r = r-1$ since $\mathbf{v} - \mathbf{w} = \mathbf{e}_s - \mathbf{e}_l$ where $s \in \{1, \dots, r\}$ and $l \notin \{1, \dots, r\}$. $\square$

In light of Theorem 62 we note that any triangulation of $\{\mathbf{v}, \text{Adj}(\mathbf{v})\}$ is unimodular where $\mathbf{v} \in \text{Inc}(\mathcal{B}_{\mathcal{M}})$. This follows from the fact that all adjacent vertices of $\mathbf{v}$ are in an $n-2$ hyperplane, and thus any full-dimensional simplex X must contain $\mathbf{v}$ which implies $G(X)$ is connected.

CHAPTER 4

Applications to Optimization Through the Structure of Matroid Polytopes

Here we present several heuristics and algorithms for non-linear matroid optimization and convex matroid maximization problems along with computational results. We give a description of our implementation of a speed-up of the algorithm proposed in [Omn03] for convex maximization and two primal heuristics, namely Local Search and Tabu Search, for convex minimization. We also give two heuristics for enumerating a subset of the projected bases.

4.1. Description of the Algorithms and Heuristics

Before we start our description of the algorithms and heuristics, we remark that all of them rely on the geometry of the 1-skeleton graph of the matroid polytope [Sch03]. Recall [Lemma 17](#) from [Chapter 1](#), which states that the edges correspond to bases exchanges. These graphs have a lot of special structure. For example, these graphs are always Hamiltonian [HH72], and it is known that each two-dimensional face of every matroid polytope is either a triangle or a quadrilateral which means the graphs obtained when projecting are quite dense and easy to traverse [BGW07].

Given $\mathbf{v}_1, \dots, \mathbf{v}_k$, we define $\text{cone}(\mathbf{v}_1, \dots, \mathbf{v}_k) := \left\{ \sum_{i=1}^k \lambda_i \mathbf{v}_i \mid \lambda_i \geq 0 \right\}$. If $\mathcal{P} \subseteq \mathbb{R}^n$ is a polytope, we say two vertices $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{P}$ are *adjacent* if they are contained in a one-dimensional face. Next we give a vital but elementary proposition necessary for our algorithms.

PROPOSITION 76. *Let $\mathcal{P} \subseteq \mathbb{R}^n$ be a polytope, $\mathbf{v}_1, \mathbf{v}_2$ vertices of $\mathcal{P}$, and $W \in \mathbb{R}^{d \times n}$. Then*

$$W\mathbf{v}_2 = W\mathbf{v}_1 + \sum_{\mathbf{v} \text{ adjacent to } \mathbf{v}_1} \lambda_{\mathbf{v}} (W\mathbf{v} - W\mathbf{v}_1).$$

PROOF. Let $\tilde{\mathbf{v}}_1, \dots, \tilde{\mathbf{v}}_k$ be adjacent to $\mathbf{v}_1$. It follows from convexity of $\mathcal{P}$ that there exists $\lambda_1, \dots, \lambda_k \geq 0$ such that

$$\begin{aligned} \mathbf{v}_2 &= \lambda_1(\tilde{\mathbf{v}}_1 - \mathbf{v}_1) + \dots + \lambda_k(\tilde{\mathbf{v}}_k - \mathbf{v}_1) + \mathbf{v}_1 \\ \implies W\mathbf{v}_2 &= \lambda_1 W(\tilde{\mathbf{v}}_1 - \mathbf{v}_1) + \dots + \lambda_k W(\tilde{\mathbf{v}}_k - \mathbf{v}_1) + W\mathbf{v}_1. \end{aligned}$$

□

Recall from [Chapter 1](#) the definitions $W\mathcal{P}_{\mathcal{M}} := \{W\mathbf{e}_B \mid B \in \mathcal{B}_{\mathcal{M}}\} \subseteq \mathbb{R}^d$, $\#W\mathcal{P}_{\mathcal{M}}$ to mean the cardinality of $W\mathcal{P}_{\mathcal{M}}$, and $\overline{W\mathcal{P}_{\mathcal{M}}} := \text{conv}(W\mathcal{P}_{\mathcal{M}})$. [Proposition 76](#) implies the following lemma.

LEMMA 77. *Let $\mathcal{P} \subseteq \mathbb{R}^n$ be a polytope, $\mathbf{v}$ a vertex of $\mathcal{P}$, and $W \in \mathbb{R}^{d \times n}$. Then $W\mathbf{v} \in \text{relint}(W\mathcal{P})$ if and only if $\text{cone}(W\tilde{\mathbf{v}} - W\mathbf{v} \mid \tilde{\mathbf{v}} \text{ adjacent to } \mathbf{v})$.*

Define $\text{Adj}(B) := \{\tilde{B} \in \mathcal{B}_{\mathcal{M}} \mid \mathbf{e}_{\tilde{B}} - \mathbf{e}_B = \mathbf{e}_i - \mathbf{e}_j\}$. Using [Lemma 17](#) we get the following corollaries of [Proposition 76](#) and [Lemma 77](#).

COROLLARY 78. *Let $\mathcal{M}$ be a matroid on the ground set $[n]$, $W \in \mathbb{R}^{d \times n}$, $B, \tilde{B} \in \mathcal{B}_{\mathcal{M}}$.*

$$W\mathbf{e}_{\tilde{B}} = W\mathbf{e}_B + \sum_{B' \in \text{Adj}(B)} \lambda_{B'}(W\mathbf{e}_{B'} - W\mathbf{e}_B)$$

where $\lambda_{B'} \geq 0$.

COROLLARY 79. *Let $\mathcal{M}$ be a matroid on the ground set $[n]$ and $W \in \mathbb{R}^{d \times n}$ and $B \in \mathcal{B}_{\mathcal{M}}$. $W\mathbf{e}_B \in \text{relint}(\overline{W\mathcal{P}(\mathcal{M})})$ if and only if $\text{cone}(W\mathbf{e}_{B'} - W\mathbf{e}_B \mid B' \in \text{Adj}(B)) = \mathbb{R}^d$.*

In order to prove that some of our heuristics are efficient, it is necessary to consider the *generalized unary encoding* of the weightings $W \in \mathbb{R}^{d \times n}$ [[BLOW08](#)]. We consider weightings $W_{i,j}$ of the form

$$W_{i,j} = \sum_{k=1}^p \delta_{i,j}^k a_k$$

where $p \geq 1$ is a fixed integer, the distinct positive integers a_k are binary encoded, and the integers $\delta_{i,j}^k$ are unary encoded. We can think of $\delta^k := ((\delta_{i,j}^k)) \in \mathbb{Z}^{d \times n}$ for $1 \leq k \leq p$ as unary-encoded matrices, with binary encoded $\mathbf{a} \in \mathbb{Z}^p$ where $a_k > 0$ for $1 \leq k \leq p$. Then we have $W := ((W_{i,j})) = \sum_{k=1}^p a_k \delta^k$. The generalized unary encoding includes three special cases:

1. **Unary-encoded weights:** With $p = 1$ and $a_1 = 1$, we get the ordinary model of unary-encoded $W = \delta^1$.
2. **Binary-encoded $\{0, a_1, \dots, a_p\}$ -valued weights:** With $\delta^k \geq 0$ for all k , and $\sum_{k=1}^p \delta_{i,j}^k \leq 1$, for all i, j , we get the case of all $W_{i,j}$ in the set $\{0, a_1, \dots, a_p\}$ having binary-encoded elements.
3. **$0 \setminus 1$ -valued weights:** This is, of course, the important common special case of 1 and 2.

With W encoded in this way it is possible to bound the size of $\#W\mathcal{P}_{\mathcal{M}}$.

LEMMA 80. *Let d and p be fixed, $\mathcal{M}$ a matroid on n elements, and a generalized unary encoded matrix $W := ((W_{i,j})) = \sum_{k=1}^p a_k \delta^k \in \mathbb{Z}^{d \times n}$. Then $\#W\mathcal{P}_{\mathcal{M}}$ is polynomially bounded in n , the binary encoding of a_k , and the unary encoding of δ^k , for $1 \leq k \leq p$.*

PROOF. Let $\beta_k = \max_{i,j} \delta_{i,j}^k$, for $1 \leq k \leq p$. Then

$$\begin{aligned} W\mathcal{P}_{\mathcal{M}} &= \{ W\mathbf{e}_B \mid B \in \mathcal{B}_{\mathcal{M}} \} \\ &\subseteq \left\{ \sum_{k=1}^p \lambda_k a_k \mid \lambda \in \{0, 1, \dots, \beta_k \text{rank}(\mathcal{M})\}^p, \sum_{k=1}^p \lambda_k = \text{rank}(\mathcal{M}) \right\}^d \end{aligned}$$

□

The previous lemma bounds the size of $\#W\mathcal{P}_{\mathcal{M}}$, yet the number of integer points in the smallest rectangular region containing $W\mathcal{P}_{\mathcal{M}}$ may be exponential in the input. Some of our heuristics depend on the number of integers points in the smallest rectangular region containing $W\mathcal{P}_{\mathcal{M}}$ to be polynomially bounded in order to prove efficiency.

LEMMA 81 ([BLOW08]). *Let d and p be fixed, $\mathcal{M}$ a matroid on n elements, and $W \in \mathbb{Z}^{d \times n}$. Then $\#\overline{W\mathcal{P}_{\mathcal{M}}} \cap \mathbb{Z}^d$ is polynomially bounded in n , and the unary encoding of W , for $1 \leq k \leq p$.*

PROOF. Then

$$\begin{aligned} W\mathcal{P}_{\mathcal{M}} &= \{ W\mathbf{e}_B \mid B \in \mathcal{B}_{\mathcal{M}} \} \\ &\subseteq \left\{ \sum_{k=1}^p \lambda_k \mid \lambda \in \{0, 1, \dots, \beta_k \text{rank}(\mathcal{M})\}^p, \sum_{k=1}^p \lambda_k = \text{rank}(\mathcal{M}) \right\}^d. \end{aligned}$$

□

4.1.1. Local and Tabu Search. First we present **Heuristic 82** that starts at any base of $\mathcal{M}$ and proceeds by pivoting to its neighbors on $\mathcal{P}(\mathcal{M})$ as long as the pivot decreases the given function f through the weighting W .

In the majority of our experiments f is concave or convex. In the case where f is concave, there will be a minimum on the boundary of $\overline{W\mathcal{P}_{\mathcal{M}}}$. If f is convex, then the minima may be in the interior of $\overline{W\mathcal{P}_{\mathcal{M}}}$. We emphasize that **Heuristic 82** is not guaranteed to terminate at an optimum.

HEURISTIC 82 (Local Search).

LS($\mathcal{M}, W, f, B$)

Input: Matroid $\mathcal{M}$ on n elements, $B \in \mathcal{B}_{\mathcal{M}}$, $W \in \mathbb{R}^{d \times n}$, $f : \mathbb{R}^d \rightarrow \mathbb{R}$.

Output: A base $B' \in \mathcal{B}_{\mathcal{M}}$ such that $f(W\mathbf{e}_{\tilde{B}}) \geq f(W\mathbf{e}_{B'}) \forall \tilde{B} \in \text{Adj}(B')$

```

1:  $B' := B$ 
2: repeat
3:    $OLDB' := B'$ 
4:   if  $f(W\mathbf{e}_{\tilde{B}}) < f(W\mathbf{e}_{B'})$  for some  $\tilde{B} \in \text{Adj}(B')$  then
5:      $B' := \tilde{B}$ 
6: until  $B' = OLDB'$ 
7: return  $B'$ 

```

If the objective function f is linear, then **Heuristic 82** follows the non-degenerate steps of the simplex method. The following is an application of the well-known fact that the simplex method with a linear objective will terminate at a minimum [Sch86b].

LEMMA 83. *Let $\mathcal{M}$ be a matroid on $[n]$, $B \in \mathcal{B}_{\mathcal{M}}$, $W \in \mathbb{R}^{d \times n}$, and $f(\mathbf{x}) = \mathbf{c} \cdot \mathbf{x}$, where $\mathbf{c} \in \mathbb{R}^d$. LS($\mathcal{M}, W, g, B$) terminates at $\hat{B}$ where $g(W\mathbf{e}_{\hat{B}}) = \min\{g(W\mathbf{e}_B) \mid B \in \mathcal{B}_{\mathcal{M}}\}$.*

Lemma 83 is used by MOCHA in the implementation of our heuristics. For example, we can easily compute a tight rectangular region containing $W\mathcal{P}$. We also use **Lemma 83** to obtain an integer point on the boundary of $W\mathcal{P}_{\mathcal{M}}$, a prerequisite for some of our heuristics.

In [section 4.3](#), we will show the practical limitations of [Heuristic 82](#) and we give a theoretical bound on the running time of the previous algorithm given restrictions on W and d .

LEMMA 84. *For every fixed d and p , $\mathcal{M}$ a matroid on $[n]$ given by an independence oracle, a generalized unary encoded matrix $W := ((W_{i,j})) = \sum_{k=1}^p a_k \delta^k \in \mathbb{Z}^{d \times n}$, and f given by comparison oracle, then [Heuristic 82](#) terminates in time polynomial in n , the binary encoding of a_k , and the unary encoding of δ^k , for $1 \leq k \leq p$.*

PROOF. For any $B \in \mathcal{B}_{\mathcal{M}}$ we can enumerate $\text{Adj}(B)$ in polynomial time via an oracle for $\mathcal{M}$ by testing which of $\{B \setminus \{i\} \cup \{j\} \mid i \in B, j \in [n] \setminus B\}$ are bases. Thus, we can pivot in polynomial time in the input. Since [Heuristic 82](#) always pivots to an adjacent base that is smaller, through f and W , than the current base, each point of $W\mathcal{P}_{\mathcal{M}}$ will be visited at most once. By [Lemma 80](#), $\#W\mathcal{P}_{\mathcal{M}} \cap \mathbb{Z}^d$ is polynomially bounded in the input. Also, the arithmetic in [Heuristic 82](#) is polynomial in n and the binary encoding of a_k . $\square$

Tabu search was first presented in [[Glo86](#)] and has been widely used in combinatorial optimization. It begins at base $B \in \mathcal{B}_{\mathcal{M}}$ and pivots to an adjacent base B' if it is smaller than some other adjacent base, through f and the weighting W . This differs from [Heuristic 82](#) in that we allow pivots that are not necessarily smaller than the current base, through f and the weighting W . We record the smallest value $f(W\mathbf{e}_B)$ encountered and terminate after L pivots with no update to the minimum encountered. As for [Heuristic 82](#), for the majority of our experiments f is concave or convex. We emphasize that [Heuristic 85](#) is not guaranteed to terminate at the optimum for general f .

HEURISTIC 85 (Tabu Search).

TS($\mathcal{M}, B, W, L$)

Input: Matroid $\mathcal{M}$ on $[n]$, $B \in \mathcal{B}_{\mathcal{M}}$, $W \in \mathbb{R}^{d \times n}$, $L \in \mathbb{N}$.

Output: $B' \in \mathcal{B}_{\mathcal{M}}$

- 1: $B' := B$
- 2: VIS $:= \{B\}$
- 3: CURMIN $:= f(W\mathbf{e}_{B'})$.

```

4: repeat
5:   if  $f(\mathbf{We}_{\tilde{B}}) < f(\mathbf{We}_{\hat{B}})$  for some  $\tilde{B}, \hat{B} \in \text{Adj}(B') \setminus \text{VIS}$  then
6:      $\text{VIS} := \text{VIS} \cup \{\tilde{B}\}$ 
7:      $B' := \tilde{B}$ 
8:   if  $f(\mathbf{We}_{B'}) < \text{CURMIN}$  then
9:      $\text{CURMIN} := f(\mathbf{We}_{B'})$ .
10: until  $L$  pivots with CURMIN unchanged.
11: return  $B'$ 

```

Heuristic 85 can also be modified such that we mark $\mathbf{We}_B$ as visited, not B , and only pivot to B such that $\mathbf{We}_B$ is unvisited. In this way, we could also prove that the running time of the modified algorithm is polynomial in the input, given the same restrictions for **Lemma 87**. Although, restricting to visiting new $\mathbf{We}_B$ could limit the possibilities of quickly converging to the optimum though.

We observed through our computational experiments that **Heuristic 82** and **Heuristic 85** work very well when optimizing concave or convex functions f over projected matroid polytopes as we will show in **section 4.3**. However, when f is arbitrary then the previous algorithms may not reach the optimum.

4.1.2. Listing Heuristics. In the remainder of the present section, we describe algorithms and heuristics aimed at tackling problems with an arbitrary balancing function f or using Pareto or minmax optimization. The following algorithms do not explicitly evaluate f but are presented as *listing algorithms*. Note that if we have all of the projected matroid bases then it is simple to extract the Pareto optima through a straightforward pairwise comparison. Likewise for optimizing over f or with a min-max objective, we simply evaluate over all projected bases found. This is a similar methodology as [BLOW08] where the authors prove efficient deterministic algorithms, given sufficient conditions on the input, to list all projected bases.

In [BLOW08] the authors used matroid intersections to solve the problem: Given $x \in \mathbb{R}^d$ find $B \in \mathcal{B}_M$ such that $\mathbf{We}_B = x$ if such a B exists. Guided by the success of our previous

heuristics, we had the new idea of using either **Heuristic 85** or **Heuristic 82** to find a $B \in \mathcal{B}_{\mathcal{M}}$ such that $W\mathbf{e}_B = \mathbf{x}$. The novelty of our heuristic is that we use fast convex optimization as a subroutine to solve non-linear and Pareto optimization problems.

Given an $\mathbf{x}' \in \mathbb{Z}^d$, **Heuristic 86** forms a convex function $f_{\mathbf{x}'}$ that minimizes at $\mathbf{x}'$ and $f_{\mathbf{x}'}(\mathbf{x}') = 0$. Our heuristic calls **Heuristic 82** or **Heuristic 85** with the function $f_{\mathbf{x}'}$ and if it returns a base $B \in \mathcal{B}_{\mathcal{M}}$ such that $f_{\mathbf{x}'}(W\mathbf{e}_B) = 0$ then $W\mathbf{e}_B = \mathbf{x}$.

HEURISTIC 86 (Pivot Test).

PT($\mathcal{M}, B, W, t, S$)

Input: Matroid $\mathcal{M}$ on n elements, $B \in \mathcal{B}_{\mathcal{M}}$, $W \in \mathbb{R}^{d \times n}$, $t \in \mathbb{N}$, finite set $S \subseteq \mathbb{Z}^d$.

Output: $\text{PT} \subseteq \mathcal{B}_{\mathcal{M}}$ such that $\{W\mathbf{e}_B \mid B \in \text{PT}\} \subseteq S$.

```

1: PT :=  $\emptyset$ 
2: for each  $\mathbf{x}' \in S$  do
3:    $f_{\mathbf{x}'}(\mathbf{x}) := \sum_{i=1}^d (x_i - x'_i)^2$ 
4:   for  $1, \dots, t$  do
5:      $B' :=$  random basis of  $\mathcal{M}$ 
6:      $B := \text{LS}(\mathcal{M}, W, f_{\mathbf{x}'}, B')$  (Or use Heuristic 85)
7:     if  $f(W\mathbf{e}_B) = 0$  then
8:       PT := PT  $\cup$   $B$ 
9:     Break for loop
10: return PT

```

LEMMA 87. *For every fixed d and p , $\mathcal{M}$ a matroid on $[n]$ given by an independence oracle, a generalized unary encoded matrix $W := ((W_{i,j})) = \sum_{k=1}^p a_k \delta^k \in \mathbb{Z}^{d \times n}$, a finite set $S \subset \mathbb{Z}^d$, then **Heuristic 86** terminates in time polynomial in n , t , the size of S , the binary encoding of a_k , and the unary encoding of δ^k , for $1 \leq k \leq p$.*

PROOF. This follows from the fact that **Heuristic 82** and **Heuristic 85** are polynomial under these assumptions. Moreover, we call **Heuristic 82** or **Heuristic 85** at most $t|S|$ times. $\square$

Typically we wish to use **Heuristic 86** to try to enumerate a subset of $W_{\mathcal{P}}$. Thus, if we further assume W is unary encoded, then the smallest box containing $\overline{W\mathcal{P}_{\mathcal{M}}}$ is polynomial which we use as our input S in **Heuristic 86**.

For general non-linear multi-criteria matroid optimization problems, when the function f we are minimizing is concave, then some optimum will be a vertex of $\overline{W\mathcal{P}_{\mathcal{M}}}$. Thus, in such a case, it is sufficient to enumerate the vertices of $\overline{W\mathcal{P}_{\mathcal{M}}}$ and test f over those points. Furthermore, some vertices are also Pareto optimal, but in general, the vertices are not all the Pareto optima. Later we will give an algorithm that uses the vertices of $W\mathcal{P}_{\mathcal{M}}$ to facilitate finding other Pareto optima. We will prove that the following algorithm will enumerate all vertices of $\overline{W\mathcal{P}_{\mathcal{M}}}$ and possibly other integral points on the boundary of $\overline{W\mathcal{P}_{\mathcal{M}}}$. **Algorithm 88** starts at a base $B \in \mathcal{B}_{\mathcal{M}}$ such that $W\mathbf{e}_B$ is on the boundary of $\overline{W\mathcal{P}_{\mathcal{M}}}$. If $B' \in \text{Adj}(B)$, $W\mathbf{e}_{B'}$ is a newly seen point and is on the boundary of $\overline{W\mathcal{P}_{\mathcal{M}}}$, we record B' and $W\mathbf{e}_{B'}$, pivot to B' , and continue.

ALGORITHM 88 (Projected Boundary).

PB($\mathcal{M}, W, B$)

Input: Matroid $\mathcal{M}$ on n elements, $W \in \mathbb{R}^{d \times n}$, $B \in \mathcal{B}_{\mathcal{M}}$ such that $W\mathbf{e}_B$ is an extreme point of $\overline{W\mathcal{P}_{\mathcal{M}}}$.

Output: $\text{CH} \subseteq \mathcal{B}_{\mathcal{M}}$ such that $\{W\mathbf{e}_B \mid B \in \text{CH}\} \supseteq \text{vert}(\overline{W\mathcal{P}_{\mathcal{M}}})$.

- 1: $\text{CH} := \{B'\}$
- 2: Mark B' as unvisited in CH
- 3: $\text{PB} := \{W\mathbf{e}_{B'}\}$
- 4: **repeat**
- 5: $B' :=$ first unvisited base in CH
- 6: Mark B' as visited.
- 7: **for** $\hat{B} \in \text{Adj}(B')$ **do**
- 8: **if** $W\mathbf{e}_{\hat{B}}$ is an extreme point of $\overline{W\mathcal{P}_{\mathcal{M}}}$ **then**
- 9: **if** $W\mathbf{e}_{\hat{B}} \notin \text{PB}$ **then**
- 10: $\text{CH} := \text{CH} \cup \{\hat{B}\}$

```

11:      Mark  $\hat{B}$  as unvisited.
12:       $PB := PB \cup \{W\mathbf{e}_{\hat{B}}\}$ 
13:  until  $PB$  unchanged.
14:  return  $CH$ 

```

In [section 4.3](#) we will see that [Algorithm 88](#) works well. We believe this is due to the following: the projected bases are often highly clustered and with much fewer projected bases near the boundary of $\overline{WP_{\mathcal{M}}}$. To see how large $W^{-1}\mathbf{x}$, for $\mathbf{x} \in \mathbb{R}^d$, can be, see Figure 1 in [\[GLM08\]](#). The following lemma proves that the output of [Algorithm 88](#) will contain all the vertices of the convex hull of $WP_{\mathcal{M}}$.

LEMMA 89. *Let $\mathcal{M}$ be a matroid on $[n]$, $W \in \mathbb{R}^{d \times n}$, $B \in \mathcal{B}_{\mathcal{M}}$ such that $W\mathbf{e}_B$ is an extremal point of $\overline{WP_{\mathcal{M}}}$. Then*

$$\text{vert}(\overline{WP_{\mathcal{M}}}) \subseteq W(PB(\mathcal{M}, W, B)) \subseteq (WP_{\mathcal{M}} \cap \partial \overline{WP_{\mathcal{M}}}).$$

PROOF. Let $B', \hat{B} \in \mathcal{B}_{\mathcal{M}}$ where $W\mathbf{e}_{B'}$ is an extremal point and $W\mathbf{e}_{\hat{B}}$ is a vertex of $\overline{WP_{\mathcal{M}}}$. Moreover let $W\mathbf{e}_{B'}$ and $W\mathbf{e}_{\hat{B}}$ be on a 1-face F of $\overline{WP_{\mathcal{M}}}$. By [Corollary 78](#) there exists $B'' \in \text{Adj}(B')$ such that either $W\mathbf{e}_{B''} = W\mathbf{e}_{\hat{B}}$ or $W\mathbf{e}_{B''}$ is a positive convex combination of $W\mathbf{e}_{B'}$ and $W\mathbf{e}_{\hat{B}}$, i.e. $W\mathbf{e}_{B''} \in F$. Thus $W\mathbf{e}_{B''}$ is extremal and [Algorithm 88](#) can always pivot towards every vertex of $\overline{WP_{\mathcal{M}}}$ and will output all vertices of $\overline{WP_{\mathcal{M}}}$. $\square$




FIGURE 4.1. A pivot from B' to B'' towards $\hat{B}$ in [Algorithm 88](#).

In [\[Onn03\]](#) it was shown that the number of vertices of $WP_{\mathcal{M}}$ is polynomially bounded when the number of weightings d is fixed (see Proposition 2.1 and Lemma 2.3). Moreover the author showed that the vertices of $WP_{\mathcal{M}}$ can be found in polynomial time. A similar

result can be found in [BLOW08]. **Algorithm 88** can (and does in practice) pick up non-vertex integral boundary points. Hence, a bound on the number of vertices is not sufficient to guarantee efficiency of the algorithm.

LEMMA 90. *For fixed d and p , $\mathcal{M}$ a matroid on $[n]$ given by an independence oracle, a generalized unary encoded matrix $W := ((W_{i,j})) = \sum_{k=1}^p a_k \delta^k \in \mathbb{Z}^{d \times n}$, **Algorithm 88** runs in time polynomial in n , the binary encoding of a_k , and the unary encoding of δ^k , for $1 \leq k \leq p$.*

PROOF. Since $\#W\mathcal{P}_{\mathcal{M}} \cap \mathbb{Z}^d$ is polynomially bounded and $\#\text{Adj}(B) \leq n^2$ we only have to show that deciding if $W\mathbf{e}_{\tilde{B}}$ is an extremal point of $\overline{W\mathcal{P}_{\mathcal{M}}}$ is polynomial in the input. This is equivalent to deciding if $\text{cone}\{W\mathbf{e}_B - W\mathbf{e}_{\tilde{B}} \mid B \in \text{Adj}(\tilde{B})\}$ is pointed. We can decide this by solving the feasibility problem:

$$\left\{ \mathbf{x}, -\mathbf{x} \in \text{cone}\{W\mathbf{e}_B - W\mathbf{e}_{\tilde{B}} \mid B \in \text{Adj}(\tilde{B})\} \mid \mathbf{x} > 0 \right\}$$

□

We will show in **section 4.3** the effectiveness of **Heuristic 86** (PT) in enumerating all projected bases. When finding Pareto optima though it is not necessary to enumerate *all* projected bases. The following heuristic combines **Algorithm 88** and **Heuristic 86** to find a set of points which could be Pareto optima of $W\mathcal{P}_{\mathcal{M}}$. We observe that the min-max optimum is also a Pareto optima. Hence **Algorithm 88** and **Heuristic 86** could return the min-max optimum of $W\mathcal{P}_{\mathcal{M}}$. The main idea is that if we have the boundary points, then the regions where other Pareto optima lay is determined by the convex hulls of points derived from a triangulation of the boundary.

HEURISTIC 91 (Boundary and Triangular Region Pareto Test).

BTRPT($\mathcal{M}, W, t$)

Input: Matroid $\mathcal{M}$ on n elements, $W \in \mathbb{R}^{d \times n}$, $t \in \mathbb{N}$.

Output: $\text{PO} \subseteq \mathcal{B}_{\mathcal{M}}$

1: $\text{PO} := \text{PB}(\mathcal{M}, W)$

2: **for** $B \in \text{PO}$ **do**

```

3:  for  $B' \in \text{PO} \setminus B$  do
4:    if  $\text{We}_{B'} \geq_{\text{Pareto}} \text{We}_B$  then
5:       $\text{PO} := \text{PO} \setminus B'$ 
6:  $\text{PPO} := \text{WPO}$  projected on the hyperplane  $H := \{x \mid \sum x_i = 1\}$ .
7: Triangulate  $\text{PPO}$ .
8: for each facet  $F$  of  $\text{PPO}$  do
9:    $\text{TRIPOINTS} := \{\text{Bases in PO corresponding to vertices of } F\}$ 
10:  for  $i = 1, \dots, d$  do
11:     $X_i^{\max} := \max(\text{We}_B)_i \mid B \in \text{TRIPOINTS}$ 
12:   $\text{PO} := \text{PT}(\mathcal{M}, B, W, t, \text{conv}(\text{TRIPOINTS} \cup X^{\max})) \cup \text{PO}$ 
13: for  $B \in \text{PO}$  do
14:  for  $B' \in \text{PO} \setminus B$  do
15:    if  $\text{We}_{B'} \geq_{\text{Pareto}} \text{We}_B$  then
16:       $\text{PO} := \text{PO} \setminus B'$ 
17: return  $\text{PO}$ 

```




FIGURE 4.2. Pareto optima boundary points and blue regions where other Pareto optima may lay.

Except for the boundary points, **Heuristic 91** is not guaranteed to return points that are Pareto optima since **Heuristic 86** (PT) may miss some projected bases in the test regions.

Motivated by the need of optimizing any function f or exploring *all* Pareto or minmax optima, we devised a variation of the breadth-first search algorithm that would limit the search to some depth of the tree. Our intuition was that the graph of base exchanges is highly connected, thus small depth was all that is necessary for listing a large proportion of all projected bases, as we will see in [section 4.3.](#):

HEURISTIC 92 (Different Fiber BFS).

DFBFS($\mathcal{M}, W, B, d, l, \text{PB}$)

Input: Matroid $\mathcal{M}$ on n elements, $W \in \mathbb{R}^{d \times n}$, $B \in \mathcal{B}_{\mathcal{M}}$, $d, l \in \mathbb{N}$, $\text{PB} \subseteq \mathcal{WP}_{\mathcal{M}}$.

Output: $\text{DFBFS} \subseteq \{W\mathbf{e}_B \mid B \in \mathcal{B}_{\mathcal{M}}\}$

```

1: if  $l \geq d$  then
2:   return  $\emptyset$ 
3:  $\text{PB} := \text{PB} \cup \{W\mathbf{e}_B\}$ 
4:  $\text{PTT} := \emptyset$ 
5: for each  $B' \in \text{Adj}(B)$  do
6:   if  $W\mathbf{e}_{B'} \notin \text{PB}$  then
7:     if  $W\mathbf{e}_{B'} \neq W\mathbf{e}_B$  then
8:        $\text{PB} := \text{PB} \cup W\mathbf{e}_{B'}$ 
9:        $\text{PTT} := \text{PTT} \cup B'$ 
10: for each  $B' \in \text{PTT}$  do
11:    $\text{PB} := \text{PB} \cup \text{DFBFS}(\mathcal{M}, W, B', d, l + 1, \text{PB})$ .
12: return  $\text{PB}$ 

```

Heuristic 92 begins at a base $B \in \mathcal{B}_{\mathcal{M}}$. It adds the projected point $W\mathbf{e}_B$ to the set PB , short for projected bases. It then enumerates all adjacent bases $\text{Adj}(B)$ of B . If a neighbor B' 's projected point $W\mathbf{e}_{B'} \notin \text{PB}$ and $W\mathbf{e}_B \neq W\mathbf{e}_{B'}$, then we add $W\mathbf{e}_{B'}$ to PB and recursively call DFBFS on B' . We allow a parameter d which determines the recursive depth allowed. **Heuristic 92** takes its name, different fiber breadth-first-search, from the fact that we do not allow pivots that evaluate the same under the weighting W . As a small consequence,

this guarantees that the number of times DFBFS is called is bounded by $\#\mathcal{WP}_{\mathcal{M}}$ a useful fact if the input is bounded as in [Lemma 80](#).

4.2. Software Implementation

Our heuristics are implemented in C++ and take advantage of the object-oriented paradigm. The software MOCHA¹[\[DHLO09\]](#) reads in either a vectorial matroid, represented by an $m \times n$ floating-point matrix or a graphical matroid, represented by an $n \times n$ adjacency matrix. The weightings are read in as a $d \times n$ floating point matrix.

For vectorial matroids, the rank of a subset of columns is computed using LAPACK, a standard and robust linear algebra package [\[ABB⁺99\]](#). For vectorial matroids with elements in $\mathbb{Z}$, MOCHA has the option of using the GMP arbitrary-precision software to perform Gaussian elimination using exact arithmetic [\[Gea09\]](#). Enumeration of the neighbors of a base B is done by calculating the rank of $B \setminus i \cup j$ for all $i \in B$ and $j \in [n] \setminus B$. If it is full rank then it is returned as a neighboring base. Random bases are determined by randomly choosing a $\text{rank}(\mathcal{M})$ sized subset $A \subseteq [n]$ and checking if $\text{rank}(A) = \text{rank}(\mathcal{M})$, repeating until such an A is found.

For graphical matroids, the rank of the matroid, and any $A \subseteq [n]$, is determined by calculating the size of a spanning forest via breadth-first-search. To enumerate the neighbors of $B \in \mathcal{B}_{\mathcal{M}}$ we first calculate all paths in B using dynamic programming. Adding any element (edge) $j \notin B$ to B will create a cycle C . We use our pre-calculated all paths of B to quickly determine C . Then all subsets $B \setminus i \cup j$ where $i \in C$ will be an adjacent base to B . This may not be the optimal solution to graphical matroid adjacency enumeration, but it is straight-forward.

The Projected Boundary algorithm ([Algorithm 88](#)) is only implemented for $d = 2$. This is due to the computational expense of determining if $W\mathbf{e}_{\hat{B}}$ is an extreme point of $\overline{\mathcal{WP}_{\mathcal{M}}}$ on line 8. When $d = 2$ this is easy to check by sorting the projected rays between $\hat{B}$ and its neighbors. If there exists two sorted rays with angle larger than π then $W\mathbf{e}_{\hat{B}}$ is extreme.

For correctness, we compare the number of projected spanning trees found using our methods versus the real total number of projected spanning trees. We used an algorithm for generating all of the spanning trees in undirected graphs presented by Matsui [\[Mat97\]](#). The

¹Matroid Optimization Combinatorial Heuristics and Algorithms

algorithm requires $O(n + m + \tau n)$ time when the given graph has n vertices, m edges, and τ spanning trees. For outputting all of the spanning trees explicitly, this time complexity is optimal. We also implemented the asymptotic $0 \setminus 1$ polytope vertex-estimation presented in [BS07a]. This gives us the ability to estimate the number of bases of matroid polytopes in order to better understand the proportion of bases to projected bases for problems where full enumeration is intractable.

4.3. Computational Results

Now we present our experiments for which we used six roughly comparable machines (see Table 4.38). For our tests, **Heuristic 82** uses the pivot rule that $f(W\mathbf{e}_{\tilde{B}}) < f(W\mathbf{e}_{B'})$ for all $\tilde{B} \in \text{Adj}(B')$ in line 4. **Heuristic 85** uses the pivot rule that $f(W\mathbf{e}_{\tilde{B}}) < f(W\mathbf{e}_{\hat{B}})$ for all $\tilde{B}, \hat{B} \in \text{Adj}(B') \setminus \text{VIS}$ in line 5.

4.3.1. Calibration Set. Our first goal was to perform experiments on matroids for which we can compute all bases in order to better understand our heuristics and algorithms. We generated fifteen connected random graphs: *gn9e18*, *gn9e27*, *gn10e22*, *gn10e28*, *gn10e33*, *gn11e13*, *gn11e20*, *gn11e27*, *gn11e41*, *gn12e16*, *gn12e24*, *gn12e33*, *gn13e19*, *gn13e29*, *gn13e39* which we will refer to as our *calibration set*. The names of our graphs follow the simple nomenclature *gn[#nodes]e[#edges]*. We consider two, three and five criteria, i.e. number of weightings. We further consider three different ranges of integral weights for each criteria. For the calibration set we adopt the following nomenclature

$$gn[\text{\#nodes}]e[\text{\#edges}]d[\text{\#criteria}]w[\text{low weight}]w[\text{high weight}]$$

where we generated random integral weightings between [low weight] and [high weight]. First we simply compare the number of spanning trees of our calibration set to the number of projected spanning trees.

- **Table 4.1** shows the calibration set with two weightings (criteria) and integral weights 0 – 20, 0 – 100, and 0 – 1000.
- **Table 4.2** shows the calibration set with three weightings (criteria) and integral weights 0 – 20, 0 – 100, and 0 – 1000.
- **Table 4.3** contains the calibration set with five weightings (criteria) and integral weights 0 – 1, 0 – 2, and 0 – 5.

We give the exact number of projected spanning trees and compare versus the exact number of spanning trees.

4.3.1.1. *Calibration Set - Heuristic 92.* There are four parameters to our implementation of **Heuristic 92** (DFBFS): number of searches N , BFS depth, boundary retry limit and random retry limit. First we attempt to find a new boundary projected bases using **Heuristic 82** (Local Search) and a random direction. For every new projected base, we run **Heuristic 92** (DFBFS) with the given depth parameter. We attempt to find N new boundary projected bases and give up if we exceed the boundary retry limit. Next our algorithm will generate a random base and project it by the weighting. If it is a new projected base we run **Heuristic 92** (DFBFS) with the given depth parameter. We attempt to find N new random projected bases and give up if we exceed the random retry limit.

- **Table 4.4** shows **Heuristic 92** (DFBFS) on the calibration set in two criteria.
- **Table 4.5** shows **Heuristic 92** (DFBFS) on the calibration set in three criteria.
- **Table 4.6** shows **Heuristic 92** (DFBFS) on the calibration set in five criteria.

We list the machine used, seconds required, number of searches, BFS depth, boundary retry limit and the random retry limit. We noticed a decrease in the proportion of projected trees found to all projected trees as the dimension and weights increased. It is also important to note, as seen in **Table 4.1** and **Table 4.2**, as the dimension and weights increase, the proportion of all projected trees to all spanning trees increases.

In our experiments, in almost all cases, **Heuristic 92** (DFBFS) terminated by exceeding the boundary and random retry limit and not the number of searches. This can be attributed to the phenomenon that bases are projected in a relatively tight area, with few projected bases near the boundary. Other conditions for picking the initial base for **Heuristic 92** (DFBFS) could yield better results. For example, keeping a random leaf of the previous truncated BFS. Or we could compute the boundary of the projected bases and attempt to find a new initial base using **Heuristic 86** (PT) on the “holes” in the convex hull.

4.3.1.2. *Calibration Set - Heuristic 86.* Our implementation of **Heuristic 86** (PT) uses **Heuristic 82** (LS) with the $2d$ directions $\{\pm \mathbf{e}_i \mid i \in \{1, \dots, d\}\}$ to compute a bounding box containing all projected base.




FIGURE 4.3. 96.18% projected bases of *gn11e41d2w0w100* using **Heuristic 92** (DFBFS)

- **Table 4.7** shows the result of **Heuristic 86** (PT) on our calibration test set with two criteria and weights 0 – 20 and 0 – 100 using **Heuristic 82** (LS) as our test method subroutine. We give the proportion of projected trees found versus all projected trees. Naturally with larger weight values, the projected trees will be contained in a larger box, requiring more time.
- **Table 4.8** shows the result of **Heuristic 86** (PT) on our calibration test set with two criteria and weights 0 – 20 using **Heuristic 85** (TS) as our test method subroutine. The run times using **Heuristic 85** (TS) are longer than **Heuristic 82** (LS), but we find nearly all the projected trees. The worse case is the sparse graph *gn11e13d2w0w20.mo* where we find only 90.24% of all projected trees.

Heuristic 86 (PT) has two major advantages;

- **Heuristic 86** uses very little memory since **Heuristic 85** (TS) and **Heuristic 82** (LS) use little memory;
- **Heuristic 86** can be distributed since one can partition up test regions and run each region as a separate instance.

4.3.1.3. *Calibration Set - Algorithm 88.* Table 4.9 shows the number of projected spanning trees found on the boundary using Algorithm 88 (PB). All computations took less than one second. To find a starting base, Algorithm 88 (PB) generates a random direction and calls Heuristic 82 (LS). It is interesting to note that the graph *gn13e39d2w0w1000* has 15,037,589 projected spanning trees, yet there are only 26 points found by Algorithm 88 (PB).

4.3.1.4. *Calibration Set - Heuristic 91.* Table 4.10 shows the results of Heuristic 91 (BTRPT) on the calibration set with weightings 0 – 20 and 0 – 100 in two criteria. We use Heuristic 85 (TS) as our subroutine in Heuristic 86 (PT). We give the seconds, the exact number of Pareto optima and the number of Pareto optima found. Heuristic 91 (BTRPT) not only found the correct number of Pareto optima, it found the correct Pareto optima in all cases. Heuristic 91 (BTRPT) can be distributed: once the boundary is computed and the regions are found, Heuristic 86 (PT) can be run as separate instances for each region (or subdivided further).

4.3.1.5. *Calibration - Local and Tabu Search For Convex Minimization.* For our experiments of Heuristic 82 (LS) and Heuristic 85 (TS) on the calibration set we minimized over two convex functions: $(\mathbf{x} - \hat{\mathbf{x}})^2$ and $(\mathbf{x} - \hat{\mathbf{x}})^4$. First we choose $\hat{\mathbf{x}} \in W\mathcal{P}_{\mathcal{M}}$ such that it is an interior point of $\overline{W\mathcal{P}_{\mathcal{M}}}$. Second we consider $\hat{\mathbf{x}}$ to be rational non-integer point which is an interior point of $\overline{W\mathcal{P}_{\mathcal{M}}}$. For the integral case, we can easily detect if we are at the minimum since our objective will evaluate to zero if so. For the rational case, we verify the global minimum by evaluating our objective on all projected spanning trees. For all tests we perform 1000 minimizations with random starts and record the number of successes. For easy reference, in the cases of Heuristic 82 (LS) we also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.




FIGURE 4.4. Green Pareto optima and blue test regions of *gn11e41d2w0w100* using Heuristic 91 (BTRPT)

- Table 4.12, Table 4.13, Table 4.14 show Heuristic 82 (LS) where the minimum is the integer point described above minimizing $(\mathbf{x} - \hat{\mathbf{x}})^2$.
- Table 4.15, Table 4.16, Table 4.17 show Heuristic 82 (LS) where the minimum is the integer point described above minimizing $(\mathbf{x} - \hat{\mathbf{x}})^4$.
- Table 4.24, Table 4.25, Table 4.26 show Heuristic 82 (LS) where the minimum is the rational point described above minimizing $(\mathbf{x} - \hat{\mathbf{x}})^2$.
- Table 4.27, Table 4.28, Table 4.29 show Heuristic 82 (LS) where the minimum is the rational point described above minimizing $(\mathbf{x} - \hat{\mathbf{x}})^4$.
- Table 4.18, Table 4.19, Table 4.20 show Heuristic 85 (TS) where the minimum is the integer point described above minimizing $(\mathbf{x} - \hat{\mathbf{x}})^2$.
- Table 4.21, Table 4.22, Table 4.23 show Heuristic 85 (TS) where the minimum is the integer point described above minimizing $(\mathbf{x} - \hat{\mathbf{x}})^4$.
- Table 4.30, Table 4.31, Table 4.32 show Heuristic 85 (TS) where the minimum is the rational point described above minimizing $(\mathbf{x} - \hat{\mathbf{x}})^2$.

- [Table 4.33](#), [Table 4.34](#), [Table 4.35](#) show [Heuristic 85](#) (TS) where the minimum is the rational point described above minimizing $(\mathbf{x} - \hat{\mathbf{x}})^4$.

We first observe that the success of [Heuristic 82](#) (LS) and [Heuristic 85](#) (TS) decrease as the proportion of projected spanning trees to all spanning trees increases. Second, for [Heuristic 85](#) (TS) we see a noticeable increase in the number of successes as the Tabu Limit ranges from 1 to 10 indicating that in many cases, a low Tabu Limit is sufficient to reach the global minimum.

4.3.2. Pushing the Limits.

4.3.2.1. *Sparse Graphs - Solids.* We now compare sparse graphs versus the dense graphs presented in the calibration set. We chose fifteen planar graphs of the 1-skeleton of three-dimensional polytopes. [Table 4.11](#) shows the result of [Heuristic 92](#) (DFBFS) on these examples. For some of these graphs, the number of spanning trees can be explicitly enumerated, hence we give the exact number of projected spanning trees. When the weights take value from 0 – 1000 we omit the last five graphs as DFBFS exceeded over a day of computation or the machines(*fuzzy*, *truth*) ran out of memory.

4.3.2.2. *Experimental Design.* In [\[BLMA⁺08\]](#) the authors first proposed using non-linear matroid optimization to solve the statistical experimental design problem, which is explained in detail in [\[FH80\]](#). The experimental design problem can be briefly described as attempting to learn an unknown system whose output $\mathbf{y}$ is an unknown function Φ with input $\mathbf{x} = (x_1, \dots, x_k) \in \mathbb{R}^k$. To learn the system, experiments are performed using input $\mathbf{p}_i = (p_{i,1}, \dots, p_{i,k})$ and the output $\mathbf{y}_i = \Phi(\mathbf{p}_i)$ is measured. Then, based on the experiments we wish to *fit a model* for the system, that is, determine an estimation $\hat{\Phi}$ of the function Φ such that it: lies in a prescribed class of functions, is consistent with the outcomes of the experiments, and minimizes the *aberration* (some suitable criteria) among models in the class.

A deterministic polynomial time algorithm was proved in [\[BLMA⁺08\]](#) which solves non-linear matroid optimization over arbitrary matroids (presented by evaluation oracle) when the number of weightings are fixed and the weightings are a general unary encoding. If d is the number of weightings, the weights only take value on p many integers, and the matroid

is on n elements, then their algorithm for arbitrary matroids uses the matroid intersection algorithm n^{p^d} a polynomial number of times.

However, for vector matroids they provide a faster algorithm (compared to their algorithm for arbitrary matroids) for unary encoded weights. If ω is the size of the maximum unary encoded element of the weights, then this improved algorithm requires solving a $(\text{rank}(\mathcal{M})\omega + 1)^d \times (\text{rank}(\mathcal{M})\omega + 1)^d$ Vandemonde equation. This non-linear vector matroid optimization algorithm has been implemented on IBM's Blue Gene/L computer [GLM08]. At the authors disposal was the 360TF, 128K processor Blue Gene/L supercomputer at Lawrence Livermore National Laboratories. We present now two instances P_{20} and P_{28} which were solved using the deterministic vector matroid algorithm on said supercomputer. Each is an instance of an experimental design problem encoded as a vector matroid non-linear optimization program. Both their algorithm and our heuristics (Heuristic 92 and Heuristic 86) do not optimize a particular objective function, but instead list the projected bases (note our heuristic is not guaranteed to find all projected bases).

Table 4.36 shows the size of the instances P_{20} and P_{28} , the true number of projected bases and the number of seconds required by Blue Gene/L to find the projected bases. We also show two runs of Heuristic 92 (DFBFS) using Fuzzy with the number of projected bases found and seconds required. What is noteworthy is that Heuristic 92 found all the projected bases in at least one of the two runs. Moreover, the other runs of Heuristic 92 found nearly all the projected bases. This demonstrates that our heuristics are useful when solving these difficult problems, especially in light of the fact that they were computed using a relatively cheap computer system. An additional point is that the deterministic vector matroid algorithm requires very high precision in order to solve the large Vandemonde systems derived from the matroid optimization problem. On the other hand, our heuristics only depend numerically on solving at most $\text{rank}(\mathcal{M}) \times \text{rank}(\mathcal{M})$ linear systems. In fact, our software has an option to use exact arithmetic, albeit with a slow down in solution times. We emphasize the drastic difference in running times and computational power between Blue Gene/L and Fuzzy.

In Table 4.37 we show the result of Heuristic 86 (PT) on P_{20} . We divided up the region of feasible projected bases into twelve disjoint regions and ran twelve instances of Heuristic 86 (PT) concurrently. We report the times and the number of projected bases

found for each region. We also give the total number of projected bases found which is the exact number of projected bases. For the search in [Heuristic 86](#) (PT) we used [Heuristic 85](#) (TS) 10 times on each point with a retry limit of 20. We performed this experiment to exhibit the effectiveness and ease by which [Heuristic 86](#) can be distributed.

4.3.3. Discussion. Non-linear matroid optimization problems are very difficult. They are NP-complete in general and exponential in some cases [\[BLMA⁺08\]](#). Our goal was to present and explore the practicality of new heuristics and algorithms for solving these problems. The effectiveness of our new techniques rely on two important properties: traversing using the local adjacency inherent in matroids is easy, and that although there may be exponentially many bases, the number of projected bases is manageable.

The fact that many of our heuristics are not guaranteed to find the optimal solution or list all the projected bases is countered by the fact that they are not overly complex and very fast. For instance, [Heuristic 92](#) (DFBFS) is not guaranteed to find all the projected bases but in many of the tests it found a large portion of them. In practice, as seen in [subsubsection 4.3.2.2](#), since [Heuristic 92](#) (DFBFS) finds a large portion of the projected bases, we would simply run the heuristic many times and union the results.

Most all of our heuristics and algorithms complement each other. For example, we can start by finding the vertices using [Algorithm 88](#) (PB) then enumerate as many projected bases as possible using [Heuristic 92](#) (DFBFS). Next we can run [Heuristic 86](#) (PT) on any regions we suspect there to be more projected bases.

Although we do not prove it theoretically, our heuristics and algorithms use minimal memory. For instance, the driving part of all of our heuristics and algorithms is enumerating neighboring bases, which is quadratic in the number of elements of the matroid. The biggest memory hog is [Heuristic 92](#) (DFBFS) which at worst stores all the projected bases found thus far.

This leads to the ease by which [Heuristic 86](#) (PT) can be distributed. In fact, the only overhead would be combining the solutions, which would be a simple union. Ideally if we had X machines with Y processors, we can first find a box containing the potential projected bases using $2n$ linear programs using [Heuristic 82](#) (LS). Next we partition the

region of potential projected bases into XY regions and run an instance of **Heuristic 86** (PT) for each.

TABLE 4.1. Calibration test set: Exact #Spanning trees vs. #projected spanning trees in 2 criteria.

Name	Nodes	Edges	Weight range	#Spanning Trees	#Projected Trees	Percent
gn9e18d2w0w20	9	18	0 – 20	2,981	1,742	58.44
gn9e27d2w0w20	9	27	0 – 20	372,320	6,346	01.70
gn10e22d2w0w20	10	22	0 – 20	53,357	3,957	07.42
gn10e28d2w0w20	10	28	0 – 20	800,948	7,131	00.89
gn10e33d2w0w20	10	33	0 – 20	3,584,016	10,833	00.30
gn11e13d2w0w20	11	13	0 – 20	41	41	100.00
gn11e20d2w0w20	11	20	0 – 20	6,939	2,173	31.32
gn11e27d2w0w20	11	27	0 – 20	284,730	7,515	02.64
gn11e41d2w0w20	11	41	0 – 20	90,922,271	16,457	00.02
gn12e16d2w0w20	12	16	0 – 20	162	154	95.06
gn12e24d2w0w20	12	24	0 – 20	208,380	5,647	02.71
gn12e33d2w0w20	12	33	0 – 20	4,741,624	9,364	00.20
gn13e19d2w0w20	13	19	0 – 20	3,401	1,608	47.28
gn13e29d2w0w20	13	29	0 – 20	1,543,340	8,474	00.55
gn13e39d2w0w20	13	39	0 – 20	131,807,934	18,468	00.01
gn9e18d2w0w100	9	18	0 – 100	2,981	2,858	95.87
gn9e27d2w0w100	9	27	0 – 100	372,320	89,092	23.93
gn10e22d2w0w100	10	22	0 – 100	53,357	37,204	69.73
gn10e28d2w0w100	10	28	0 – 100	800,948	101,334	12.65
gn10e33d2w0w100	10	33	0 – 100	3,584,016	166,427	04.64
gn11e13d2w0w100	11	13	0 – 100	41	41	100.00
gn11e20d2w0w100	11	20	0 – 100	6,939	6,580	94.83
gn11e27d2w0w100	11	27	0 – 100	284,730	81,803	28.73
gn11e41d2w0w100	11	41	0 – 100	90,922,271	309,961	00.34
gn12e16d2w0w100	12	16	0 – 100	162	162	100.00
gn12e24d2w0w100	12	24	0 – 100	208,380	92,813	44.54
gn12e33d2w0w100	12	33	0 – 100	4,741,624	192,122	04.05
gn13e19d2w0w100	13	19	0 – 100	3,401	3,255	95.71
gn13e29d2w0w100	13	29	0 – 100	1,543,340	164,617	10.67
gn13e39d2w0w100	13	39	0 – 100	131,807,934	315,881	00.24
gn9e18d2w0w1000	9	18	0 – 1000	2,981	2,981	100.00
gn9e27d2w0w1000	9	27	0 – 1000	372,320	364,382	97.87
gn10e22d2w0w1000	10	22	0 – 1000	53,357	52,990	99.31
gn10e28d2w0w1000	10	28	0 – 1000	800,948	756,013	94.39
gn10e33d2w0w1000	10	33	0 – 1000	3,584,016	2,726,287	76.07
gn11e13d2w0w1000	11	13	0 – 1000	41	41	100.00
gn11e20d2w0w1000	11	20	0 – 1000	6,939	6,921	99.74
gn11e27d2w0w1000	11	27	0 – 1000	284,730	279,308	98.10
gn11e41d2w0w1000	11	41	0 – 1000	90,922,271	13,884,793	15.27
gn12e16d2w0w1000	12	16	0 – 1000	162	162	100.00
gn12e24d2w0w1000	12	24	0 – 1000	208,380	205,690	98.71
gn12e33d2w0w1000	12	33	0 – 1000	4,741,624	3,680,313	77.62
gn13e19d2w0w1000	13	19	0 – 1000	3,401	3,401	100.00
gn13e29d2w0w1000	13	29	0 – 1000	1,543,340	1,396,180	90.46
gn13e39d2w0w1000	13	39	0 – 1000	131,807,934	15,037,589	11.41

4.3.4. Tables.

TABLE 4.2. Calibration test set: Exact #Spanning trees vs. #projected spanning trees in 3 criteria.

Name	Nodes	Edges	Weight range	#Spanning Trees	#Projected Trees	Percent
gn9e18d3w0w20	9	18	0 – 20	2,981	2,905	97.45
gn9e27d3w0w20	9	27	0 – 20	372,320	134,039	36.00
gn10e22d3w0w20	10	22	0 – 20	53,357	42,887	80.38
gn10e28d3w0w20	10	28	0 – 20	800,948	238,529	29.78
gn10e33d3w0w20	10	33	0 – 20	3,584,016	411,730	11.49
gn11e13d3w0w20	11	13	0 – 20	41	41	100.00
gn11e20d3w0w20	11	20	0 – 20	6,939	6,603	95.16
gn11e27d3w0w20	11	27	0 – 20	284,730	146,672	51.51
gn11e41d3w0w20	11	41	0 – 20	90,922,271	959,469	01.06
gn12e16d3w0w20	12	16	0 – 20	162	162	100.00
gn12e24d3w0w20	12	24	0 – 20	208,380	111,201	53.36
gn12e33d3w0w20	12	33	0 – 20	4,741,624	470,609	09.93
gn13e19d3w0w20	13	19	0 – 20	3,401	3,358	98.74
gn13e29d3w0w20	13	29	0 – 20	1,543,340	264,949	17.18
gn13e39d3w0w20	13	39	0 – 20	131,807,934	930,322	00.71
gn9e18d3w0w100	9	18	0 – 100	2,981	2,981	100.00
gn9e27d3w0w100	9	27	0 – 100	372,320	367,313	98.66
gn10e22d3w0w100	10	22	0 – 100	53,357	53,289	99.87
gn10e28d3w0w100	10	28	0 – 100	800,948	786,781	98.23
gn10e33d3w0w100	10	33	0 – 100	3,584,016	3,351,096	93.01
gn11e13d3w0w100	11	13	0 – 100	41	41	100.00
gn11e20d3w0w100	11	20	0 – 100	6,939	6,939	100.00
gn11e27d3w0w100	11	27	0 – 100	284,730	281,303	98.80
gn11e41d3w0w100	11	41	0 – 100	90,922,271	35,943,327	39.53
gn12e16d3w0w100	12	16	0 – 100	162	162	100.00
gn12e24d3w0w100	12	24	0 – 100	208,380	207,143	99.41
gn12e33d3w0w100	12	33	0 – 100	4,741,624	4,740,880	99.98
gn13e19d3w0w100	13	19	0 – 100	3,401	3,401	100.00
gn13e29d3w0w100	13	29	0 – 100	1,543,340	1,484,719	96.20
gn13e39d3w0w100	13	39	0 – 100	131,807,934	44,757,592	33.96
gn9e18d3w0w1000	9	18	0 – 1000	2,981	2,981	100.00
gn9e27d3w0w1000	9	27	0 – 1000	372,320	372,320	100.00
gn10e22d3w0w1000	10	22	0 – 1000	53,357	53,357	100.00
gn10e28d3w0w1000	10	28	0 – 1000	800,948	800,946	99.99
gn10e33d3w0w1000	10	33	0 – 1000	3,584,016	3,583,757	99.99
gn11e13d3w0w1000	11	13	0 – 1000	41	41	100.00
gn11e20d3w0w1000	11	20	0 – 1000	6,939	6,939	100.00
gn11e27d3w0w1000	11	27	0 – 1000	284,730	284,730	100.00
gn11e41d3w0w1000	11	41	0 – 1000	90,922,271	90,699,181	99.75
gn12e16d3w0w1000	12	16	0 – 1000	162	162	100.00
gn12e24d3w0w1000	12	24	0 – 1000	208,380	208,356	99.99
gn12e33d3w0w1000	12	33	0 – 1000	4,741,624	4,740,880	99.98
gn13e19d3w0w1000	13	19	0 – 1000	3,401	3,401	100.00
gn13e29d3w0w1000	13	29	0 – 1000	1,543,340	1,543,304	99.99
gn13e39d3w0w1000	13	39	0 – 1000	131,807,934	131,464,478	99.74

TABLE 4.3. Calibration test set: Exact #Spanning trees vs. #projected spanning trees in 5 criteria.

Name	Nodes	Edges	Weight range	#Spanning Trees	#Projected Trees	Percent
gn9e18d5w0w1	9	18	0 – 1	2,981	1,099	36.87
gn9e27d5w0w1	9	27	0 – 1	372,320	8,205	2.20
gn10e22d5w0w1	10	22	0 – 1	53,357	4,746	8.89
gn10e28d5w0w1	10	28	0 – 1	800,948	10,898	1.36
gn10e33d5w0w1	10	33	0 – 1	3,584,016	15,482	0.43
gn11e13d5w0w1	11	13	0 – 1	41	27	65.85
gn11e20d5w0w1	11	20	0 – 1	6,939	1,866	26.89
gn11e27d5w0w1	11	27	0 – 1	284,730	5,828	2.05
gn11e41d5w0w1	11	41	0 – 1	90,922,271	36,847	0.04
gn12e16d5w0w1	12	16	0 – 1	162	116	71.61
gn12e24d5w0w1	12	24	0 – 1	208,380	6,738	3.23
gn12e33d5w0w1	12	33	0 – 1	4,741,624	20,857	0.44
gn13e19d5w0w1	13	19	0 – 1	3,401	1,149	33.78
gn13e29d5w0w1	13	29	0 – 1	1,543,340	111,116	7.20
gn13e39d5w0w1	13	39	0 – 1	131,807,934	34,392	0.03
gn9e18d5w0w2	9	18	0 – 2	2,981	2,276	76.35
gn9e27d5w0w2	9	27	0 – 2	372,320	43,029	11.56
gn10e22d5w0w2	10	22	0 – 2	53,357	14,623	27.41
gn10e28d5w0w2	10	28	0 – 2	800,948	66,190	8.26
gn10e33d5w0w2	10	33	0 – 2	3,584,016	105,309	2.94
gn11e13d5w0w2	11	13	0 – 2	41	41	100.00
gn11e20d5w0w2	11	20	0 – 2	6,939	3,761	54.20
gn11e27d5w0w2	11	27	0 – 2	284,730	39,292	13.80
gn11e41d5w0w2	11	41	0 – 2	90,922,271	290,555	0.32
gn12e16d5w0w2	12	16	0 – 2	162	160	98.77
gn12e24d5w0w2	12	24	0 – 2	208,380	34,057	16.34
gn12e33d5w0w2	12	33	0 – 2	4,741,624	120,211	2.54
gn13e19d5w0w2	13	19	0 – 2	3,401	2,613	76.83
gn13e29d5w0w2	13	29	0 – 2	1,543,340	98,285	6.37
gn13e39d5w0w2	13	39	0 – 2	131,807,934	348,703	0.26
gn9e18d5w0w5	9	18	0 – 5	2,981	2,960	99.30
gn9e27d5w0w5	9	27	0 – 5	372,320	271,048	72.80
gn10e22d5w0w5	10	22	0 – 5	53,357	49,463	92.70
gn10e28d5w0w5	10	28	0 – 5	800,948	493,565	61.62
gn10e33d5w0w5	10	33	0 – 5	3,584,016	1,294,875	36.13
gn11e13d5w0w5	11	13	0 – 5	41	41	100.00
gn11e20d5w0w5	11	20	0 – 5	6,939	5,975	86.11
gn11e27d5w0w5	11	27	0 – 5	284,730	222,974	78.31
gn11e41d5w0w5	11	41	0 – 5	90,922,271	4,711,354	5.18
gn12e16d5w0w5	12	16	0 – 5	162	162	100.00
gn12e24d5w0w5	12	24	0 – 5	208,380	177,845	85.35
gn12e33d5w0w5	12	33	0 – 5	4,741,624	1,489,751	31.42
gn13e19d5w0w5	13	19	0 – 5	3,401	3,381	99.41
gn13e29d5w0w5	13	29	0 – 5	1,543,340	787,196	51.01
gn13e39d5w0w5	13	39	0 – 5	131,807,934	7,737,684	5.87

TABLE 4.4. Calibration test set: **Heuristic 92** (DFBFS) vs. exact # of projected spanning trees in 2 criteria. We tried 100 searches, depth 4, boundary retry limit 100 and interior retry limit of 10000.

Name	Nodes	Edges	Weight	Computer	Seconds	#Spanning Trees	#Projected Trees	#Trees Found by DFBFS	Percent
gn9e18d2w0w100	9	18	0 - 20	Fuzzy	0	2, 981	1, 742	1, 717	98.56
gn9e27d2w0w100	9	27	0 - 20	Fuzzy	1	372, 320	6, 346	6, 258	98.61
gn10e22d2w0w100	10	22	0 - 20	Fuzzy	0	53, 357	3, 957	3, 852	97.34
gn10e28d2w0w100	10	28	0 - 20	Fuzzy	2	800, 948	7, 131	7, 074	99.20
gn10e33d2w0w100	10	33	0 - 20	Fuzzy	2	3, 584, 016	10, 833	10, 734	99.08
gn11e13d2w0w100	11	13	0 - 20	Fuzzy	0	41	41	37	90.24
gn11e20d2w0w100	11	20	0 - 20	Fuzzy	1	6, 939	2, 173	2, 096	96.45
gn11e27d2w0w100	11	27	0 - 20	Fuzzy	1	284, 730	7, 515	7, 417	98.69
gn11e41d2w0w100	11	41	0 - 20	Fuzzy	4	90, 922, 271	16, 457	16, 262	98.81
gn12e16d2w0w100	12	16	0 - 20	Fuzzy	1	162	154	153	99.35
gn12e24d2w0w100	12	24	0 - 20	Fuzzy	1	208, 380	5, 647	5, 535	98.01
gn12e33d2w0w100	12	33	0 - 20	Fuzzy	3	4, 741, 624	9, 364	9, 325	99.58
gn13e19d2w0w100	13	19	0 - 20	Fuzzy	0	3, 401	1, 608	1, 587	98.69
gn13e29d2w0w100	13	29	0 - 20	Fuzzy	2	1, 543, 340	8, 474	8, 436	99.55
gn13e39d2w0w100	13	39	0 - 20	Fuzzy	5	131, 807, 934	18, 468	18, 330	99.25
gn9e18d2w0w100	9	18	0 - 100	Fuzzy	0	2, 981	2, 858	2, 799	97.93
gn9e27d2w0w100	9	27	0 - 100	Fuzzy	6	372, 320	89, 092	81, 750	91.75
gn10e22d2w0w100	10	22	0 - 100	Fuzzy	3	53, 357	37, 204	35, 581	95.63
gn10e28d2w0w100	10	28	0 - 100	Fuzzy	8	800, 948	101, 334	95, 313	94.05
gn10e33d2w0w100	10	33	0 - 100	Fuzzy	14	3, 584, 016	166, 427	157, 685	94.74
gn11e13d2w0w100	11	13	0 - 100	Fuzzy	0	41	41	41	100.00
gn11e20d2w0w100	11	20	0 - 100	Fuzzy	1	6, 939	6, 580	6, 513	98.98
gn11e27d2w0w100	11	27	0 - 100	Fuzzy	7	284, 730	81, 803	76, 372	93.36
gn11e41d2w0w100	11	41	0 - 100	Fuzzy	32	90, 922, 271	309, 961	298, 150	96.18
gn12e16d2w0w100	12	16	0 - 100	Fuzzy	0	162	162	162	100.00
gn12e24d2w0w100	12	24	0 - 100	Fuzzy	9	208, 380	92, 813	88, 361	95.20
gn12e33d2w0w100	12	33	0 - 100	Fuzzy	17	4, 741, 624	192, 122	175, 048	91.11
gn13e19d2w0w100	13	19	0 - 100	Fuzzy	0	3, 401	3, 255	3, 219	98.89
gn13e29d2w0w100	13	29	0 - 100	Fuzzy	18	1, 543, 340	164, 617	156, 608	95.13
gn13e39d2w0w100	13	39	0 - 100	Fuzzy	37	131, 807, 934	315, 882	301, 567	95.46
gn9e18d2w0w1000	9	18	0 - 1000	Fuzzy	1	2, 981	2, 981	2, 961	99.32
gn9e27d2w0w1000	9	27	0 - 1000	Fuzzy	16	372, 320	364, 382	307, 755	84.45
gn10e22d2w0w1000	10	22	0 - 1000	Fuzzy	5	53, 357	52, 990	51, 782	97.72
gn10e28d2w0w1000	10	28	0 - 1000	Fuzzy	46	800, 948	756, 013	659, 761	87.26
gn10e33d2w0w1000	10	33	0 - 1000	Fuzzy	90	3, 584, 016	2, 726, 287	1, 628, 981	59.75
gn11e13d2w0w1000	11	13	0 - 1000	Fuzzy	0	41	41	41	100.00
gn11e20d2w0w1000	11	20	0 - 1000	Fuzzy	1	6, 939	6, 921	6, 831	98.69
gn11e27d2w0w1000	11	27	0 - 1000	Fuzzy	17	284, 730	279, 308	254, 290	91.04
gn11e41d2w0w1000	11	41	0 - 1000	Fuzzy	455	90, 922, 271	13, 884, 793	7092, 823	51.08
gn12e16d2w0w1000	12	16	0 - 1000	Fuzzy	0	162	162	162	100.00
gn12e24d2w0w1000	12	24	0 - 1000	Fuzzy	17	208, 380	205, 690	200, 129	97.29
gn12e33d2w0w1000	12	33	0 - 1000	Fuzzy	187	4, 741, 624	3, 680, 313	2, 651, 306	72.04
gn13e19d2w0w1000	13	19	0 - 1000	Fuzzy	1	3, 401	3, 401	3, 384	99.50
gn13e29d2w0w1000	13	29	0 - 1000	Fuzzy	104	1, 543, 340	1, 396, 180	1, 247, 040	89.31
gn13e39d2w0w1000	13	39	0 - 1000	Fuzzy	825	131, 807, 934	15, 037, 589	10, 260, 810	68.23

TABLE 4.5. Calibration test set: **Heuristic 92** (DFBFS) vs. exact # of projected spanning trees in 3 criteria. We tried 100 searches, depth 4, boundary retry limit 100 and interior retry limit of 10000.

Name	Nodes	Edges	Weight range	Computer	Seconds	#Spanning Trees	#Projected Trees	#Trees Found by DFBFS	Percent
gn9e18d3w0w100	9	18	0 - 20	Fuzzy	0	2, 981	2, 905	2, 854	98.24
gn9e27d3w0w100	9	27	0 - 20	Fuzzy	10	372, 320	134, 039	124, 212	92.66
gn10e22d3w0w100	10	22	0 - 20	Fuzzy	4	53, 357	42, 887	41, 649	97.11
gn10e28d3w0w100	10	28	0 - 20	Fuzzy	19	800, 948	238, 529	217, 444	91.16
gn10e33d3w0w100	10	33	0 - 20	Fuzzy	36	3, 584, 016	411, 730	374, 110	90.86
gn11e13d3w0w100	11	13	0 - 20	Fuzzy	1	41	41	41	100.00
gn11e20d3w0w100	11	20	0 - 20	Fuzzy	0	6, 939	6, 603	6, 532	98.92
gn11e27d3w0w100	11	27	0 - 20	Fuzzy	12	284, 730	146, 672	136, 807	93.27
gn11e41d3w0w100	11	41	0 - 20	Fuzzy	97	90, 922, 271	959, 469	876, 488	91.35
gn12e16d3w0w100	12	16	0 - 20	Fuzzy	1	162	162	162	100.00
gn12e24d3w0w100	12	24	0 - 20	Fuzzy	11	208, 380	111, 201	107, 064	96.27
gn12e33d3w0w100	12	33	0 - 20	Fuzzy	48	4, 741, 624	470, 609	434, 245	92.27
gn13e19d3w0w100	13	19	0 - 20	Fuzzy	1	3, 401	3, 358	3, 319	98.83
gn13e29d3w0w100	13	29	0 - 20	Fuzzy	29	1, 543, 340	264, 949	251, 408	94.88
gn13e39d3w0w100	13	39	0 - 20	Fuzzy	106	131, 807, 934	930, 322	872, 397	93.77
gn9e18d3w0w100	9	18	0 - 100	Fuzzy	0	2, 981	2, 981	2, 912	97.68
gn9e27d3w0w100	9	27	0 - 100	Fuzzy	20	372, 320	367, 313	330, 331	89.93
gn10e22d3w0w100	10	22	0 - 100	Fuzzy	4	53, 357	53, 289	52, 296	98.13
gn10e28d3w0w100	10	28	0 - 100	Fuzzy	50	800, 948	786, 781	699, 399	88.89
gn10e33d3w0w100	10	33	0 - 100	Fuzzy	133	3, 584, 016	3, 351, 096	2, 095, 143	66.66
gn11e13d3w0w100	11	13	0 - 100	Fuzzy	0	41	41	41	100.00
gn11e20d3w0w100	11	20	0 - 100	Fuzzy	1	6, 939	6, 939	6, 895	99.36
gn11e27d3w0w100	11	27	0 - 100	Fuzzy	21	284, 730	281, 303	267, 912	95.23
gn11e41d3w0w100	11	41	0 - 100	Fuzzy	801	90, 922, 271	35, 943, 327	13, 527, 453	37.14
gn12e16d3w0w100	12	16	0 - 100	Fuzzy	0	162	162	162	100.00
gn12e24d3w0w100	12	24	0 - 100	Fuzzy	18	208, 380	207, 143	202, 199	97.61
gn12e33d3w0w100	12	33	0 - 100	Fuzzy	260	4, 741, 624	4, 740, 880	3, 326, 573	75.00
gn13e19d3w0w100	13	19	0 - 100	Fuzzy	1	3, 401	3, 401	3, 385	99.52
gn13e29d3w0w100	13	29	0 - 100	Fuzzy	108	1, 543, 340	1, 484, 719	1, 318, 092	100.00
gn13e39d3w0w100	13	39	0 - 100	Fuzzy	1777	131, 807, 934	44, 757, 592	23, 502, 664	52.27
gn9e18d3w0w1000	9	18	0 - 1000	Fuzzy	1	2, 981	2, 981	2, 861	95.97
gn9e27d3w0w1000	9	27	0 - 1000	Fuzzy	18	372, 320	372, 320	331, 050	88.91
gn10e22d3w0w1000	10	22	0 - 1000	Fuzzy	4	53, 357	53, 357	51, 592	96.69
gn10e28d3w0w1000	10	28	0 - 1000	Fuzzy	52	800, 948	800, 946	721, 587	90.09
gn10e33d3w0w1000	10	33	0 - 1000	Fuzzy	135	3, 584, 016	3, 583, 757	2, 313, 881	64.56
gn11e13d3w0w1000	11	13	0 - 1000	Fuzzy	0	41	41	41	100.00
gn11e20d3w0w1000	11	20	0 - 1000	Fuzzy	1	6, 939	6, 939	6, 780	97.70
gn11e27d3w0w1000	11	27	0 - 1000	Fuzzy	20	284, 730	284, 730	263, 159	92.42
gn11e41d3w0w1000	11	41	0 - 1000	Fuzzy	1021	90, 922, 271	90, 699, 181	16, 919, 561	18.65
gn12e16d3w0w1000	12	16	0 - 1000	Fuzzy	0	162	162	162	100.00
gn12e24d3w0w1000	12	24	0 - 1000	Fuzzy	17	208, 380	208, 356	203, 347	97.59
gn12e33d3w0w1000	12	33	0 - 1000	Fuzzy	238	4, 741, 624	4, 740, 880	3, 376, 122	71.21
gn13e19d3w0w1000	13	19	0 - 1000	Fuzzy	1	3, 401	3, 401	3, 372	99.14
gn13e29d3w0w1000	13	29	0 - 1000	Fuzzy	118	1, 543, 340	1, 543, 304	1, 389, 181	90.01
gn13e39d3w0w1000	13	39	0 - 1000	Fuzzy	2315	131, 807, 934	131, 464, 478	34, 603, 989	26.32

TABLE 4.6. Calibration test set: **Heuristic 92** (DFBFS) vs. exact # of projected spanning trees in 5 criteria. We tried 100 searches, depth 4, boundary retry limit 100 and interior retry limit of 10000.

Name	Nodes	Edges	Weight	Computer	Seconds	#Spanning Trees	#Projected Trees	#Trees Found by DFBFS	Percent
gn9e18d5w0w1	9	18	0 - 1	Fuzzy	0	2, 981	1, 099	1, 092	99.36
gn9e27d5w0w1	9	27	0 - 1	Fuzzy	3	372, 320	8, 205	8, 079	98.46
gn10e22d5w0w1	10	22	0 - 1	Fuzzy	1	53, 357	4, 746	4, 704	99.12
gn10e28d5w0w1	10	28	0 - 1	Fuzzy	3	800, 948	10, 898	10, 655	97.77
gn10e33d5w0w1	10	33	0 - 1	Fuzzy	4	3, 584, 016	15, 482	15, 235	98.40
gn11e13d5w0w1	11	13	0 - 1	Fuzzy	0	41	27	27	100.00
gn11e20d5w0w1	11	20	0 - 1	Fuzzy	1	6, 939	1, 866	1, 845	98.87
gn11e27d5w0w1	11	27	0 - 1	Fuzzy	2	284, 730	5, 828	5, 702	97.84
gn11e41d5w0w1	11	41	0 - 1	Fuzzy	14	90, 922, 271	36, 847	36, 457	98.94
gn12e16d5w0w1	12	16	0 - 1	Fuzzy	0	162	116	116	100.00
gn12e24d5w0w1	12	24	0 - 1	Fuzzy	2	208, 380	6, 738	6, 665	98.92
gn12e33d5w0w1	12	33	0 - 1	Fuzzy	6	4, 741, 624	20, 857	20, 652	99.02
gn13e19d5w0w1	13	19	0 - 1	Fuzzy	1	3, 401	1, 149	1, 142	99.39
gn13e29d5w0w1	13	29	0 - 1	Fuzzy	4	1, 543, 340	111, 116	10, 953	99.86
gn13e39d5w0w1	13	39	0 - 1	Fuzzy	16	131, 807, 934	34, 392	34, 076	99.08
gn9e18d5w0w2	9	18	0 - 2	Fuzzy	0	2, 981	2, 276	2, 253	98.99
gn9e27d5w0w2	9	27	0 - 2	Fuzzy	5	372, 320	43, 029	42, 188	98.05
gn10e22d5w0w2	10	22	0 - 2	Fuzzy	2	53, 357	14, 623	14, 205	97.14
gn10e28d5w0w2	10	28	0 - 2	Fuzzy	8	800, 948	66, 190	63, 992	96.68
gn10e33d5w0w2	10	33	0 - 2	Fuzzy	16	3, 584, 016	105, 309	103, 221	98.02
gn11e13d5w0w2	11	13	0 - 2	Fuzzy	0	41	41	41	100.00
gn11e20d5w0w2	11	20	0 - 2	Fuzzy	1	6, 939	3, 761	3, 708	98.59
gn11e27d5w0w2	11	27	0 - 2	Fuzzy	6	284, 730	39, 292	38, 228	97.29
gn11e41d5w0w2	11	41	0 - 2	Fuzzy	49	90, 922, 271	290, 555	281, 519	96.89
gn12e16d5w0w2	12	16	0 - 2	Fuzzy	0	162	160	160	100.00
gn12e24d5w0w2	12	24	0 - 2	Fuzzy	5	208, 380	34, 057	32, 993	96.88
gn12e33d5w0w2	12	33	0 - 2	Fuzzy	19	4, 741, 624	120, 211	116, 691	97.08
gn13e19d5w0w2	13	19	0 - 2	Fuzzy	1	3, 401	2, 613	2, 586	98.97
gn13e29d5w0w2	13	29	0 - 2	Fuzzy	17	1, 543, 340	98, 285	95, 528	97.19
gn13e39d5w0w2	13	39	0 - 2	Fuzzy	64	131, 807, 934	348, 703	339, 179	97.27
gn9e18d5w0w5	9	18	0 - 5	Fuzzy	0	2, 981	2, 960	2, 936	99.19
gn9e27d5w0w5	9	27	0 - 5	Fuzzy	22	372, 320	271, 048	253, 388	93.49
gn10e22d5w0w5	10	22	0 - 5	Fuzzy	5	53, 357	49, 463	48, 569	98.19
gn10e28d5w0w5	10	28	0 - 5	Fuzzy	44	800, 948	493, 565	460, 744	93.35
gn10e33d5w0w5	11	33	0 - 5	Fuzzy	114	3, 584, 016	1, 294, 875	1, 145, 396	88.46
gn11e13d5w0w5	11	13	0 - 5	Fuzzy	0	41	41	41	100.00
gn11e20d5w0w5	11	20	0 - 5	Fuzzy	1	6, 939	5, 975	5, 915	99.00
gn11e27d5w0w5	11	27	0 - 5	Fuzzy	21	284, 730	222, 974	213, 780	95.88
gn11e41d5w0w5	11	41	0 - 5	Fuzzy	561	90, 922, 271	4, 711, 354	4, 247, 218	90.15
gn12e16d5w0w5	12	16	0 - 5	Fuzzy	1	162	162	161	99.38
gn12e24d5w0w5	12	24	0 - 5	Fuzzy	19	208, 380	177, 845	173, 181	97.38
gn12e33d5w0w5	12	33	0 - 5	Fuzzy	156	4, 741, 624	1, 489, 751	1, 323, 574	88.85
gn13e19d5w0w5	13	19	0 - 5	Fuzzy	1	3, 401	3, 381	3, 360	99.38
gn13e29d5w0w5	13	29	0 - 5	Fuzzy	88	1, 543, 340	787, 196	733, 793	93.22
gn13e39d5w0w5	13	39	0 - 5	Fuzzy	907	131, 807, 934	7, 737, 684	6, 800, 052	87.88

TABLE 4.7. Calibration test set: **Heuristic 86** (Pivot Test) using **Heuristic 82** (Local Search), 10 searches per point.

Name	Nodes	Edges	Dimension	Weights	Computer	Seconds	#Projected Trees	#Trees Found by DFBFS	Percent
gn9e18d2w0w20	9	18	2	0 – 20	mocr01	92	1,742	1,121	64.35
gn9e27d2w0w20	9	27	2	0 – 20	mocr01	169	6,346	4,939	77.83
gn10e22d2w0w20	10	22	2	0 – 20	mocr01	101	3,957	3,318	83.85
gn10e28d2w0w20	10	28	2	0 – 20	mocr01	207	7,131	6,163	86.43
gn10e33d2w0w20	10	33	2	0 – 20	mocr01	356	10,833	8,842	81.62
gn11e13d2w0w20	11	13	2	0 – 20	mocr01	5	41	37	90.24
gn11e20d2w0w20	11	20	2	0 – 20	mocr01	89	2,173	1,328	61.11
gn11e27d2w0w20	11	27	2	0 – 20	mocr01	273	7,515	5,803	77.22
gn11e41d2w0w20	11	41	2	0 – 20	mocr01	806	16,457	14,747	89.61
gn12e16d2w0w20	12	16	2	0 – 20	mocr01	13	154	144	93.51
gn12e24d2w0w20	12	24	2	0 – 20	mocr01	233	5,647	3,701	65.54
gn12e33d2w0w20	12	33	2	0 – 20	mocr01	357	9,364	7,599	81.15
gn13e19d2w0w20	13	19	2	0 – 20	mocr01	68	1,608	1,051	65.36
gn13e29d2w0w20	13	29	2	0 – 20	mocr01	359	8,474	6,407	75.61
gn13e39d2w0w20	13	39	2	0 – 20	mocr01	1,160	18,468	15,082	81.67
gn9e18d2w0w100	9	18	2	0 – 100	mocr02	944	2,858	1,346	47.10
gn9e27d2w0w100	9	27	2	0 – 100	mocr02	5,923	89,092	8,760	09.83
gn10e22d2w0w100	10	22	2	0 – 100	mocr02	4,315	37,204	5,249	14.11
gn10e28d2w0w100	10	28	2	0 – 100	mocr02	6,224	101,334	8,236	08.13
gn10e33d2w0w100	10	33	2	0 – 100	mocr02	11,925	166,427	19,247	11.56
gn11e13d2w0w100	11	13	2	0 – 100	mocr02	215	41	41	100.00
gn11e20d2w0w100	11	20	2	0 – 100	mocr02	2,737	6,580	2,107	32.02
gn11e27d2w0w100	11	27	2	0 – 100	mocr02	6,327	81,803	6,790	08.30
gn11e41d2w0w100	11	41	2	0 – 100	mocr02	26,144	309,961	29,373	09.48
gn12e16d2w0w100	12	16	2	0 – 100	mocr02	272	162	155	95.68
gn12e24d2w0w100	12	24	2	0 – 100	mocr02	11,812	92,813	11,596	12.49
gn12e33d2w0w100	12	33	2	0 – 100	mocr02	15,676	192,122	16,705	08.70
gn13e19d2w0w100	13	19	2	0 – 100	mocr02	1,859	3,255	2,010	61.75
gn13e29d2w0w100	13	29	2	0 – 100	mocr02	15,365	164,617	12,986	07.89
gn13e39d2w0w100	13	39	2	0 – 100	mocr02	32,958	315,881	38,530	12.20
gn9e18d3w0w20	9	18	2	0 – 100	mocr01	3,395	2,905	1,659	57.11
gn9e27d3w0w20	9	27	2	0 – 100	mocr01	27,537	134,039	16,842	12.57
gn10e22d3w0w20	10	22	2	0 – 100	mocr01	18,141	42,887	8,564	19.97
gn10e28d3w0w20	10	28	2	0 – 100	mocr01	50,429	238,529	39,819	16.69
gn10e33d3w0w20	10	33	2	0 – 100	mocr01	76,051	411,730	47,606	11.56
gn11e13d3w0w20	11	13	2	0 – 100	mocr01	271	41	40	97.56
gn11e20d3w0w20	11	20	2	0 – 100	mocr01	6,726	6,603	2,851	43.18
gn11e27d3w0w20	11	27	2	0 – 100	mocr01	39,970	146,672	21,790	14.86
gn11e41d3w0w20	11	41	2	0 – 100	mocr01	216,987	959,469	120,142	12.52
gn12e16d3w0w20	12	16	2	0 – 100	mocr01	1,329	162	155	95.68
gn12e24d3w0w20	12	24	2	0 – 100	mocr01	32,498	111,201	16,698	15.02
gn12e33d3w0w20	12	33	2	0 – 100	mocr01	107,663	470,609	53,432	11.35
gn13e19d3w0w20	13	19	2	0 – 100	mocr01	9,980	3,358	2,255	67.15
gn13e29d3w0w20	13	29	2	0 – 100	mocr01	62,506	264,949	34,109	12.87
gn13e39d3w0w20	13	39	2	0 – 100	mocr01	232,975	930,322	126,044	13.55

TABLE 4.8. Calibration test set: **Heuristic 86** (Pivot Test) using **Heuristic 85** (Tabu Search), 10 searches per point, 20 search limit.

Name	Nodes	Edges	Dimension	Weights	Computer	Seconds	#Projected Trees	#Trees Found by DFBFS	Percent
gn9e18d2w0w20	9	18	2	0 – 20	mocr04	333	1,742	1,643	94.32
gn9e27d2w0w20	9	27	2	0 – 20	mocr04	518	6,346	6,299	99.26
gn10e22d2w0w20	10	22	2	0 – 20	mocr04	394	3,957	3,878	98.00
gn10e28d2w0w20	10	28	2	0 – 20	mocr04	712	7,131	7,085	99.35
gn10e33d2w0w20	10	33	2	0 – 20	mocr04	1,146	10,833	10,739	99.13
gn11e13d2w0w20	11	13	2	0 – 20	mocr04	41	41	37	90.24
gn11e20d2w0w20	11	20	2	0 – 20	mocr04	440	2,173	1,979	91.07
gn11e27d2w0w20	11	27	2	0 – 20	mocr04	971	7,515	7,441	99.02
gn11e41d2w0w20	11	41	2	0 – 20	mocr04	2,484	16,457	16,379	99.53
gn12e16d2w0w20	12	16	2	0 – 20	mocr04	85	154	153	99.35
gn12e24d2w0w20	12	24	2	0 – 20	mocr04	898	5,647	5,449	96.49
gn12e33d2w0w20	12	33	2	0 – 20	mocr04	1,082	9,364	9,270	99.00
gn13e19d2w0w20	13	19	2	0 – 20	mocr04	340	1,608	1,521	94.59
gn13e29d2w0w20	13	29	2	0 – 20	mocr04	1,102	8,474	8,408	99.22
gn13e39d2w0w20	13	39	2	0 – 20	mocr04	3,233	18,468	18,316	99.18
gn9e18d2w0w100	9	18	2	0 – 100	mocr04	6,231	2,858	2,556	94.32
gn9e27d2w0w100	9	27	2	0 – 100	mocr04	30,162	89,092	31,929	99.26
gn10e22d2w0w100	10	22	2	0 – 100	mocr04	25,007	37,204	20,976	98.00
gn10e28d2w0w100	10	28	2	0 – 100	mocr04	33,342	101,334	50,722	99.35
gn10e33d2w0w100	10	33	2	0 – 100	mocr04	56,517	166,427	82,275	99.13
gn11e13d2w0w100	11	13	2	0 – 100	mocr04	1,561	41	41	90.24
gn11e20d2w0w100	11	20	2	0 – 100	mocr04	17,385	6,580	4,972	91.07
gn11e27d2w0w100	11	27	2	0 – 100	mocr04	36,726	81,803	33,446	99.02
gn11e41d2w0w100	11	41	2	0 – 100	mocr04	120,393	309,961	188,619	99.53
gn12e16d2w0w100	12	16	2	0 – 100	mocr04	1,750	162	161	99.35
gn12e24d2w0w100	12	24	2	0 – 100	mocr04	62,618	92,813	47,870	96.49
gn12e33d2w0w100	12	33	2	0 – 100	mocr04	78,539	192,122	101,544	99.00
gn13e19d2w0w100	13	19	2	0 – 100	mocr04	9,703	3,255	2,467	94.59
gn13e29d2w0w100	13	29	2	0 – 100	mocr04	68,776	164,617	48,053	99.22
gn13e39d2w0w100	13	39	2	0 – 100	mocr04	138,832	315,881	197,263	99.18

TABLE 4.9. Calibration test set: **Algorithm 88** (Boundary Calculation) with 2 criteria. All times under 1 second.

Name	Nodes	Edges	Weight range	Computer	#Spanning Trees	#Projected Trees	Boundary points
gn9e18d2w0w20	9	18	0 – 20	mocr02	2,981	1,742	20
gn9e27d2w0w20	9	27	0 – 20	mocr02	372,320	6,346	29
gn10e22d2w0w20	10	22	0 – 20	mocr02	53,357	3,957	35
gn10e28d2w0w20	10	28	0 – 20	mocr02	800,948	7,131	33
gn10e33d2w0w20	10	33	0 – 20	mocr02	3,584,016	10,833	10
gn11e13d2w0w20	11	13	0 – 20	mocr02	41	41	10
gn11e20d2w0w20	11	20	0 – 20	mocr02	6,939	2,173	20
gn11e27d2w0w20	11	27	0 – 20	mocr02	284,730	7,515	27
gn11e41d2w0w20	11	41	0 – 20	mocr02	90,922,271	16,457	38
gn12e16d2w0w20	12	16	0 – 20	mocr02	162	154	13
gn12e24d2w0w20	12	24	0 – 20	mocr02	208,380	5,647	25
gn12e33d2w0w20	12	33	0 – 20	mocr02	4,741,624	9,364	35
gn13e19d2w0w20	13	19	0 – 20	mocr02	3,401	1,608	21
gn13e29d2w0w20	13	29	0 – 20	mocr02	1,543,340	8,474	36
gn13e39d2w0w20	13	39	0 – 20	mocr02	131,807,934	18,468	52
gn9e18d2w0w100	9	18	0 – 100	mocr02	2,981	2,858	19
gn9e27d2w0w100	9	27	0 – 100	mocr02	372,320	89,092	31
gn10e22d2w0w100	10	22	0 – 100	mocr02	53,357	37,204	26
gn10e28d2w0w100	10	28	0 – 100	mocr02	800,948	101,334	31
gn10e33d2w0w100	10	33	0 – 100	mocr02	3,584,016	166,427	37
gn11e13d2w0w100	11	13	0 – 100	mocr02	41	41	9
gn11e20d2w0w100	11	20	0 – 100	mocr02	6,939	6,580	22
gn11e27d2w0w100	11	27	0 – 100	mocr02	284,730	81,803	30
gn11e41d2w0w100	11	41	0 – 100	mocr02	90,922,271	309,961	41
gn12e16d2w0w100	12	16	0 – 100	mocr02	162	162	11
gn12e24d2w0w100	12	24	0 – 100	mocr02	208,380	92,813	29
gn12e33d2w0w100	12	33	0 – 100	mocr02	4,741,624	192,122	32
gn13e19d2w0w100	13	19	0 – 100	mocr02	3,401	3,255	22
gn13e29d2w0w100	13	29	0 – 100	mocr02	1,543,340	164,617	37
gn13e39d2w0w100	13	39	0 – 100	mocr02	131,807,934	315,881	42
gn9e18d2w0w1000	9	18	0 – 1000	mocr02	2,981	2,981	26
gn9e27d2w0w1000	9	27	0 – 1000	mocr02	372,320	364,382	31
gn10e22d2w0w1000	10	22	0 – 1000	mocr02	53,357	52,990	37
gn10e28d2w0w1000	10	28	0 – 1000	mocr02	800,948	756,013	9
gn10e33d2w0w1000	10	33	0 – 1000	mocr02	3,584,016	2,726,287	21
gn11e13d2w0w1000	11	13	0 – 1000	mocr02	41	41	28
gn11e20d2w0w1000	11	20	0 – 1000	mocr02	6,939	6,921	42
gn11e27d2w0w1000	11	27	0 – 1000	mocr02	284,730	279,308	13
gn11e41d2w0w1000	11	41	0 – 1000	mocr02	90,922,271	13,884,793	31
gn12e16d2w0w1000	12	16	0 – 1000	mocr02	162	162	32
gn12e24d2w0w1000	12	24	0 – 1000	mocr02	208,380	205,690	18
gn12e33d2w0w1000	12	33	0 – 1000	mocr02	4,741,624	3,680,313	36
gn13e19d2w0w1000	13	19	0 – 1000	mocr02	3,401	3,401	43
gn13e29d2w0w1000	13	29	0 – 1000	mocr02	1,543,340	1,396,180	17
gn13e39d2w0w1000	13	39	0 – 1000	mocr02	131,807,934	15,037,589	26

TABLE 4.10. Calibration test set: **Heuristic 91** (Boundary and Triangular Region Pareto Test) with 2 criteria. Internally, **Heuristic 86** (PT) used 10 searches per point, 100 pivot limit.

Name	Nodes	Edges	Weights	Computer	Seconds	# Pareto Optimum	#Computed Pareto Optimum	Percent
gn9e18d2w0w20	9	18	0 – 20	Fuzzy	40	15	15	100.00
gn9e27d2w0w20	9	27	0 – 20	Fuzzy	11	19	19	100.00
gn10e22d2w0w20	10	22	0 – 20	Fuzzy	29	19	19	100.00
gn10e28d2w0w20	10	28	0 – 20	Fuzzy	9	10	10	100.00
gn10e33d2w0w20	10	33	0 – 20	Fuzzy	59	21	21	100.00
gn11e13d2w0w20	11	13	0 – 20	Fuzzy	1	2	2	100.00
gn11e20d2w0w20	11	20	0 – 20	Fuzzy	34	17	17	100.00
gn11e27d2w0w20	11	27	0 – 20	Fuzzy	96	28	28	100.00
gn11e41d2w0w20	11	41	0 – 20	Fuzzy	96	17	17	100.00
gn12e16d2w0w20	12	16	0 – 20	Fuzzy	11	5	5	100.00
gn12e24d2w0w20	12	24	0 – 20	Fuzzy	48	17	17	100.00
gn12e33d2w0w20	12	33	0 – 20	Fuzzy	29	24	24	100.00
gn13e19d2w0w20	13	19	0 – 20	Fuzzy	44	13	13	100.00
gn13e29d2w0w20	13	29	0 – 20	Fuzzy	50	20	20	100.00
gn13e39d2w0w20	13	39	0 – 20	Fuzzy	33	19	19	100.00
gn9e18d2w0w100	9	18	0 – 100	Fuzzy	350	9	9	100.00
gn9e27d2w0w100	9	27	0 – 100	Fuzzy	1,481	43	43	100.00
gn10e22d2w0w100	10	22	0 – 100	Fuzzy	1,286	21	21	100.00
gn10e28d2w0w100	10	28	0 – 100	Fuzzy	1,251	28	28	100.00
gn10e33d2w0w100	10	33	0 – 100	Fuzzy	1,625	23	23	100.00
gn11e13d2w0w100	11	13	0 – 100	Fuzzy	376	4	4	100.00
gn11e20d2w0w100	11	20	0 – 100	Fuzzy	435	6	6	100.00
gn11e27d2w0w100	11	27	0 – 100	Fuzzy	2,002	25	25	100.00
gn11e41d2w0w100	11	41	0 – 100	Fuzzy	2,472	28	28	100.00
gn12e16d2w0w100	12	16	0 – 100	Fuzzy	155	6	6	100.00
gn12e24d2w0w100	12	24	0 – 100	Fuzzy	3,290	24	24	100.00
gn12e33d2w0w100	12	33	0 – 100	Fuzzy	2,493	37	37	100.00
gn13e19d2w0w100	13	19	0 – 100	Fuzzy	829	18	18	100.00
gn13e29d2w0w100	13	29	0 – 100	Fuzzy	3,240	39	39	100.00
gn13e39d2w0w100	13	39	0 – 100	Fuzzy	3,328	33	33	100.00

TABLE 4.11. Solids, **Heuristic 92** (DFBFS). 100 searches, depth 4, boundary retry limit 100 and interior retry limit of 10000. Note that Matsui's algorithm was only able to enumerate the spanning trees of the first six graphs.

Name	Nodes	Edges	Weight range	Computer	Seconds	#Spanning Trees	#Projected Trees	#Trees Found by DFBFS	Percent
Solid-tetrahedron-d2w0w20	4	6	0 - 20	mocr04	0	16	16	16	100.00
Solid-cube-d2w0w20	4	6	0 - 20	mocr04	0	384	321	320	99.69
Solid-ocahedron-d2w0w20	6	12	0 - 20	mocr04	0	384	296	292	98.65
Solid-truncatedtetrahedron-d2w0w20	12	18	0 - 20	mocr04	0	6,000	1,947	1,931	99.18
Solid-cuboctahedron-d2w0w20	12	24	0 - 20	mocr04	2	331,776	6,291	6,209	98.70
Solid-icosahedron-d2w0w20	12	30	0 - 20	mocr04	3	5,184,000	12,190	11,911	97.71
Solid-dodecahedron-d2w0w20	20	30	0 - 20	mocr04	4	5,184,000	11,708	11,625	99.29
Solid-truncatedoctahedron-d2w0w20	24	36	0 - 20	mocr04	11	101,154,816		16,882	
Solid-srhombicuboctahedron-d2w0w20	24	48	0 - 20	mocr04	31	301 × 10 ⁹		35,132	
Solid-snubcube-d2w0w20	24	60	0 - 20	mocr04	56	89 × 10 ¹²		56,992	
Solid-icosidodecahedron-d2w0w20	30	60	0 - 20	mocr04	78	208 × 10 ¹²		61,926	
Solid-grhombicuboctahedron-d2w0w20	48	72	0 - 20	mocr04	201	12 × 10 ¹⁵		73,166	
Solid-truncatedicosahedron-d2w0w20	60	90	0 - 20	mocr04	623	375 × 10 ¹⁸		112,292	
Solid-rhomicosidodecahedron-d2w0w20	60	120	0 - 20	mocr04	1,266	201 × 10 ²⁷		166,622	
Solid-snubdodecahedron-d2w0w20	60	150	0 - 20	mocr04	1,955	438 × 10 ³³		199,764	
Solid-tetrahedron-d2w0w100	4	6	0 - 100	mocr04	1	16	16	16	100.00
Solid-octahedron-d2w0w100	6	12	0 - 100	mocr04	0	384	383	380	99.22
Solid-cube-d2w0w100	8	12	0 - 100	mocr04	0	384	383	383	100.00
Solid-truncatedtetrahedron-d2w0w100	12	18	0 - 100	mocr04	1	6,000	5,812	5,782	99.48
Solid-cuboctahedron-d2w0w100	12	24	0 - 100	mocr04	13	331,776	93,982	90,788	96.60
Solid-icosahedron-d2w0w100	12	30	0 - 100	mocr04	27	5,184,000	207,447	200,641	96.72
Solid-dodecahedron-d2w0w100	20	30	0 - 100	mocr04	49	5,184,000	217,882	211,169	96.92
Solid-truncatedoctahedron-d2w0w100	24	36	0 - 100	mocr04	102	101,154,816	353,262	339,369	96.07
Solid-srhombicuboctahedron-d2w0w100	24	48	0 - 100	mocr04	233	301 × 10 ⁹		793,894	
Solid-snubcube-d2w0w100	24	60	0 - 100	mocr04	322	89 × 10 ¹²		1,017,998	
Solid-icosidodecahedron-d2w0w100	30	60	0 - 100	mocr04	518	208 × 10 ¹²		1,335,581	
Solid-grhombicuboctahedron-d2w0w100	48	72	0 - 100	mocr04	1409	12 × 10 ¹⁵		1,554,008	
Solid-truncatedicosahedron-d2w0w100	60	90	0 - 100	mocr04	3005	375 × 10 ¹⁸		2,440,020	
Solid-rhomicosidodecahedron-d2w0w100	60	120	0 - 100	mocr04	5772	201 × 10 ²⁷		4,327,586	
Solid-snubdodecahedron-d2w0w100	60	150	0 - 100	mocr04	7878	438 × 10 ³³		5,278,828	
Solid-tetrahedron-d2w0w1000	4	6	0 - 1000	mocr04	0	16	16	16	100.00
Solid-octahedron-d2w0w1000	6	12	0 - 1000	mocr04	1	384	384	376	97.92
Solid-cube-d2w0w1000	8	12	0 - 1000	mocr04	0	384	384	383	99.74
Solid-truncatedtetrahedron-d2w0w1000	12	18	0 - 1000	mocr04	1	6,000	6,000	5,662	94.37
Solid-cuboctahedron-d2w0w1000	12	24	0 - 1000	mocr04	34	331,776	331,776	318,156	95.89
Solid-icosahedron-d2w0w1000	12	30	0 - 1000	mocr04	328	5,184,000	5,184,000	3,263,800	62.96
Solid-dodecahedron-d2w0w1000	20	30	0 - 1000	mocr04	628	5,184,000	5,184,000	3,606,194	69.56
Solid-truncatedoctahedron-d2w0w1000	24	36	0 - 1000	mocr04	2,637	101,154,816	101,154,816	13,024,774	12.88
Solid-srhombicuboctahedron-d2w0w1000	24	48	0 - 1000	mocr04	9,354	301 × 10 ⁹		48,407,588	
Solid-snubcube-d2w0w1000	24	60	0 - 1000	mocr04	32,481	89 × 10 ¹²		83,600,261	

TABLE 4.12. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 2$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d2w0w20	2,981	1,742	58.44	91	0
gn9e27d2w0w20	372,320	6,346	01.70	167	1
gn10e22d2w0w20	53,357	3,957	07.42	316	1
gn10e28d2w0w20	800,948	7,131	00.89	252	2
gn10e33d2w0w20	3,584,016	10,833	00.30	172	1
gn11e13d2w0w20	41	41	100.00	438	1
gn11e20d2w0w20	6,939	2,173	31.32	135	1
gn11e27d2w0w20	284,730	7,515	02.64	252	2
gn11e41d2w0w20	90,922,271	16,457	00.02	190	5
gn12e16d2w0w20	162	154	95.06	546	0
gn12e24d2w0w20	208,380	5,647	02.71	98	2
gn12e33d2w0w20	4,741,624	9,364	00.20	111	4
gn13e19d2w0w20	3,401	1,608	47.28	38	1
gn13e29d2w0w20	1,543,340	8,474	00.55	100	4
gn13e39d2w0w20	131,807,934	18,468	00.01	244	4
gn9e18d2w0w100	2,981	2,858	95.87	206	0
gn9e27d2w0w100	372,320	89,092	23.93	111	2
gn10e22d2w0w100	53,357	37,204	69.73	5	1
gn10e28d2w0w100	800,948	101,334	12.65	0	2
gn10e33d2w0w100	3,584,016	166,427	04.64	12	3
gn11e13d2w0w100	41	41	100.00	1000	1
gn11e20d2w0w100	6,939	6,580	94.83	21	1
gn11e27d2w0w100	284,730	81,803	28.73	24	2
gn11e41d2w0w100	90,922,271	309,961	00.34	6	4
gn12e16d2w0w100	162	162	100.00	227	0
gn12e24d2w0w100	208,380	92,813	44.54	0	2
gn12e33d2w0w100	4,741,624	192,122	04.05	19	3
gn13e19d2w0w100	3,401	3,255	95.71	158	1
gn13e29d2w0w100	1,543,340	164,617	10.67	25	4
gn13e39d2w0w100	131,807,934	315,881	00.24	13	5
gn9e18d2w0w1000	2,981	2,981	100.00	49	0
gn9e27d2w0w1000	372,320	364,382	97.87	0	1
gn10e22d2w0w1000	53,357	52,990	99.31	5	1
gn10e28d2w0w1000	800,948	756,013	94.39	0	1
gn10e33d2w0w1000	3,584,016	2,726,287	76.07	0	2
gn11e13d2w0w1000	41	41	100.00	1000	1
gn11e20d2w0w1000	6,939	6,921	99.74	35	1
gn11e27d2w0w1000	284,730	279,308	98.10	0	2
gn11e41d2w0w1000	90,922,271	13,884,793	15.27	0	2
gn12e16d2w0w1000	162	162	100.00	484	1
gn12e24d2w0w1000	208,380	205,690	98.71	0	2
gn12e33d2w0w1000	4,741,624	3,680,313	77.62	0	3
gn13e19d2w0w1000	3,401	3,401	100.00	11	2
gn13e29d2w0w1000	1,543,340	1,396,180	90.46	5	3
gn13e39d2w0w1000	131,807,934	15,037,589	11.41	0	3

TABLE 4.13. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 3$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d3w0w20	2,981	2,905	97.45	187	1
gn9e27d3w0w20	372,320	134,039	36.00	2	2
gn10e22d3w0w20	53,357	42,887	80.38	0	2
gn10e28d3w0w20	800,948	238,529	29.78	0	3
gn10e33d3w0w20	3,584,016	411,730	11.49	0	3
gn11e13d3w0w20	41	41	100.00	389	0
gn11e20d3w0w20	6,939	6,603	95.16	0	2
gn11e27d3w0w20	284,730	146,672	51.51	60	2
gn11e41d3w0w20	90,922,271	959,469	01.06	8	5
gn12e16d3w0w20	162	162	100.00	677	1
gn12e24d3w0w20	208,380	111,201	53.36	13	3
gn12e33d3w0w20	4,741,624	470,609	09.93	18	3
gn13e19d3w0w20	3,401	3,358	98.74	106	2
gn13e29d3w0w20	1,543,340	264,949	17.18	18	4
gn13e39d3w0w20	131,807,934	930,322	00.71	7	4
gn9e18d3w0w100	2,981	2,981	100.00	32	1
gn9e27d3w0w100	372,320	367,313	98.66	0	2
gn10e22d3w0w100	53,357	53,289	99.87	29	1
gn10e28d3w0w100	800,948	786,781	98.23	0	2
gn10e33d3w0w100	3,584,016	3,351,096	93.01	0	3
gn11e13d3w0w100	41	41	100.00	834	1
gn11e20d3w0w100	6,939	6,939	100.00	181	2
gn11e27d3w0w100	284,730	281,303	98.80	3	1
gn11e41d3w0w100	90,922,271	35,943,327	39.53	0	4
gn12e16d3w0w100	162	162	100.00	568	1
gn12e24d3w0w100	208,380	207,143	99.41	0	2
gn12e33d3w0w100	4,741,624	4,740,880	99.98	0	6
gn13e19d3w0w100	3,401	3,401	100.00	30	2
gn13e29d3w0w100	1,543,340	1,484,719	96.20	0	3
gn13e39d3w0w100	131,807,934	44,757,592	33.96	0	6
gn9e18d3w0w1000	2,981	2,981	100.00	260	1
gn9e27d3w0w1000	372,320	372,320	100.00	0	2
gn10e22d3w0w1000	53,357	53,357	100.00	12	1
gn10e28d3w0w1000	800,948	800,946	99.99	0	2
gn10e33d3w0w1000	3,584,016	3,583,757	99.99	0	2
gn11e13d3w0w1000	41	41	100.00	1000	1
gn11e20d3w0w1000	6,939	6,939	100.00	24	1
gn11e27d3w0w1000	284,730	284,730	100.00	0	2
gn11e41d3w0w1000	90,922,271	90,699,181	99.75	0	3
gn12e16d3w0w1000	162	162	100.00	504	0
gn12e24d3w0w1000	208,380	208,356	99.99	13	2
gn12e33d3w0w1000	4,741,624	4,740,880	99.98	0	2
gn13e19d3w0w1000	3,401	3,401	100.00	101	1
gn13e29d3w0w1000	1,543,340	1,543,304	99.99	1	3
gn13e39d3w0w1000	131,807,934	131,464,478	99.74	0	3

TABLE 4.14. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 5$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d5w0w1	2,981	1,099	36.87	160	1
gn9e27d5w0w1	372,320	8,205	2.20	333	2
gn10e22d5w0w1	53,357	4,746	8.89	197	2
gn10e28d5w0w1	800,948	10,898	1.36	122	3
gn10e33d5w0w1	3,584,016	15,482	0.43	110	4
gn11e13d5w0w1	41	27	65.85	517	0
gn11e20d5w0w1	6,939	1,866	26.89	241	2
gn11e27d5w0w1	284,730	5,828	2.05	178	2
gn11e41d5w0w1	90,922,271	36,847	0.04	195	6
gn12e16d5w0w1	162	116	71.61	748	1
gn12e24d5w0w1	208,380	6,738	3.23	529	2
gn12e33d5w0w1	4,741,624	20,857	0.44	353	5
gn13e19d5w0w1	3,401	1,149	33.78	685	2
gn13e29d5w0w1	1,543,340	111,116	7.20	395	4
gn13e39d5w0w1	131,807,934	34,392	0.03	203	6
gn9e18d5w0w2	2,981	2,276	76.35	42	1
gn9e27d5w0w2	372,320	43,029	11.55	159	3
gn10e22d5w0w2	53,357	14,623	27.40	284	2
gn10e28d5w0w2	800,948	66,190	8.26	112	2
gn10e33d5w0w2	3,584,016	105,309	2.93	53	4
gn11e13d5w0w2	41	41	100.00	523	1
gn11e20d5w0w2	6,939	3,761	54.20	132	1
gn11e27d5w0w2	284,730	39,292	13.79	228	2
gn11e41d5w0w2	90,922,271	290,555	0.31	394	4
gn12e16d5w0w2	162	160	98.76	645	1
gn12e24d5w0w2	208,380	34,057	16.34	236	3
gn12e33d5w0w2	4,741,624	120,211	2.53	49	4
gn13e19d5w0w2	3,401	2,613	76.83	65	2
gn13e29d5w0w2	1,543,340	98,285	6.36	146	4
gn13e39d5w0w2	131,807,934	348,703	0.26	352	4
gn9e18d5w0w5	2,981	2,960	99.29	229	1
gn9e27d5w0w5	372,320	271,048	72.79	0	2
gn10e22d5w0w5	53,357	49,463	92.70	94	2
gn10e28d5w0w5	800,948	493,565	61.62	10	2
gn10e33d5w0w5	3,584,016	1,294,875	36.12	16	3
gn11e13d5w0w5	41	41	100.00	1000	1
gn11e20d5w0w5	6,939	5,975	86.10	15	1
gn11e27d5w0w5	284,730	222,974	78.31	0	3
gn11e41d5w0w5	90,922,271	4,711,354	5.18	0	7
gn12e16d5w0w5	162	162	100.00	604	2
gn12e24d5w0w5	208,380	177,845	85.34	42	3
gn12e33d5w0w5	4,741,624	1,489,751	31.41	10	4
gn13e19d5w0w5	3,401	3,381	99.41	67	1
gn13e29d5w0w5	1,543,340	787,196	51.00	52	4
gn13e39d5w0w5	131,807,934	7,737,684	5.87	0	6

TABLE 4.15. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 2$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d2w0w20	2,981	1,742	58.44	77	1
gn9e27d2w0w20	372,320	6,346	01.70	176	2
gn10e22d2w0w20	53,357	3,957	07.42	362	1
gn10e28d2w0w20	800,948	7,131	00.89	273	2
gn10e33d2w0w20	3,584,016	10,833	00.30	175	1
gn11e13d2w0w20	41	41	100.00	453	1
gn11e20d2w0w20	6,939	2,173	31.32	75	1
gn11e27d2w0w20	284,730	7,515	02.64	249	1
gn11e41d2w0w20	90,922,271	16,457	00.02	217	5
gn12e16d2w0w20	162	154	95.06	573	1
gn12e24d2w0w20	208,380	5,647	02.71	119	2
gn12e33d2w0w20	4,741,624	9,364	00.20	119	4
gn13e19d2w0w20	3,401	1,608	47.28	42	1
gn13e29d2w0w20	1,543,340	8,474	00.55	118	4
gn13e39d2w0w20	131,807,934	18,468	00.01	228	3
gn9e18d2w0w100	2,981	2,858	95.87	85	1
gn9e27d2w0w100	372,320	89,092	23.93	72	2
gn10e22d2w0w100	53,357	37,204	69.73	3	1
gn10e28d2w0w100	800,948	101,334	12.65	1	2
gn10e33d2w0w100	3,584,016	166,427	04.64	31	3
gn11e13d2w0w100	41	41	100.00	1000	0
gn11e20d2w0w100	6,939	6,580	94.83	18	1
gn11e27d2w0w100	284,730	81,803	28.73	32	2
gn11e41d2w0w100	90,922,271	309,961	00.34	2	5
gn12e16d2w0w100	162	162	100.00	239	1
gn12e24d2w0w100	208,380	92,813	44.54	0	2
gn12e33d2w0w100	4,741,624	192,122	04.05	18	3
gn13e19d2w0w100	3,401	3,255	95.71	193	2
gn13e29d2w0w100	1,543,340	164,617	10.67	11	4
gn13e39d2w0w100	131,807,934	315,881	00.24	4	4
gn9e18d2w0w1000	2,981	2,981	100.00	63	1
gn9e27d2w0w1000	372,320	364,382	97.87	0	1
gn10e22d2w0w1000	53,357	52,990	99.31	3	2
gn10e28d2w0w1000	800,948	756,013	94.39	0	1
gn10e33d2w0w1000	3,584,016	2,726,287	76.07	0	2
gn11e13d2w0w1000	41	41	100.00	1000	0
gn11e20d2w0w1000	6,939	6,921	99.74	35	1
gn11e27d2w0w1000	284,730	279,308	98.10	0	2
gn11e41d2w0w1000	90,922,271	13,884,793	15.27	0	2
gn12e16d2w0w1000	162	162	100.00	485	1
gn12e24d2w0w1000	208,380	205,690	98.71	0	2
gn12e33d2w0w1000	4,741,624	3,680,313	77.62	0	2
gn13e19d2w0w1000	3,401	3,401	100.00	24	1
gn13e29d2w0w1000	1,543,340	1,396,180	90.46	5	3
gn13e39d2w0w1000	131,807,934	15,037,589	11.41	0	4

TABLE 4.16. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 3$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d3w0w20	2,981	2,905	97.45	275	1
gn9e27d3w0w20	372,320	134,039	36.00	0	2
gn10e22d3w0w20	53,357	42,887	80.38	0	1
gn10e28d3w0w20	800,948	238,529	29.78	0	2
gn10e33d3w0w20	3,584,016	411,730	11.49	0	2
gn11e13d3w0w20	41	41	100.00	323	0
gn11e20d3w0w20	6,939	6,603	95.16	0	2
gn11e27d3w0w20	284,730	146,672	51.51	40	2
gn11e41d3w0w20	90,922,271	959,469	01.06	0	4
gn12e16d3w0w20	162	162	100.00	681	1
gn12e24d3w0w20	208,380	111,201	53.36	9	3
gn12e33d3w0w20	4,741,624	470,609	09.93	13	2
gn13e19d3w0w20	3,401	3,358	98.74	55	2
gn13e29d3w0w20	1,543,340	264,949	17.18	11	4
gn13e39d3w0w20	131,807,934	930,322	00.71	20	4
gn9e18d3w0w100	2,981	2,981	100.00	40	1
gn9e27d3w0w100	372,320	367,313	98.66	19	2
gn10e22d3w0w100	53,357	53,289	99.87	52	2
gn10e28d3w0w100	800,948	786,781	98.23	0	2
gn10e33d3w0w100	3,584,016	3,351,096	93.01	0	3
gn11e13d3w0w100	41	41	100.00	849	0
gn11e20d3w0w100	6,939	6,939	100.00	129	1
gn11e27d3w0w100	284,730	281,303	98.80	26	2
gn11e41d3w0w100	90,922,271	35,943,327	39.53	0	3
gn12e16d3w0w100	162	162	100.00	473	1
gn12e24d3w0w100	208,380	207,143	99.41	0	2
gn12e33d3w0w100	4,741,624	4,740,880	99.98	0	5
gn13e19d3w0w100	3,401	3,401	100.00	17	1
gn13e29d3w0w100	1,543,340	1,484,719	96.20	0	2
gn13e39d3w0w100	131,807,934	44,757,592	33.96	0	6
gn9e18d3w0w1000	2,981	2,981	100.00	136	1
gn9e27d3w0w1000	372,320	372,320	100.00	0	1
gn10e22d3w0w1000	53,357	53,357	100.00	25	2
gn10e28d3w0w1000	800,948	800,946	99.99	0	2
gn10e33d3w0w1000	3,584,016	3,583,757	99.99	0	2
gn11e13d3w0w1000	41	41	100.00	1000	1
gn11e20d3w0w1000	6,939	6,939	100.00	21	2
gn11e27d3w0w1000	284,730	284,730	100.00	0	3
gn11e41d3w0w1000	90,922,271	90,699,181	99.75	0	3
gn12e16d3w0w1000	162	162	100.00	509	1
gn12e24d3w0w1000	208,380	208,356	99.99	26	2
gn12e33d3w0w1000	4,741,624	4,740,880	99.98	0	2
gn13e19d3w0w1000	3,401	3,401	100.00	127	2
gn13e29d3w0w1000	1,543,340	1,543,304	99.99	0	2
gn13e39d3w0w1000	131,807,934	131,464,478	99.74	0	4

TABLE 4.17. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 5$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d5w0w1	2,981	1,099	36.87	170	1
gn9e27d5w0w1	372,320	8,205	2.20	338	2
gn10e22d5w0w1	53,357	4,746	8.89	191	2
gn10e28d5w0w1	800,948	10,898	1.36	156	3
gn10e33d5w0w1	3,584,016	15,482	0.43	114	4
gn11e13d5w0w1	41	27	65.85	555	1
gn11e20d5w0w1	6,939	1,866	26.89	224	2
gn11e27d5w0w1	284,730	5,828	2.05	181	3
gn11e41d5w0w1	90,922,271	36,847	0.04	213	6
gn12e16d5w0w1	162	116	71.61	755	1
gn12e24d5w0w1	208,380	6,738	3.23	502	2
gn12e33d5w0w1	4,741,624	20,857	0.44	375	5
gn13e19d5w0w1	3,401	1,149	33.78	694	3
gn13e29d5w0w1	1,543,340	111,116	7.20	390	4
gn13e39d5w0w1	131,807,934	34,392	0.03	252	6
gn9e18d5w0w2	2,981	2,276	76.35	92	0
gn9e27d5w0w2	372,320	43,029	11.55	168	2
gn10e22d5w0w2	53,357	14,623	27.40	272	3
gn10e28d5w0w2	800,948	66,190	8.26	134	3
gn10e33d5w0w2	3,584,016	105,309	2.93	72	3
gn11e13d5w0w2	41	41	100.00	503	1
gn11e20d5w0w2	6,939	3,761	54.20	130	1
gn11e27d5w0w2	284,730	39,292	13.79	238	2
gn11e41d5w0w2	90,922,271	290,555	0.31	390	5
gn12e16d5w0w2	162	160	98.76	668	1
gn12e24d5w0w2	208,380	34,057	16.34	280	3
gn12e33d5w0w2	4,741,624	120,211	2.53	58	4
gn13e19d5w0w2	3,401	2,613	76.83	61	1
gn13e29d5w0w2	1,543,340	98,285	6.36	196	4
gn13e39d5w0w2	131,807,934	348,703	0.26	374	4
gn9e18d5w0w5	2,981	2,960	99.29	89	1
gn9e27d5w0w5	372,320	271,048	72.79	0	2
gn10e22d5w0w5	53,357	49,463	92.70	56	2
gn10e28d5w0w5	800,948	493,565	61.62	41	2
gn10e33d5w0w5	3,584,016	1,294,875	36.12	9	4
gn11e13d5w0w5	41	41	100.00	1000	1
gn11e20d5w0w5	6,939	5,975	86.10	26	2
gn11e27d5w0w5	284,730	222,974	78.31	0	3
gn11e41d5w0w5	90,922,271	4,711,354	5.18	12	6
gn12e16d5w0w5	162	162	100.00	795	1
gn12e24d5w0w5	208,380	177,845	85.34	41	3
gn12e33d5w0w5	4,741,624	1,489,751	31.41	3	5
gn13e19d5w0w5	3,401	3,381	99.41	77	2
gn13e29d5w0w5	1,543,340	787,196	51.00	52	4
gn13e39d5w0w5	131,807,934	7,737,684	5.87	0	7

TABLE 4.18. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 2$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d2w0w20	0(0)	66(1)	295(1)	295(1)	348(2)	383(2)	484(3)	499(2)	600(3)	619(3)	656(5)	976(13)	1000(22)
gn9e27d2w0w20	5(0)	172(1)	271(2)	327(2)	466(2)	506(2)	573(3)	515(4)	566(4)	648(4)	796(8)	929(17)	1000(33)
gn10e22d2w0w20	56(0)	349(1)	400(1)	448(1)	503(2)	671(2)	661(3)	709(4)	706(3)	717(4)	800(6)	915(14)	915(27)
gn10e28d2w0w20	4(0)	275(1)	292(2)	315(3)	331(3)	395(4)	470(4)	438(4)	469(4)	516(5)	717(9)	985(23)	1000(39)
gn10e33d2w0w20	18(1)	167(1)	237(2)	360(3)	396(2)	450(3)	457(3)	535(4)	642(4)	664(5)	765(10)	948(23)	986(42)
gn11e13d2w0w20	324(0)	440(0)	466(0)	623(1)	613(1)	628(2)	642(2)	785(2)	883(3)	1000(3)	1000(5)	1000(9)	1000(18)
gn11e20d2w0w20	0(1)	135(1)	145(1)	272(2)	313(2)	315(3)	306(3)	328(3)	311(3)	300(4)	669(8)	744(15)	745(27)
gn11e27d2w0w20	17(1)	236(1)	316(2)	383(2)	400(2)	440(3)	495(4)	528(4)	594(5)	639(5)	811(9)	932(20)	991(39)
gn11e41d2w0w20	0(1)	189(4)	215(4)	290(4)	368(5)	381(6)	419(7)	456(8)	545(8)	587(10)	807(17)	956(35)	1000(62)
gn12e16d2w0w20	143(0)	579(0)	549(0)	590(1)	715(2)	795(2)	871(2)	867(2)	923(3)	914(3)	901(5)	913(11)	904(22)
gn12e24d2w0w20	10(1)	100(1)	137(2)	185(2)	206(3)	265(4)	284(4)	297(4)	326(5)	349(5)	385(10)	505(21)	531(39)
gn12e33d2w0w20	0(0)	128(2)	131(3)	128(4)	142(5)	155(5)	153(5)	138(6)	160(7)	146(8)	249(14)	792(31)	815(52)
gn13e19d2w0w20	12(1)	34(1)	77(1)	148(2)	218(3)	204(3)	225(3)	206(3)	248(4)	250(4)	443(8)	459(18)	450(32)
gn13e29d2w0w20	0(0)	117(3)	284(4)	278(5)	298(5)	335(6)	294(6)	340(7)	301(7)	301(8)	302(12)	631(32)	1000(67)
gn13e39d2w0w20	5(1)	201(3)	327(3)	426(4)	456(4)	484(5)	553(7)	614(7)	646(8)	662(8)	850(16)	967(35)	1000(66)
gn9e18d2w0w100	0(0)	180(1)	183(1)	290(1)	563(2)	556(2)	658(2)	759(3)	762(2)	1000(3)	1000(5)	1000(11)	1000(20)
gn9e27d2w0w100	0(1)	95(1)	91(2)	100(3)	97(3)	79(4)	96(3)	88(4)	90(4)	79(5)	82(9)	87(19)	80(33)
gn10e22d2w0w100	0(0)	9(1)	4(1)	12(2)	7(2)	5(3)	4(3)	4(3)	4(3)	3(4)	3(8)	16(18)	19(38)
gn10e28d2w0w100	0(1)	0(2)	0(2)	0(2)	2(3)	7(3)	7(4)	5(4)	5(5)	20(6)	18(12)	28(26)	37(53)
gn10e33d2w0w100	0(0)	4(2)	6(2)	7(3)	30(4)	42(4)	50(5)	48(5)	52(5)	55(6)	130(10)	190(24)	417(52)
gn11e13d2w0w100	309(0)	1000(1)	1000(0)	1000(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(2)	1000(2)	1000(4)	1000(8)	1000(18)
gn11e20d2w0w100	1(0)	19(1)	40(1)	34(1)	149(2)	148(2)	174(3)	201(3)	191(4)	168(4)	252(8)	375(19)	748(43)
gn11e27d2w0w100	0(0)	33(2)	27(2)	35(3)	26(3)	31(4)	27(4)	38(4)	35(5)	31(6)	30(11)	33(27)	34(53)
gn11e41d2w0w100	0(0)	2(3)	5(4)	2(4)	0(5)	2(5)	6(7)	6(8)	2(8)	3(9)	44(17)	32(42)	47(74)
gn12e16d2w0w100	81(1)	228(0)	354(1)	326(1)	360(1)	382(2)	901(3)	895(2)	893(3)	921(3)	1000(6)	1000(12)	1000(23)
gn12e24d2w0w100	0(0)	0(2)	0(2)	1(3)	5(3)	7(4)	4(5)	11(5)	11(6)	13(7)	70(12)	251(32)	316(55)
gn12e33d2w0w100	0(1)	14(2)	22(3)	20(3)	24(4)	32(5)	71(5)	76(7)	87(8)	113(9)	143(16)	209(38)	202(67)
gn13e19d2w0w100	2(0)	156(1)	154(2)	161(2)	146(2)	161(3)	158(3)	165(3)	164(3)	157(4)	263(7)	257(16)	283(31)
gn13e29d2w0w100	0(0)	33(3)	28(4)	24(4)	59(4)	35(5)	36(6)	49(6)	47(7)	53(8)	45(12)	278(33)	272(53)
gn13e39d2w0w100	0(1)	14(3)	18(4)	20(5)	29(6)	41(7)	35(8)	63(9)	60(10)	66(11)	113(19)	256(45)	368(94)
gn9e18d2w0w1000	34(1)	40(1)	60(1)	64(2)	125(2)	128(2)	128(2)	145(2)	150(4)	143(3)	110(6)	443(15)	460(24)
gn9e27d2w0w1000	0(0)	0(1)	0(1)	0(2)	0(2)	0(2)	0(3)	0(4)	0(4)	0(5)	17(10)	23(25)	63(49)
gn10e22d2w0w1000	0(0)	5(1)	7(2)	10(2)	35(2)	30(2)	35(3)	55(4)	48(4)	51(5)	68(10)	65(22)	58(44)
gn10e28d2w0w1000	0(0)	0(1)	0(2)	0(3)	0(3)	0(3)	0(4)	0(5)	0(5)	0(6)	8(11)	11(27)	23(51)
gn10e33d2w0w1000	0(1)	0(2)	0(2)	0(2)	0(2)	0(3)	0(3)	0(5)	0(5)	0(6)	0(13)	6(32)	2(67)
gn11e13d2w0w1000	215(0)	1000(1)	1000(0)	1000(1)	1000(1)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(4)	1000(9)	1000(17)
gn11e20d2w0w1000	0(0)	29(1)	24(2)	93(2)	101(2)	93(3)	102(3)	99(3)	109(4)	102(4)	372(9)	687(18)	757(34)
gn11e27d2w0w1000	0(1)	0(1)	0(2)	0(2)	0(3)	0(3)	0(4)	0(5)	0(5)	0(5)	1(11)	28(29)	53(61)
gn11e41d2w0w1000	0(1)	0(1)	0(2)	0(3)	0(4)	0(4)	0(6)	0(6)	0(8)	0(8)	2(17)	2(40)	15(87)
gn12e16d2w0w1000	114(0)	474(1)	512(1)	542(2)	887(1)	1000(2)	1000(2)	1000(2)	1000(3)	1000(3)	1000(5)	1000(12)	1000(22)
gn12e24d2w0w1000	0(0)	0(1)	1(2)	14(3)	12(3)	7(4)	3(4)	21(4)	19(6)	17(6)	30(12)	119(28)	192(56)
gn12e33d2w0w1000	0(0)	0(1)	0(2)	0(3)	0(4)	0(5)	0(6)	0(6)	0(7)	0(7)	0(16)	0(40)	0(77)
gn13e19d2w0w1000	21(1)	33(1)	49(1)	33(2)	46(3)	28(3)	56(4)	44(4)	37(4)	40(5)	137(11)	235(25)	501(49)
gn13e29d2w0w1000	0(1)	8(2)	5(3)	9(4)	27(4)	23(5)	29(6)	30(7)	23(9)	24(9)	61(16)	369(44)	626(84)
gn13e39d2w0w1000	0(0)	0(3)	0(3)	0(4)	0(5)	0(7)	1(7)	0(8)	0(9)	1(10)	0(19)	56(49)	64(98)

TABLE 4.19. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 3$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d3w0w1000	22(1)	204(0)	311(1)	300(1)	298(2)	341(2)	313(2)	313(2)	296(3)	330(3)	620(5)	1000(14)	1000(23)
gn9e27d3w0w1000	0(0)	3(1)	0(1)	33(2)	26(2)	30(3)	45(4)	52(4)	47(5)	63(5)	123(12)	226(24)	323(44)
gn10e22d3w0w1000	0(0)	2(1)	1(1)	3(2)	4(3)	1(3)	5(3)	5(4)	7(4)	7(5)	19(9)	64(24)	141(39)
gn10e28d3w0w1000	0(0)	0(2)	0(2)	10(3)	17(4)	20(4)	21(4)	29(5)	37(5)	25(6)	44(13)	78(30)	185(56)
gn10e33d3w0w1000	0(1)	0(2)	0(3)	0(4)	0(4)	3(5)	1(5)	1(5)	1(7)	1(7)	2(13)	30(28)	24(48)
gn11e13d3w0w1000	271(1)	390(1)	461(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(2)	1000(2)	1000(3)	1000(4)	1000(9)	1000(19)
gn11e20d3w0w1000	0(0)	0(1)	0(1)	0(2)	0(2)	0(3)	0(3)	8(3)	44(3)	37(4)	45(8)	456(22)	907(46)
gn11e27d3w0w1000	0(0)	71(2)	49(3)	66(3)	53(4)	60(4)	47(4)	63(5)	67(6)	59(7)	115(11)	149(26)	143(56)
gn11e41d3w0w1000	0(1)	3(3)	7(4)	16(5)	11(6)	10(6)	11(7)	8(8)	16(9)	14(10)	43(20)	78(49)	77(85)
gn12e16d3w0w1000	45(0)	707(1)	687(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(3)	1000(3)	1000(3)	1000(6)	1000(12)	1000(24)
gn12e24d3w0w1000	0(0)	9(2)	15(2)	16(3)	7(3)	9(4)	18(5)	11(5)	20(6)	12(7)	18(13)	22(26)	23(46)
gn12e33d3w0w1000	4(1)	13(2)	20(2)	18(3)	16(4)	30(4)	28(5)	25(6)	25(7)	34(7)	76(16)	138(36)	246(65)
gn13e19d3w0w1000	0(0)	116(1)	149(2)	170(3)	323(3)	443(3)	469(5)	671(6)	671(6)	745(6)	1000(12)	1000(21)	1000(37)
gn13e29d3w0w1000	0(1)	18(3)	33(3)	43(4)	44(5)	34(6)	44(7)	43(7)	45(8)	52(8)	155(16)	133(40)	239(73)
gn13e39d3w0w1000	0(1)	10(3)	16(3)	22(5)	22(6)	28(6)	41(7)	45(9)	47(10)	57(11)	89(20)	246(46)	352(84)
gn9e18d3w0w100	0(0)	43(1)	29(1)	31(1)	33(1)	37(2)	37(3)	38(3)	35(2)	32(3)	46(4)	37(13)	34(24)
gn9e27d3w0w100	0(0)	0(1)	0(2)	0(2)	41(3)	40(4)	35(5)	64(5)	79(5)	74(6)	147(11)	256(26)	752(55)
gn10e22d3w0w100	0(1)	35(1)	76(2)	84(3)	113(2)	119(3)	122(4)	224(4)	218(4)	232(5)	272(9)	897(23)	984(40)
gn10e28d3w0w100	0(0)	0(1)	0(2)	0(3)	0(3)	0(4)	0(5)	0(5)	0(6)	0(5)	0(12)	0(29)	0(51)
gn10e33d3w0w100	0(0)	0(2)	0(3)	0(3)	0(4)	0(5)	0(5)	0(6)	0(6)	31(7)	29(15)	32(33)	44(66)
gn11e13d3w0w100	309(0)	924(0)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(1)	1000(2)	1000(2)	1000(4)	1000(10)	1000(19)
gn11e20d3w0w100	33(1)	213(2)	197(1)	222(2)	208(2)	269(3)	305(3)	346(4)	357(4)	334(4)	364(9)	718(21)	701(34)
gn11e27d3w0w100	0(1)	8(1)	3(2)	7(2)	10(3)	7(4)	11(4)	8(5)	8(6)	14(6)	42(12)	41(31)	42(51)
gn11e41d3w0w100	0(0)	0(2)	0(3)	0(4)	0(5)	0(6)	0(7)	0(8)	0(9)	0(10)	0(21)	0(47)	0(108)
gn12e16d3w0w100	7(0)	554(1)	550(1)	569(2)	557(2)	578(2)	769(3)	1000(3)	1000(4)	1000(4)	1000(6)	1000(12)	1000(24)
gn12e24d3w0w100	0(1)	0(2)	0(3)	0(3)	0(3)	0(4)	0(4)	0(5)	0(6)	0(6)	0(12)	0(28)	0(48)
gn12e33d3w0w100	0(1)	0(4)	0(4)	0(5)	0(6)	0(6)	0(6)	0(6)	0(7)	0(8)	0(12)	0(27)	0(52)
gn13e19d3w0w100	2(1)	18(2)	31(2)	26(2)	28(3)	39(4)	31(4)	32(4)	36(5)	27(5)	182(11)	677(27)	737(43)
gn13e29d3w0w100	0(0)	0(2)	0(2)	0(3)	1(4)	1(5)	3(5)	2(6)	0(7)	0(8)	2(14)	2(30)	0(56)
gn13e39d3w0w100	0(0)	0(5)	0(5)	6(7)	5(8)	4(10)	4(11)	8(11)	16(13)	24(14)	79(24)	103(51)	235(110)
gn9e18d3w0w1000	0(0)	234(1)	235(1)	249(1)	589(2)	597(2)	751(2)	773(3)	773(3)	791(3)	1000(5)	1000(11)	1000(22)
gn9e27d3w0w1000	0(0)	0(1)	0(2)	0(2)	0(2)	0(3)	0(4)	0(4)	26(5)	24(6)	35(10)	44(28)	41(62)
gn10e22d3w0w1000	0(0)	22(1)	55(2)	50(2)	42(2)	39(3)	36(3)	53(4)	29(4)	33(5)	122(9)	122(21)	125(36)
gn10e28d3w0w1000	0(0)	0(2)	0(1)	0(2)	0(3)	0(4)	0(4)	0(5)	0(6)	0(6)	0(12)	0(30)	0(62)
gn10e33d3w0w1000	0(0)	1000(0)	1000(1)	1000(1)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(4)	1000(10)	1000(19)
gn11e13d3w0w1000	0(0)	20(1)	26(2)	53(2)	40(3)	58(3)	42(3)	38(3)	69(4)	63(4)	209(8)	628(22)	622(36)
gn11e20d3w0w1000	0(0)	0(2)	0(2)	0(3)	12(4)	6(5)	7(5)	9(6)	12(6)	8(7)	6(13)	8(37)	12(60)
gn11e27d3w0w1000	0(0)	0(2)	0(3)	0(4)	0(5)	0(6)	0(7)	0(8)	0(9)	0(9)	0(22)	0(53)	0(103)
gn11e41d3w0w1000	0(0)	0(2)	0(3)	0(4)	0(5)	0(6)	0(7)	0(8)	0(9)	1000(3)	1000(6)	1000(13)	1000(24)
gn12e16d3w0w1000	73(1)	487(1)	513(1)	874(2)	864(2)	860(2)	1000(3)	1000(3)	1000(3)	42(7)	123(15)	277(33)	365(60)
gn12e24d3w0w1000	0(0)	13(1)	19(2)	9(3)	10(4)	15(4)	32(5)	33(6)	34(6)	0(7)	0(17)	5(43)	20(84)
gn12e33d3w0w1000	0(1)	0(1)	0(2)	0(3)	0(4)	0(5)	0(6)	0(7)	0(8)	0(7)	0(17)	5(43)	20(84)
gn13e19d3w0w1000	2(0)	94(1)	163(2)	226(2)	288(3)	295(3)	460(4)	469(4)	477(5)	583(5)	816(9)	1000(21)	1000(38)
gn13e29d3w0w1000	0(1)	3(2)	0(3)	0(4)	4(4)	2(5)	0(6)	2(7)	1(8)	1(8)	2(16)	0(35)	2(67)
gn13e39d3w0w1000	0(0)	0(2)	0(3)	0(4)	0(5)	0(6)	0(7)	0(9)	0(10)	0(11)	0(21)	0(56)	0(108)

TABLE 4.20. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 5$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d5w0w1	0(0)	152(1)	223(1)	291(1)	316(2)	321(2)	327(2)	346(2)	319(2)	348(3)	325(4)	337(10)	321(21)
gn9e27d5w0w1	0(0)	323(1)	662(2)	878(3)	905(3)	974(3)	970(4)	966(4)	969(4)	980(5)	971(8)	968(19)	974(35)
gn10e22d5w0w1	0(1)	211(1)	316(1)	562(2)	668(3)	674(4)	739(4)	729(4)	852(5)	935(5)	1000(8)	1000(17)	1000(32)
gn10e28d5w0w1	0(0)	126(2)	141(2)	249(3)	309(3)	354(4)	367(4)	420(5)	449(5)	458(6)	756(11)	1000(26)	1000(47)
gn10e33d5w0w1	0(0)	98(2)	226(3)	313(3)	401(4)	458(5)	470(5)	605(7)	591(7)	647(7)	938(13)	998(26)	995(49)
gn11e13d5w0w1	231(0)	619(1)	838(1)	905(1)	896(1)	927(1)	1000(2)	1000(2)	1000(3)	1000(3)	1000(4)	1000(10)	1000(20)
gn11e20d5w0w1	97(1)	235(1)	341(1)	463(3)	651(2)	680(3)	675(3)	811(4)	815(4)	930(5)	984(8)	991(15)	989(29)
gn11e27d5w0w1	0(1)	175(2)	278(2)	388(3)	450(3)	522(4)	512(5)	572(5)	587(5)	618(6)	814(11)	845(23)	813(45)
gn11e41d5w0w1	0(1)	189(4)	349(5)	411(6)	528(7)	570(8)	571(8)	604(10)	644(10)	705(11)	873(20)	1000(45)	1000(79)
gn12e16d5w0w1	41(0)	779(1)	986(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(3)	1000(3)	1000(3)	1000(5)	1000(12)	1000(25)
gn12e24d5w0w1	24(0)	540(2)	747(3)	840(3)	902(4)	933(5)	966(5)	978(5)	971(6)	973(5)	988(10)	989(23)	984(44)
gn12e33d5w0w1	0(0)	354(4)	566(4)	644(5)	645(5)	680(6)	747(7)	773(8)	787(8)	803(9)	830(15)	808(32)	815(62)
gn13e19d5w0w1	3(0)	677(2)	681(3)	800(3)	926(3)	1000(4)	1000(4)	1000(5)	1000(5)	1000(8)	1000(8)	1000(18)	1000(36)
gn13e29d5w0w1	0(1)	360(3)	495(3)	635(5)	650(4)	724(5)	741(6)	765(7)	754(7)	765(7)	886(14)	881(29)	895(53)
gn13e39d5w0w1	0(1)	227(4)	359(5)	378(6)	461(6)	482(8)	530(9)	585(9)	604(10)	595(11)	676(19)	792(43)	831(84)
gn9e18d5w0w2	0(0)	44(1)	55(1)	36(1)	40(1)	35(2)	96(2)	122(2)	107(3)	101(3)	162(5)	139(12)	164(23)
gn9e27d5w0w2	0(1)	171(1)	486(2)	599(3)	679(3)	686(4)	716(4)	728(4)	698(5)	723(5)	808(9)	795(19)	789(36)
gn10e22d5w0w2	1(0)	309(2)	394(2)	453(2)	440(3)	440(3)	627(4)	671(4)	680(5)	681(5)	923(9)	1000(20)	1000(35)
gn10e28d5w0w2	0(1)	122(2)	187(3)	358(3)	356(4)	422(4)	472(5)	542(6)	601(7)	688(11)	725(23)	735(43)	735(43)
gn10e33d5w0w2	0(1)	65(3)	254(3)	316(4)	399(4)	349(5)	381(5)	371(6)	379(6)	408(7)	419(27)	414(51)	414(51)
gn11e13d5w0w2	265(0)	587(0)	706(1)	1000(1)	1000(2)	1000(1)	1000(2)	1000(2)	1000(2)	1000(2)	1000(4)	1000(11)	1000(20)
gn11e20d5w0w2	6(0)	109(1)	223(1)	371(3)	331(2)	388(3)	390(4)	420(4)	661(5)	641(5)	638(8)	627(17)	642(34)
gn11e27d5w0w2	5(0)	233(2)	411(3)	497(3)	483(4)	586(4)	631(4)	671(5)	639(6)	641(6)	678(10)	698(22)	698(43)
gn11e41d5w0w2	0(0)	396(3)	468(4)	555(5)	555(5)	595(6)	614(7)	679(8)	662(9)	688(9)	828(18)	895(40)	890(76)
gn12e16d5w0w2	139(0)	663(1)	673(2)	1000(2)	1000(2)	1000(2)	1000(3)	1000(2)	1000(3)	1000(3)	1000(6)	1000(12)	1000(24)
gn12e24d5w0w2	8(1)	238(2)	325(2)	383(3)	457(4)	489(4)	535(4)	538(6)	578(6)	602(6)	686(11)	699(24)	728(45)
gn12e33d5w0w2	0(1)	55(3)	132(4)	288(5)	385(6)	412(6)	397(7)	386(7)	389(9)	397(9)	407(14)	460(32)	457(60)
gn13e19d5w0w2	13(1)	63(1)	104(1)	174(2)	165(3)	158(3)	166(3)	158(3)	161(4)	203(5)	441(9)	405(19)	426(36)
gn13e29d5w0w2	1(0)	152(3)	278(3)	372(4)	450(5)	487(6)	580(7)	578(7)	627(9)	673(9)	937(17)	994(35)	997(62)
gn13e39d5w0w2	2(0)	355(3)	440(4)	578(5)	627(6)	674(6)	748(8)	803(8)	801(10)	802(10)	966(19)	975(41)	979(79)
gn9e18d5w0w5	0(1)	231(1)	228(2)	221(2)	253(2)	221(2)	231(3)	229(3)	256(3)	250(3)	453(7)	1000(18)	1000(28)
gn9e27d5w0w5	0(0)	0(2)	0(2)	0(2)	0(3)	0(4)	0(4)	0(5)	0(6)	0(6)	0(11)	0(25)	0(47)
gn10e22d5w0w5	0(1)	104(2)	194(2)	203(2)	212(3)	226(3)	335(4)	330(4)	346(4)	380(5)	563(9)	741(20)	757(36)
gn10e28d5w0w5	0(0)	19(1)	19(2)	45(2)	30(4)	38(4)	40(5)	43(5)	51(6)	55(6)	89(12)	299(29)	415(56)
gn10e33d5w0w5	0(0)	17(2)	15(3)	29(3)	21(4)	14(5)	16(6)	14(6)	25(6)	39(8)	83(15)	416(36)	919(79)
gn11e13d5w0w5	231(0)	1000(1)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(2)	1000(4)	1000(9)	1000(19)	1000(19)
gn11e20d5w0w5	8(1)	24(1)	21(2)	115(2)	129(3)	261(4)	260(4)	285(4)	309(5)	274(5)	720(10)	1000(22)	1000(38)
gn11e27d5w0w5	0(0)	0(2)	5(3)	1(3)	2(4)	1(4)	5(5)	3(6)	5(7)	15(8)	21(14)	72(31)	88(56)
gn11e41d5w0w5	0(1)	3(4)	39(5)	55(6)	127(7)	137(8)	136(9)	156(11)	181(12)	225(14)	229(23)	325(51)	398(96)
gn12e16d5w0w5	48(0)	622(1)	590(1)	1000(2)	1000(2)	1000(2)	1000(3)	1000(3)	1000(3)	1000(6)	1000(13)	1000(25)	1000(25)
gn12e24d5w0w5	5(1)	51(2)	65(3)	93(3)	89(5)	93(5)	109(5)	119(6)	137(7)	155(8)	321(16)	717(38)	988(68)
gn12e33d5w0w5	0(1)	8(3)	39(5)	40(5)	59(6)	91(7)	92(8)	112(10)	119(9)	122(11)	364(19)	555(42)	959(88)
gn13e19d5w0w5	0(0)	72(2)	83(2)	84(2)	122(3)	173(3)	162(4)	188(4)	237(4)	293(5)	683(12)	941(26)	1000(45)
gn13e29d5w0w5	0(1)	46(3)	83(4)	90(5)	88(6)	127(6)	118(7)	140(7)	164(8)	167(9)	293(19)	751(47)	932(84)
gn13e39d5w0w5	0(0)	0(4)	1(6)	2(7)	14(8)	14(10)	15(10)	31(11)	18(12)	21(13)	65(23)	181(54)	318(106)

TABLE 4.21. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 2$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d2w0w20	0(1)	95(1)	271(1)	339(1)	404(2)	400(2)	517(3)	516(2)	588(3)	592(3)	612(5)	910(13)	1000(25)
gn9e27d2w0w20	7(0)	159(1)	297(1)	343(2)	485(3)	519(3)	551(3)	554(3)	563(4)	660(4)	839(8)	935(18)	1000(35)
gn10e22d2w0w20	44(0)	352(1)	395(2)	458(2)	521(2)	667(3)	658(3)	716(3)	719(3)	695(4)	817(7)	887(16)	889(29)
gn10e28d2w0w20	8(1)	269(2)	355(2)	357(2)	417(3)	496(4)	570(4)	560(4)	570(5)	628(5)	772(10)	975(22)	1000(41)
gn10e33d2w0w20	15(0)	164(1)	270(2)	333(3)	362(3)	432(4)	438(4)	534(5)	683(6)	713(6)	847(10)	977(24)	985(44)
gn11e13d2w0w20	294(0)	428(1)	442(1)	619(1)	602(2)	607(2)	620(1)	772(2)	892(3)	1000(3)	1000(5)	1000(10)	1000(19)
gn11e20d2w0w20	0(0)	78(1)	104(1)	248(2)	287(3)	287(3)	298(3)	284(3)	294(4)	269(4)	573(7)	659(16)	648(28)
gn11e27d2w0w20	11(1)	257(1)	320(2)	339(2)	359(3)	417(3)	505(3)	519(5)	628(4)	627(8)	794(10)	945(22)	995(42)
gn11e41d2w0w20	0(1)	208(4)	272(4)	326(5)	402(6)	425(6)	414(7)	466(8)	549(9)	562(9)	776(17)	964(37)	1000(65)
gn12e16d2w0w20	142(1)	556(1)	568(2)	567(1)	689(2)	782(2)	842(2)	860(3)	901(3)	900(3)	924(6)	916(12)	906(23)
gn12e24d2w0w20	11(0)	123(1)	153(2)	207(3)	226(3)	275(3)	268(4)	306(4)	331(6)	358(5)	415(10)	555(22)	580(41)
gn12e33d2w0w20	0(0)	112(3)	147(4)	156(4)	150(5)	130(6)	153(6)	165(7)	172(7)	157(9)	307(15)	493(31)	641(56)
gn13e19d2w0w20	12(0)	40(1)	89(1)	141(2)	203(2)	199(3)	199(3)	200(3)	216(4)	247(4)	448(9)	448(20)	435(35)
gn13e29d2w0w20	0(1)	138(4)	298(4)	301(5)	289(5)	332(6)	322(6)	333(7)	325(7)	329(8)	317(12)	731(36)	1000(69)
gn13e39d2w0w20	4(1)	240(3)	343(4)	400(4)	451(5)	513(6)	567(7)	599(8)	616(8)	682(9)	853(17)	955(39)	1000(71)
gn9e18d2w0w100	0(0)	75(1)	87(1)	175(2)	153(2)	457(2)	501(2)	824(2)	841(2)	987(4)	1000(5)	1000(11)	1000(21)
gn9e27d2w0w100	0(1)	95(2)	92(2)	97(3)	80(3)	69(4)	69(3)	73(5)	105(5)	80(5)	80(9)	80(18)	109(33)
gn10e22d2w0w100	0(0)	8(0)	4(1)	4(2)	5(2)	8(2)	10(3)	6(3)	9(4)	9(5)	3(8)	251(23)	285(45)
gn10e28d2w0w100	0(0)	0(2)	1(2)	0(3)	4(3)	7(4)	7(4)	5(5)	5(6)	8(6)	11(11)	29(31)	182(58)
gn10e33d2w0w100	0(1)	28(2)	43(3)	59(3)	68(4)	61(4)	72(5)	66(5)	84(6)	70(6)	77(11)	409(27)	537(49)
gn11e13d2w0w100	320(0)	1000(1)	1000(1)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(2)	1000(4)	1000(9)	1000(18)
gn11e20d2w0w100	3(1)	16(0)	39(2)	33(2)	157(2)	155(2)	171(2)	189(3)	177(4)	277(5)	283(8)	742(21)	1000(42)
gn11e27d2w0w100	0(0)	23(2)	26(2)	27(3)	28(3)	21(4)	29(4)	28(5)	36(5)	28(6)	79(12)	280(33)	323(58)
gn11e41d2w0w100	0(0)	0(3)	3(4)	1(4)	2(6)	7(6)	20(8)	25(8)	13(9)	16(10)	55(18)	63(42)	84(79)
gn12e16d2w0w100	77(0)	236(1)	332(1)	328(1)	354(1)	454(2)	882(3)	907(3)	902(3)	963(3)	1000(6)	1000(13)	1000(24)
gn12e24d2w0w100	0(0)	0(2)	2(3)	0(2)	3(3)	2(5)	3(5)	3(6)	9(6)	9(6)	69(12)	206(32)	484(63)
gn12e33d2w0w100	0(0)	16(2)	16(3)	21(4)	30(5)	35(5)	58(6)	61(6)	74(8)	76(9)	119(16)	362(39)	545(75)
gn13e19d2w0w100	3(0)	181(2)	176(1)	191(2)	184(2)	169(3)	196(3)	183(4)	195(4)	185(4)	304(8)	324(16)	310(32)
gn13e29d2w0w100	0(0)	10(3)	20(3)	15(4)	28(5)	39(6)	28(6)	32(7)	39(7)	34(8)	43(12)	41(28)	33(51)
gn13e39d2w0w100	0(1)	6(3)	11(4)	17(5)	34(7)	27(8)	40(8)	48(9)	55(10)	51(11)	121(20)	233(49)	396(94)
gn9e18d2w0w1000	36(0)	59(1)	57(2)	56(1)	131(1)	127(2)	113(2)	137(2)	128(3)	128(4)	137(6)	189(13)	229(22)
gn9e27d2w0w1000	0(0)	0(1)	0(2)	0(2)	0(3)	0(3)	0(3)	0(5)	0(5)	0(5)	0(5)	223(28)	296(52)
gn10e22d2w0w1000	0(1)	1(1)	5(1)	6(2)	24(2)	35(2)	34(4)	32(4)	46(4)	50(5)	57(10)	79(23)	101(49)
gn10e28d2w0w1000	0(0)	0(1)	0(2)	0(3)	0(3)	0(3)	0(4)	0(5)	0(6)	0(7)	0(12)	0(29)	0(58)
gn10e33d2w0w1000	0(1)	0(1)	0(2)	0(2)	0(3)	0(4)	0(4)	0(5)	0(5)	0(6)	0(14)	2(35)	7(74)
gn11e13d2w0w1000	194(0)	1000(0)	1000(1)	1000(1)	1000(1)	1000(2)	1000(1)	1000(1)	1000(1)	1000(2)	1000(4)	1000(9)	1000(18)
gn11e20d2w0w1000	0(1)	25(1)	54(1)	92(2)	88(3)	82(3)	90(3)	114(4)	86(5)	99(5)	305(9)	688(21)	775(36)
gn11e27d2w0w1000	0(1)	0(1)	0(2)	0(2)	9(3)	12(3)	6(4)	7(4)	9(5)	9(6)	33(11)	53(29)	69(62)
gn11e41d2w0w1000	0(1)	0(1)	0(3)	0(4)	0(4)	0(5)	0(6)	0(7)	0(8)	0(9)	0(17)	15(46)	20(92)
gn12e16d2w0w1000	102(0)	462(1)	537(1)	527(1)	653(2)	901(3)	893(2)	900(3)	1000(3)	1000(4)	1000(6)	1000(12)	1000(24)
gn12e24d2w0w1000	0(1)	0(1)	3(2)	5(2)	8(3)	7(4)	6(4)	10(5)	13(6)	7(6)	21(13)	72(30)	153(55)
gn12e33d2w0w1000	0(1)	0(2)	0(2)	0(3)	0(4)	0(5)	0(5)	0(7)	0(7)	0(7)	0(17)	0(47)	0(83)
gn13e19d2w0w1000	17(1)	20(1)	22(2)	16(2)	22(3)	22(3)	27(4)	32(4)	25(4)	27(5)	20(11)	291(28)	512(53)
gn13e29d2w0w1000	0(0)	7(2)	13(3)	19(4)	25(5)	20(5)	40(7)	29(7)	40(8)	50(9)	63(17)	162(38)	336(74)
gn13e39d2w0w1000	0(0)	0(3)	0(3)	1(5)	0(6)	0(6)	0(8)	2(8)	2(10)	1(10)	1(20)	51(53)	57(108)

TABLE 4.22. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 3$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d3w0w1000	13(0)	252(1)	253(1)	305(2)	320(1)	357(2)	338(2)	656(3)	662(3)	660(3)	856(6)	1000(14)	1000(25)
gn9e27d3w0w1000	0(0)	0(1)	4(2)	4(3)	8(3)	11(4)	19(3)	13(4)	15(5)	11(6)	57(11)	467(28)	645(49)
gn10e22d3w0w1000	0(0)	0(1)	1(1)	2(2)	2(3)	13(3)	12(4)	12(4)	5(4)	24(5)	74(10)	578(29)	930(52)
gn10e28d3w0w1000	0(0)	0(2)	0(3)	26(3)	21(4)	20(4)	16(5)	18(5)	12(6)	49(13)	24(7)	67(28)	59(52)
gn10e33d3w0w1000	0(0)	0(2)	0(3)	0(4)	10(5)	18(5)	18(6)	31(6)	23(7)	57(9)	91(14)	136(30)	148(52)
gn11e13d3w0w1000	255(0)	346(1)	335(0)	545(1)	546(1)	514(2)	550(2)	544(2)	519(2)	546(2)	572(4)	559(10)	573(19)
gn11e20d3w0w1000	0(1)	0(1)	0(2)	0(2)	0(2)	0(3)	0(3)	0(3)	0(4)	0(5)	0(8)	222(29)	439(48)
gn11e27d3w0w1000	0(0)	30(2)	33(3)	42(3)	33(4)	44(4)	54(5)	67(6)	69(6)	55(6)	159(12)	229(29)	369(56)
gn11e41d3w0w1000	0(1)	0(3)	0(5)	5(5)	6(6)	7(7)	13(8)	15(10)	12(10)	12(13)	14(21)	21(48)	23(81)
gn12e16d3w0w1000	50(0)	677(1)	700(2)	1000(2)	1000(2)	1000(3)	1000(2)	1000(2)	1000(3)	1000(3)	1000(5)	1000(13)	1000(25)
gn12e24d3w0w1000	0(0)	10(2)	6(3)	14(3)	14(4)	13(4)	18(5)	14(5)	21(6)	11(7)	15(12)	31(26)	26(49)
gn12e33d3w0w1000	5(0)	10(2)	18(2)	7(3)	8(5)	22(5)	22(6)	40(6)	42(7)	33(8)	51(16)	131(39)	146(69)
gn13e19d3w0w1000	0(1)	43(2)	49(2)	47(2)	179(3)	185(3)	216(4)	288(5)	510(7)	567(8)	917(13)	1000(23)	1000(40)
gn13e29d3w0w1000	0(1)	21(3)	44(3)	40(5)	29(6)	37(6)	45(6)	52(7)	55(8)	60(9)	149(17)	251(38)	377(78)
gn13e39d3w0w1000	0(1)	16(3)	20(4)	14(5)	32(6)	41(7)	50(8)	76(9)	73(9)	95(11)	144(21)	299(48)	452(91)
gn9e18d3w0w100	0(0)	44(1)	37(0)	35(1)	41(2)	30(2)	34(2)	39(2)	36(2)	38(4)	42(6)	83(15)	82(25)
gn9e27d3w0w100	0(0)	19(2)	18(2)	23(2)	37(3)	32(4)	36(4)	32(5)	70(5)	74(6)	78(13)	414(30)	710(53)
gn10e22d3w0w100	1(1)	55(1)	67(2)	58(3)	55(3)	70(4)	74(3)	70(5)	182(5)	212(6)	289(9)	806(26)	1000(44)
gn10e28d3w0w100	0(1)	0(3)	0(3)	0(3)	0(3)	0(4)	0(5)	0(5)	0(6)	0(6)	0(14)	0(29)	0(53)
gn10e33d3w0w100	0(1)	0(2)	0(3)	0(2)	0(4)	15(5)	23(6)	26(7)	24(7)	26(8)	20(16)	27(47)	20(88)
gn11e13d3w0w100	297(1)	923(0)	1000(0)	1000(1)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(3)	1000(4)	1000(10)	1000(20)
gn11e20d3w0w100	33(0)	129(1)	198(1)	209(2)	196(3)	205(3)	234(3)	254(4)	258(4)	253(4)	608(10)	712(20)	690(33)
gn11e27d3w0w100	0(1)	19(1)	20(2)	28(3)	22(4)	26(4)	25(5)	33(6)	20(7)	29(8)	62(14)	72(29)	56(52)
gn11e41d3w0w100	0(0)	0(3)	0(3)	0(5)	0(5)	0(7)	0(8)	0(8)	0(9)	0(10)	0(22)	2(60)	3(112)
gn12e16d3w0w100	8(1)	479(1)	476(1)	469(1)	454(2)	886(3)	1000(3)	1000(3)	1000(3)	1000(4)	1000(7)	1000(12)	1000(25)
gn12e24d3w0w100	0(1)	0(1)	5(2)	6(3)	8(3)	4(5)	3(5)	8(6)	6(7)	5(7)	6(13)	7(29)	6(56)
gn12e33d3w0w100	0(0)	0(4)	0(5)	0(5)	0(5)	0(6)	0(7)	0(7)	0(8)	0(8)	0(14)	0(29)	0(57)
gn13e19d3w0w100	2(0)	13(2)	9(2)	13(2)	13(3)	13(3)	17(5)	9(4)	10(5)	14(6)	46(13)	955(30)	1000(49)
gn13e29d3w0w100	0(0)	0(2)	0(3)	3(4)	3(4)	1(5)	3(6)	4(6)	15(7)	3(8)	14(14)	13(30)	17(56)
gn13e39d3w0w100	0(1)	0(5)	0(5)	0(7)	1(8)	4(9)	1(11)	2(12)	6(13)	5(15)	53(27)	40(67)	63(118)
gn9e18d3w0w1000	0(0)	148(1)	138(1)	152(2)	495(2)	483(3)	1000(3)	1000(4)	1000(4)	1000(4)	1000(6)	1000(12)	1000(23)
gn9e27d3w0w1000	0(1)	0(1)	0(2)	0(3)	0(3)	0(4)	0(5)	0(5)	0(6)	0(6)	0(11)	61(28)	107(56)
gn10e22d3w0w1000	0(0)	25(1)	43(2)	36(2)	59(3)	100(3)	90(3)	67(4)	85(4)	71(4)	142(9)	442(23)	645(44)
gn10e28d3w0w1000	0(0)	0(1)	0(2)	0(3)	0(3)	0(4)	0(4)	0(5)	0(5)	0(6)	0(16)	0(31)	0(70)
gn10e33d3w0w1000	0(0)	0(1)	0(2)	0(3)	0(4)	0(4)	0(5)	0(5)	0(6)	0(8)	0(16)	5(40)	1(81)
gn11e13d3w0w1000	305(0)	1000(1)	1000(1)	1000(1)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(4)	1000(10)	1000(20)
gn11e20d3w0w1000	0(0)	21(1)	25(2)	40(2)	46(2)	35(3)	40(3)	32(4)	36(4)	34(5)	212(9)	271(21)	447(39)
gn11e27d3w0w1000	0(0)	2(1)	3(3)	2(3)	3(3)	3(4)	5(5)	4(5)	1(6)	5(13)	2(15)	2(39)	1(70)
gn11e41d3w0w1000	0(1)	0(2)	0(3)	0(4)	0(5)	0(6)	0(7)	0(8)	0(9)	0(10)	0(21)	0(55)	0(111)
gn12e16d3w0w1000	82(0)	511(1)	489(1)	840(2)	845(2)	856(2)	857(3)	861(3)	857(3)	847(3)	1000(6)	1000(13)	1000(24)
gn12e24d3w0w1000	0(0)	23(1)	23(3)	18(3)	31(3)	24(5)	30(5)	22(6)	23(7)	25(7)	23(18)	27(34)	18(56)
gn12e33d3w0w1000	0(1)	0(2)	0(3)	0(4)	0(5)	0(5)	0(6)	0(7)	0(8)	0(9)	0(18)	0(51)	0(97)
gn13e19d3w0w1000	0(1)	140(1)	242(2)	300(2)	370(3)	433(4)	538(4)	571(4)	562(5)	578(6)	785(10)	1000(22)	1000(39)
gn13e29d3w0w1000	0(0)	1(2)	3(2)	5(3)	3(4)	6(5)	7(6)	4(7)	7(8)	7(9)	16(17)	21(40)	23(71)
gn13e39d3w0w1000	0(1)	0(3)	0(4)	0(4)	0(6)	0(8)	0(8)	0(9)	0(10)	0(12)	0(23)	0(57)	0(112)

TABLE 4.23. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior integer point that is the image of some base. Here $d = 5$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d5w0w1	0(0)	159(1)	245(1)	242(1)	303(1)	344(2)	346(2)	312(2)	320(3)	352(3)	319(5)	339(12)	349(22)
gn9e27d5w0w1	0(0)	332(2)	646(3)	865(3)	921(3)	978(3)	974(4)	960(4)	972(5)	970(6)	980(9)	971(20)	972(38)
gn10e22d5w0w1	0(1)	220(1)	313(2)	590(2)	645(3)	637(3)	713(4)	750(4)	876(5)	925(5)	1000(9)	1000(18)	1000(35)
gn10e28d5w0w1	0(1)	127(1)	155(2)	251(3)	303(4)	356(4)	402(4)	398(5)	472(6)	475(6)	722(12)	1000(28)	1000(49)
gn10e33d5w0w1	0(0)	111(3)	230(3)	303(4)	397(5)	450(5)	463(5)	631(6)	659(7)	703(8)	922(14)	997(29)	995(53)
gn11e13d5w0w1	248(0)	588(1)	845(1)	885(1)	891(2)	923(2)	1000(2)	1000(2)	1000(2)	1000(2)	1000(4)	1000(11)	1000(21)
gn11e20d5w0w1	96(0)	218(1)	354(2)	421(2)	641(3)	665(3)	678(3)	819(4)	819(5)	928(5)	982(8)	991(17)	990(31)
gn11e27d5w0w1	0(1)	162(2)	299(3)	413(3)	454(3)	487(4)	529(4)	579(5)	576(5)	644(6)	802(12)	801(26)	823(48)
gn11e41d5w0w1	0(1)	204(4)	401(5)	420(7)	529(7)	565(8)	573(8)	600(10)	642(11)	712(12)	863(22)	1000(48)	1000(84)
gn12e16d5w0w1	50(1)	762(1)	991(1)	1000(2)	1000(2)	1000(2)	1000(3)	1000(2)	1000(3)	1000(3)	1000(5)	1000(14)	1000(26)
gn12e24d5w0w1	25(1)	522(2)	720(3)	872(3)	909(4)	931(4)	954(5)	964(6)	974(6)	971(7)	985(10)	991(24)	990(46)
gn12e33d5w0w1	1(1)	402(4)	553(4)	634(5)	670(6)	704(7)	741(8)	759(8)	773(9)	791(10)	812(16)	806(35)	809(65)
gn13e19d5w0w1	3(0)	666(2)	665(2)	801(3)	914(3)	1000(4)	1000(5)	1000(4)	1000(5)	1000(5)	1000(10)	1000(20)	1000(38)
gn13e29d5w0w1	1(0)	377(3)	521(3)	614(5)	626(5)	761(6)	752(7)	773(7)	763(8)	794(9)	874(15)	885(31)	891(57)
gn13e39d5w0w1	0(0)	228(4)	370(5)	407(6)	458(7)	488(8)	510(9)	574(10)	573(11)	569(12)	698(21)	797(50)	843(88)
gn9e18d5w0w2	0(1)	107(1)	104(2)	80(2)	110(2)	106(2)	135(3)	150(3)	147(3)	151(4)	266(6)	253(14)	265(25)
gn9e27d5w0w2	0(0)	150(2)	438(2)	482(3)	621(4)	648(4)	628(4)	659(5)	643(5)	673(6)	750(9)	719(21)	738(39)
gn10e22d5w0w2	0(0)	264(1)	385(2)	430(2)	432(3)	430(4)	550(4)	582(4)	589(5)	576(5)	880(10)	1000(22)	1000(38)
gn10e28d5w0w2	0(0)	115(2)	220(3)	340(3)	382(4)	457(5)	482(6)	491(5)	561(8)	603(7)	677(12)	746(26)	733(48)
gn10e33d5w0w2	0(1)	64(2)	270(3)	315(5)	391(5)	383(5)	402(6)	385(6)	382(13)	368(7)	382(13)	411(29)	416(55)
gn11e13d5w0w2	278(1)	610(1)	685(0)	1000(2)	1000(2)	1000(1)	1000(2)	1000(2)	1000(3)	1000(3)	1000(5)	1000(11)	1000(21)
gn11e20d5w0w2	7(1)	146(2)	220(2)	356(3)	352(2)	409(4)	397(3)	400(4)	626(5)	639(5)	643(9)	603(19)	630(36)
gn11e27d5w0w2	6(0)	239(2)	368(2)	450(4)	459(3)	573(4)	615(4)	629(6)	627(7)	647(6)	690(11)	707(24)	705(46)
gn11e41d5w0w2	0(1)	382(4)	450(4)	492(5)	564(6)	596(7)	595(7)	620(9)	664(9)	686(11)	840(19)	885(43)	899(80)
gn12e16d5w0w2	163(0)	666(1)	671(2)	1000(2)	1000(2)	1000(2)	1000(2)	1000(3)	1000(3)	1000(3)	1000(6)	1000(14)	1000(26)
gn12e24d5w0w2	8(0)	223(2)	340(3)	401(3)	493(3)	457(5)	526(5)	550(5)	588(6)	622(6)	686(11)	701(26)	716(50)
gn12e33d5w0w2	0(0)	54(3)	146(5)	282(6)	361(6)	378(7)	350(7)	381(8)	389(9)	413(10)	443(16)	455(35)	441(64)
gn13e19d5w0w2	15(0)	67(1)	115(2)	148(3)	166(3)	145(4)	171(3)	157(4)	164(5)	211(5)	432(9)	429(20)	434(37)
gn13e29d5w0w2	0(1)	190(3)	288(4)	352(5)	457(6)	523(7)	549(7)	581(8)	615(9)	649(10)	923(18)	996(38)	998(66)
gn13e39d5w0w2	4(0)	407(3)	446(4)	575(6)	634(7)	680(7)	727(9)	777(9)	814(10)	805(11)	957(21)	975(45)	975(85)
gn9e18d5w0w5	0(0)	74(1)	125(1)	128(2)	131(2)	115(3)	129(3)	148(3)	209(3)	181(4)	316(7)	1000(19)	1000(31)
gn9e27d5w0w5	0(0)	0(1)	0(2)	0(3)	0(3)	0(4)	0(5)	0(5)	0(6)	0(6)	0(13)	0(25)	0(52)
gn10e22d5w0w5	0(1)	62(2)	173(2)	238(2)	231(3)	261(3)	387(4)	395(4)	442(5)	425(5)	682(10)	964(22)	974(38)
gn10e28d5w0w5	0(0)	24(1)	47(2)	32(3)	32(4)	38(4)	32(5)	35(6)	39(7)	50(6)	112(13)	226(31)	360(60)
gn10e33d5w0w5	0(0)	9(2)	16(3)	20(3)	21(4)	12(5)	20(5)	40(7)	37(8)	35(8)	162(17)	389(39)	833(82)
gn11e13d5w0w5	258(0)	1000(0)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(2)	1000(3)	1000(5)	1000(10)	1000(21)
gn11e20d5w0w5	11(0)	29(1)	22(2)	150(3)	137(2)	288(3)	293(4)	321(4)	325(5)	308(6)	626(11)	1000(24)	1000(42)
gn11e27d5w0w5	0(0)	0(2)	1(3)	3(3)	0(5)	4(5)	5(7)	1(6)	29(7)	14(8)	20(16)	20(34)	19(58)
gn11e41d5w0w5	0(1)	11(4)	61(6)	65(7)	97(8)	105(9)	173(11)	162(12)	189(14)	248(15)	317(25)	526(56)	780(113)
gn12e16d5w0w5	43(0)	782(1)	788(1)	1000(2)	1000(2)	1000(3)	1000(3)	1000(3)	1000(3)	1000(4)	1000(6)	1000(14)	1000(26)
gn12e24d5w0w5	6(0)	51(2)	81(3)	87(4)	99(5)	105(6)	128(6)	133(7)	221(8)	234(8)	368(16)	577(36)	955(74)
gn12e33d5w0w5	0(1)	4(3)	7(5)	9(5)	41(7)	48(8)	54(9)	61(10)	84(12)	84(12)	228(22)	660(50)	953(93)
gn13e19d5w0w5	0(1)	85(2)	87(2)	78(2)	146(3)	185(3)	194(4)	177(5)	204(5)	213(5)	399(10)	982(27)	981(46)
gn13e29d5w0w5	0(1)	55(3)	84(4)	106(5)	119(6)	151(7)	153(7)	170(8)	157(9)	183(10)	344(21)	714(49)	945(92)
gn13e39d5w0w5	0(1)	0(5)	1(6)	5(8)	19(8)	18(10)	9(11)	24(12)	17(13)	26(14)	99(25)	211(59)	543(125)

TABLE 4.24. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior rational non-integer point. Here $d = 2$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d2w0w20	2,981	1,742	58.44	144	1
gn9e27d2w0w20	372,320	6,346	01.70	273	1
gn10e22d2w0w20	53,357	3,957	07.42	203	2
gn10e28d2w0w20	800,948	7,131	00.89	364	2
gn10e33d2w0w20	3,584,016	10,833	00.30	216	2
gn11e13d2w0w20	41	41	100.00	252	1
gn11e20d2w0w20	6,939	2,173	31.32	125	2
gn11e27d2w0w20	284,730	7,515	02.64	181	1
gn11e41d2w0w20	90,922,271	16,457	00.02	318	2
gn12e16d2w0w20	162	154	95.06	240	1
gn12e24d2w0w20	208,380	5,647	02.71	166	2
gn12e33d2w0w20	4,741,624	9,364	00.20	219	2
gn13e19d2w0w20	3,401	1,608	47.28	78	1
gn13e29d2w0w20	1,543,340	8,474	00.55	199	2
gn13e39d2w0w20	131,807,934	18,468	00.01	232	3
gn9e18d2w0w100	2,981	2,858	95.87	440	1
gn9e27d2w0w100	372,320	89,092	23.93	0	1
gn10e22d2w0w100	53,357	37,204	69.73	47	2
gn10e28d2w0w100	800,948	101,334	12.65	0	2
gn10e33d2w0w100	3,584,016	166,427	04.64	16	2
gn11e13d2w0w100	41	41	100.00	283	1
gn11e20d2w0w100	6,939	6,580	94.83	0	1
gn11e27d2w0w100	284,730	81,803	28.73	0	2
gn11e41d2w0w100	90,922,271	309,961	00.34	21	3
gn12e16d2w0w100	162	162	100.00	526	1
gn12e24d2w0w100	208,380	92,813	44.54	0	2
gn12e33d2w0w100	4,741,624	192,122	04.05	7	2
gn13e19d2w0w100	3,401	3,255	95.71	78	1
gn13e29d2w0w100	1,543,340	164,617	10.67	8	2
gn13e39d2w0w100	131,807,934	315,881	00.24	34	3
gn9e18d2w0w1000	2,981	2,981	100.00	75	1
gn9e27d2w0w1000	372,320	364,382	97.87	0	1
gn10e22d2w0w1000	53,357	52,990	99.31	0	1
gn10e28d2w0w1000	800,948	756,013	94.39	0	2
gn10e33d2w0w1000	3,584,016	2,726,287	76.07	0	2
gn11e13d2w0w1000	41	41	100.00	766	0
gn11e20d2w0w1000	6,939	6,921	99.74	12	1
gn11e27d2w0w1000	284,730	279,308	98.10	0	1
gn11e41d2w0w1000	90,922,271	13,884,793	15.27	0	2
gn12e16d2w0w1000	162	162	100.00	355	1
gn12e24d2w0w1000	208,380	205,690	98.71	2	2
gn12e33d2w0w1000	4,741,624	3,680,313	77.62	0	3
gn13e19d2w0w1000	3,401	3,401	100.00	1	1
gn13e29d2w0w1000	1,543,340	1,396,180	90.46	1	2
gn13e39d2w0w1000	131,807,934	15,037,589	11.41	0	3

TABLE 4.25. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior rational non-integer point. Here $d = 3$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d3w0w20	2,981	2,905	97.45	62	1
gn9e27d3w0w20	372,320	134,039	36.00	0	1
gn10e22d3w0w20	53,357	42,887	80.38	18	2
gn10e28d3w0w20	800,948	238,529	29.78	0	2
gn10e33d3w0w20	3,584,016	411,730	11.49	0	3
gn11e13d3w0w20	41	41	100.00	183	0
gn11e20d3w0w20	6,939	6,603	95.16	9	1
gn11e27d3w0w20	284,730	146,672	51.51	1	2
gn11e41d3w0w20	90,922,271	959,469	01.06	14	2
gn12e16d3w0w20	162	162	100.00	206	1
gn12e24d3w0w20	208,380	111,201	53.36	32	2
gn12e33d3w0w20	4,741,624	470,609	09.93	25	2
gn13e19d3w0w20	3,401	3,358	98.74	17	1
gn13e29d3w0w20	1,543,340	264,949	17.18	23	2
gn13e39d3w0w20	131,807,934	930,322	00.71	14	3
gn9e18d3w0w100	2,981	2,981	100.00	33	1
gn9e27d3w0w100	372,320	367,313	98.66	0	2
gn10e22d3w0w100	53,357	53,289	99.87	2	1
gn10e28d3w0w100	800,948	786,781	98.23	0	2
gn10e33d3w0w100	3,584,016	3,351,096	93.01	0	2
gn11e13d3w0w100	41	41	100.00	700	0
gn11e20d3w0w100	6,939	6,939	100.00	0	2
gn11e27d3w0w100	284,730	281,303	98.80	0	1
gn11e41d3w0w100	90,922,271	35,943,327	39.53	0	3
gn12e16d3w0w100	162	162	100.00	825	1
gn12e24d3w0w100	208,380	207,143	99.41	3	2
gn12e33d3w0w100	4,741,624	4,740,880	99.98	0	3
gn13e19d3w0w100	3,401	3,401	100.00	59	1
gn13e29d3w0w100	1,543,340	1,484,719	96.20	0	3
gn13e39d3w0w100	131,807,934	44,757,592	33.96	0	4
gn9e18d3w0w1000	2,981	2,981	100.00	235	1
gn9e27d3w0w1000	372,320	372,320	100.00	0	1
gn10e22d3w0w1000	53,357	53,357	100.00	10	1
gn10e28d3w0w1000	800,948	800,946	99.99	0	2
gn10e33d3w0w1000	3,584,016	3,583,757	99.99	0	2
gn11e13d3w0w1000	41	41	100.00	410	1
gn11e20d3w0w1000	6,939	6,939	100.00	0	1
gn11e27d3w0w1000	284,730	284,730	100.00	1	2
gn11e41d3w0w1000	90,922,271	90,699,181	99.75	0	3
gn12e16d3w0w1000	162	162	100.00	583	1
gn12e24d3w0w1000	208,380	208,356	99.99	0	2
gn12e33d3w0w1000	4,741,624	4,740,880	99.98	0	3
gn13e19d3w0w1000	3,401	3,401	100.00	51	1
gn13e29d3w0w1000	1,543,340	1,543,304	99.99	0	3
gn13e39d3w0w1000	131,807,934	131,464,478	99.74	0	3

TABLE 4.26. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior rational non-integer point. Here $d = 5$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d5w0w1	2,981	1,099	36.87	624	2
gn9e27d5w0w1	372,320	8,205	2.20	766	2
gn10e22d5w0w1	53,357	4,746	8.89	693	2
gn10e28d5w0w1	800,948	10,898	1.36	774	2
gn10e33d5w0w1	3,584,016	15,482	0.43	956	3
gn11e13d5w0w1	41	27	65.85	504	0
gn11e20d5w0w1	6,939	1,866	26.89	529	2
gn11e27d5w0w1	284,730	5,828	2.05	608	2
gn11e41d5w0w1	90,922,271	36,847	0.04	953	3
gn12e16d5w0w1	162	116	71.61	286	1
gn12e24d5w0w1	208,380	6,738	3.23	860	3
gn12e33d5w0w1	4,741,624	20,857	0.44	957	3
gn13e19d5w0w1	3,401	1,149	33.78	448	1
gn13e29d5w0w1	1,543,340	111,116	7.20	746	3
gn13e39d5w0w1	131,807,934	34,392	0.03	623	5
gn9e18d5w0w2	2,981	2,276	76.35	308	1
gn9e27d5w0w2	372,320	43,029	11.55	323	2
gn10e22d5w0w2	53,357	14,623	27.40	97	2
gn10e28d5w0w2	800,948	66,190	8.26	268	2
gn10e33d5w0w2	3,584,016	105,309	2.93	189	3
gn11e13d5w0w2	41	41	100.00	706	1
gn11e20d5w0w2	6,939	3,761	54.20	177	1
gn11e27d5w0w2	284,730	39,292	13.79	257	3
gn11e41d5w0w2	90,922,271	290,555	0.31	430	4
gn12e16d5w0w2	162	160	98.76	245	1
gn12e24d5w0w2	208,380	34,057	16.34	146	2
gn12e33d5w0w2	4,741,624	120,211	2.53	404	3
gn13e19d5w0w2	3,401	2,613	76.83	28	2
gn13e29d5w0w2	1,543,340	98,285	6.36	288	4
gn13e39d5w0w2	131,807,934	348,703	0.26	361	4
gn9e18d5w0w5	2,981	2,960	99.29	32	1
gn9e27d5w0w5	372,320	271,048	72.79	0	2
gn10e22d5w0w5	53,357	49,463	92.70	0	1
gn10e28d5w0w5	800,948	493,565	61.62	9	2
gn10e33d5w0w5	3,584,016	1,294,875	36.12	1	3
gn11e13d5w0w5	41	41	100.00	574	0
gn11e20d5w0w5	6,939	5,975	86.10	14	1
gn11e27d5w0w5	284,730	222,974	78.31	0	2
gn11e41d5w0w5	90,922,271	4,711,354	5.18	33	4
gn12e16d5w0w5	162	162	100.00	194	0
gn12e24d5w0w5	208,380	177,845	85.34	0	2
gn12e33d5w0w5	4,741,624	1,489,751	31.41	27	4
gn13e19d5w0w5	3,401	3,381	99.41	48	1
gn13e29d5w0w5	1,543,340	787,196	51.00	25	4
gn13e39d5w0w5	131,807,934	7,737,684	5.87	16	5

TABLE 4.27. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior rational non-integer point. Here $d = 2$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d2w0w20	2,981	1,742	58.44	161	1
gn9e27d2w0w20	372,320	6,346	01.70	211	1
gn10e22d2w0w20	53,357	3,957	07.42	194	1
gn10e28d2w0w20	800,948	7,131	00.89	363	1
gn10e33d2w0w20	3,584,016	10,833	00.30	254	2
gn11e13d2w0w20	41	41	100.00	210	1
gn11e20d2w0w20	6,939	2,173	31.32	124	1
gn11e27d2w0w20	284,730	7,515	02.64	234	1
gn11e41d2w0w20	90,922,271	16,457	00.02	390	2
gn12e16d2w0w20	162	154	95.06	260	1
gn12e24d2w0w20	208,380	5,647	02.71	166	1
gn12e33d2w0w20	4,741,624	9,364	00.20	224	2
gn13e19d2w0w20	3,401	1,608	47.28	82	2
gn13e29d2w0w20	1,543,340	8,474	00.55	172	1
gn13e39d2w0w20	131,807,934	18,468	00.01	240	3
gn9e18d2w0w100	2,981	2,858	95.87	373	1
gn9e27d2w0w100	372,320	89,092	23.93	5	1
gn10e22d2w0w100	53,357	37,204	69.73	53	1
gn10e28d2w0w100	800,948	101,334	12.65	27	2
gn10e33d2w0w100	3,584,016	166,427	04.64	12	1
gn11e13d2w0w100	41	41	100.00	330	0
gn11e20d2w0w100	6,939	6,580	94.83	0	1
gn11e27d2w0w100	284,730	81,803	28.73	0	1
gn11e41d2w0w100	90,922,271	309,961	00.34	18	3
gn12e16d2w0w100	162	162	100.00	537	1
gn12e24d2w0w100	208,380	92,813	44.54	0	1
gn12e33d2w0w100	4,741,624	192,122	04.05	5	2
gn13e19d2w0w100	3,401	3,255	95.71	89	1
gn13e29d2w0w100	1,543,340	164,617	10.67	18	1
gn13e39d2w0w100	131,807,934	315,881	00.24	34	3
gn9e18d2w0w1000	2,981	2,981	100.00	71	1
gn9e27d2w0w1000	372,320	364,382	97.87	0	2
gn10e22d2w0w1000	53,357	52,990	99.31	0	1
gn10e28d2w0w1000	800,948	756,013	94.39	0	2
gn10e33d2w0w1000	3,584,016	2,726,287	76.07	0	1
gn11e13d2w0w1000	41	41	100.00	434	0
gn11e20d2w0w1000	6,939	6,921	99.74	12	1
gn11e27d2w0w1000	284,730	279,308	98.10	0	2
gn11e41d2w0w1000	90,922,271	13,884,793	15.27	0	2
gn12e16d2w0w1000	162	162	100.00	346	1
gn12e24d2w0w1000	208,380	205,690	98.71	0	1
gn12e33d2w0w1000	4,741,624	3,680,313	77.62	0	2
gn13e19d2w0w1000	3,401	3,401	100.00	5	1
gn13e29d2w0w1000	1,543,340	1,396,180	90.46	0	1
gn13e39d2w0w1000	131,807,934	15,037,589	11.41	0	3

TABLE 4.28. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior rational non-integer point. Here $d = 3$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d3w0w20	2,981	2,905	97.45	0	1
gn9e27d3w0w20	372,320	134,039	36.00	8	2
gn10e22d3w0w20	53,357	42,887	80.38	18	1
gn10e28d3w0w20	800,948	238,529	29.78	0	2
gn10e33d3w0w20	3,584,016	411,730	11.49	1	2
gn11e13d3w0w20	41	41	100.00	185	0
gn11e20d3w0w20	6,939	6,603	95.16	8	1
gn11e27d3w0w20	284,730	146,672	51.51	3	2
gn11e41d3w0w20	90,922,271	959,469	01.06	19	3
gn12e16d3w0w20	162	162	100.00	116	1
gn12e24d3w0w20	208,380	111,201	53.36	20	2
gn12e33d3w0w20	4,741,624	470,609	09.93	19	2
gn13e19d3w0w20	3,401	3,358	98.74	18	1
gn13e29d3w0w20	1,543,340	264,949	17.18	30	3
gn13e39d3w0w20	131,807,934	930,322	00.71	9	3
gn9e18d3w0w100	2,981	2,981	100.00	36	1
gn9e27d3w0w100	372,320	367,313	98.66	2	2
gn10e22d3w0w100	53,357	53,289	99.87	3	1
gn10e28d3w0w100	800,948	786,781	98.23	0	2
gn10e33d3w0w100	3,584,016	3,351,096	93.01	0	2
gn11e13d3w0w100	41	41	100.00	699	1
gn11e20d3w0w100	6,939	6,939	100.00	0	1
gn11e27d3w0w100	284,730	281,303	98.80	0	2
gn11e41d3w0w100	90,922,271	35,943,327	39.53	0	3
gn12e16d3w0w100	162	162	100.00	668	1
gn12e24d3w0w100	208,380	207,143	99.41	0	2
gn12e33d3w0w100	4,741,624	4,740,880	99.98	0	3
gn13e19d3w0w100	3,401	3,401	100.00	10	1
gn13e29d3w0w100	1,543,340	1,484,719	96.20	0	2
gn13e39d3w0w100	131,807,934	44,757,592	33.96	0	3
gn9e18d3w0w1000	2,981	2,981	100.00	135	1
gn9e27d3w0w1000	372,320	372,320	100.00	0	2
gn10e22d3w0w1000	53,357	53,357	100.00	16	1
gn10e28d3w0w1000	800,948	800,946	99.99	0	1
gn10e33d3w0w1000	3,584,016	3,583,757	99.99	0	2
gn11e13d3w0w1000	41	41	100.00	432	0
gn11e20d3w0w1000	6,939	6,939	100.00	0	1
gn11e27d3w0w1000	284,730	284,730	100.00	3	2
gn11e41d3w0w1000	90,922,271	90,699,181	99.75	0	3
gn12e16d3w0w1000	162	162	100.00	576	1
gn12e24d3w0w1000	208,380	208,356	99.99	2	2
gn12e33d3w0w1000	4,741,624	4,740,880	99.98	0	2
gn13e19d3w0w1000	3,401	3,401	100.00	5	1
gn13e29d3w0w1000	1,543,340	1,543,304	99.99	0	2
gn13e39d3w0w1000	131,807,934	131,464,478	99.74	0	3

TABLE 4.29. Calibration test set: Local Search **Heuristic 82** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior rational non-integer point. Here $d = 5$ and all tests run on Fuzzy. Shown are the number of spanning trees and projected bases of each instance and the number of successes out of 1000 Local Searches and seconds to perform the 1000 searches. We also repeat the exact number of spanning trees and exact number of projected spanning trees and the proportion (given as percentage) between them.

Name	#Spanning Trees	#Projected Trees	Percent	#Successes out of 1000	Seconds
gn9e18d5w0w1	2,981	1,099	36.87	394	1
gn9e27d5w0w1	372,320	8,205	2.20	709	2
gn10e22d5w0w1	53,357	4,746	8.89	711	2
gn10e28d5w0w1	800,948	10,898	1.36	749	2
gn10e33d5w0w1	3,584,016	15,482	0.43	950	2
gn11e13d5w0w1	41	27	65.85	526	0
gn11e20d5w0w1	6,939	1,866	26.89	533	1
gn11e27d5w0w1	284,730	5,828	2.05	593	2
gn11e41d5w0w1	90,922,271	36,847	0.04	940	4
gn12e16d5w0w1	162	116	71.61	286	1
gn12e24d5w0w1	208,380	6,738	3.23	865	3
gn12e33d5w0w1	4,741,624	20,857	0.44	961	3
gn13e19d5w0w1	3,401	1,149	33.78	430	2
gn13e29d5w0w1	1,543,340	111,116	7.20	780	3
gn13e39d5w0w1	131,807,934	34,392	0.03	581	4
gn9e18d5w0w2	2,981	2,276	76.35	281	2
gn9e27d5w0w2	372,320	43,029	11.55	323	2
gn10e22d5w0w2	53,357	14,623	27.40	84	2
gn10e28d5w0w2	800,948	66,190	8.26	241	3
gn10e33d5w0w2	3,584,016	105,309	2.93	238	3
gn11e13d5w0w2	41	41	100.00	702	0
gn11e20d5w0w2	6,939	3,761	54.20	106	2
gn11e27d5w0w2	284,730	39,292	13.79	243	2
gn11e41d5w0w2	90,922,271	290,555	0.31	469	4
gn12e16d5w0w2	162	160	98.76	176	1
gn12e24d5w0w2	208,380	34,057	16.34	119	2
gn12e33d5w0w2	4,741,624	120,211	2.53	420	3
gn13e19d5w0w2	3,401	2,613	76.83	23	1
gn13e29d5w0w2	1,543,340	98,285	6.36	309	3
gn13e39d5w0w2	131,807,934	348,703	0.26	359	4
gn9e18d5w0w5	2,981	2,960	99.29	11	1
gn9e27d5w0w5	372,320	271,048	72.79	0	2
gn10e22d5w0w5	53,357	49,463	92.70	0	2
gn10e28d5w0w5	800,948	493,565	61.62	34	2
gn10e33d5w0w5	3,584,016	1,294,875	36.12	18	3
gn11e13d5w0w5	41	41	100.00	484	1
gn11e20d5w0w5	6,939	5,975	86.10	5	1
gn11e27d5w0w5	284,730	222,974	78.31	0	2
gn11e41d5w0w5	90,922,271	4,711,354	5.18	41	5
gn12e16d5w0w5	162	162	100.00	161	0
gn12e24d5w0w5	208,380	177,845	85.34	4	3
gn12e33d5w0w5	4,741,624	1,489,751	31.41	10	4
gn13e19d5w0w5	3,401	3,381	99.41	74	1
gn13e29d5w0w5	1,543,340	787,196	51.00	19	3
gn13e39d5w0w5	131,807,934	7,737,684	5.87	15	4

TABLE 4.30. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior rational non-integer point that is the image of some base. Here $d = 2$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d2w0w20	24(0)	153(0)	198(0)	203(2)	191(2)	222(2)	534(2)	564(2)	599(3)	621(2)	773(5)	1000(12)	1000(21)
gn9e27d2w0w20	8(0)	255(1)	279(1)	318(2)	374(3)	443(3)	483(4)	526(3)	581(4)	605(5)	888(8)	989(18)	1000(31)
gn10e22d2w0w20	16(1)	200(1)	246(1)	405(2)	491(2)	499(3)	518(3)	550(3)	602(3)	651(4)	711(7)	730(15)	734(27)
gn10e28d2w0w20	45(0)	350(1)	473(2)	585(2)	680(3)	737(3)	743(3)	769(4)	784(5)	832(5)	958(9)	1000(19)	1000(36)
gn10e33d2w0w20	17(0)	245(1)	329(1)	414(3)	449(2)	501(3)	541(4)	571(4)	589(5)	630(5)	934(11)	997(22)	998(40)
gn11e13d2w0w20	248(0)	252(0)	254(1)	306(1)	354(1)	364(1)	338(2)	655(3)	694(2)	721(2)	853(5)	851(9)	857(18)
gn11e20d2w0w20	0(0)	132(1)	149(2)	147(2)	158(2)	209(3)	213(3)	209(4)	223(3)	242(4)	568(9)	578(16)	557(28)
gn11e27d2w0w20	33(0)	223(1)	247(2)	249(2)	354(3)	355(3)	401(4)	434(4)	481(5)	489(5)	854(11)	963(22)	970(40)
gn11e41d2w0w20	15(1)	321(2)	479(2)	555(3)	619(3)	603(4)	712(6)	785(6)	817(7)	870(8)	970(13)	1000(30)	1000(55)
gn12e16d2w0w20	51(0)	231(1)	272(1)	604(1)	654(2)	859(2)	835(2)	814(3)	833(3)	831(3)	827(5)	824(11)	840(22)
gn12e24d2w0w20	30(0)	167(1)	191(2)	290(2)	320(3)	338(3)	429(4)	416(4)	420(5)	493(6)	613(10)	761(22)	782(41)
gn12e33d2w0w20	21(1)	214(2)	355(2)	412(2)	500(3)	498(4)	578(5)	607(5)	596(5)	621(6)	805(11)	859(25)	860(47)
gn13e19d2w0w20	6(0)	101(1)	145(2)	132(2)	137(3)	140(3)	158(3)	160(3)	162(4)	162(4)	147(8)	161(18)	160(33)
gn13e29d2w0w20	35(0)	194(2)	260(2)	295(3)	388(4)	463(4)	496(5)	558(6)	540(6)	629(6)	832(12)	987(29)	1000(54)
gn13e39d2w0w20	20(1)	239(2)	319(3)	416(3)	478(4)	579(6)	623(6)	672(7)	704(7)	734(8)	897(16)	998(35)	1000(65)
gn9e18d2w0w100	65(0)	417(1)	427(1)	421(1)	484(1)	479(2)	664(2)	682(2)	680(3)	669(2)	872(5)	963(11)	1000(21)
gn9e27d2w0w100	0(1)	0(1)	0(2)	0(2)	0(2)	25(3)	14(3)	24(3)	20(4)	16(4)	29(8)	31(16)	28(30)
gn10e22d2w0w100	0(0)	46(1)	56(2)	56(2)	50(3)	53(2)	59(3)	39(3)	33(4)	45(4)	43(8)	48(19)	52(34)
gn10e28d2w0w100	0(0)	0(1)	4(1)	5(2)	13(3)	10(4)	17(4)	19(4)	13(5)	24(6)	128(11)	205(26)	250(52)
gn10e33d2w0w100	2(1)	13(1)	9(2)	15(2)	6(3)	12(4)	23(4)	24(5)	21(5)	28(6)	58(11)	127(28)	198(51)
gn11e13d2w0w100	261(0)	366(1)	446(0)	880(1)	1000(2)	1000(2)	1000(1)	1000(1)	1000(2)	1000(3)	1000(4)	1000(9)	1000(18)
gn11e20d2w0w100	0(0)	0(1)	14(1)	28(2)	38(2)	54(2)	59(3)	49(3)	70(4)	59(4)	163(8)	419(20)	635(37)
gn11e27d2w0w100	0(1)	0(1)	0(2)	0(2)	2(2)	1(3)	1(4)	4(4)	5(5)	7(5)	19(11)	115(28)	234(57)
gn11e41d2w0w100	14(0)	20(2)	16(3)	25(3)	34(4)	64(4)	66(5)	60(6)	60(7)	61(8)	107(17)	219(43)	334(92)
gn12e16d2w0w100	17(0)	536(1)	496(1)	550(2)	537(2)	556(2)	524(2)	533(2)	505(2)	537(3)	1000(6)	1000(13)	1000(23)
gn12e24d2w0w100	0(1)	0(1)	0(2)	0(2)	1(3)	2(4)	0(4)	5(5)	17(5)	13(6)	13(12)	55(29)	99(52)
gn12e33d2w0w100	0(0)	11(1)	6(2)	7(3)	12(4)	14(4)	11(5)	19(6)	15(7)	20(9)	75(16)	182(40)	282(69)
gn13e19d2w0w100	20(1)	78(1)	86(1)	98(2)	86(2)	102(3)	95(3)	91(3)	89(4)	104(4)	88(7)	65(16)	96(32)
gn13e29d2w0w100	0(0)	16(1)	13(2)	24(3)	28(3)	22(4)	25(5)	20(5)	25(6)	23(6)	41(12)	42(25)	40(47)
gn13e39d2w0w100	1(1)	25(2)	25(3)	26(4)	22(4)	33(5)	35(7)	37(8)	57(9)	80(9)	124(20)	286(51)	483(97)
gn9e18d2w0w1000	0(0)	107(1)	71(1)	66(1)	155(1)	163(2)	172(2)	141(2)	283(3)	333(3)	347(5)	421(11)	862(26)
gn9e27d2w0w1000	0(0)	0(1)	0(1)	0(2)	0(2)	0(3)	0(3)	0(4)	0(4)	0(5)	0(11)	0(26)	0(44)
gn10e22d2w0w1000	0(1)	0(1)	0(1)	0(2)	0(2)	0(2)	0(3)	0(4)	0(4)	0(4)	3(9)	42(22)	98(45)
gn10e28d2w0w1000	0(1)	0(2)	0(2)	0(3)	0(3)	0(3)	0(3)	0(5)	0(5)	0(6)	0(11)	0(27)	90(59)
gn10e33d2w0w1000	0(0)	0(1)	0(2)	0(2)	0(3)	0(4)	0(4)	0(4)	0(5)	0(6)	0(12)	0(28)	0(61)
gn11e13d2w0w1000	294(1)	762(1)	875(1)	872(1)	1000(1)	1000(2)	1000(1)	1000(1)	1000(2)	1000(2)	1000(3)	1000(9)	1000(18)
gn11e20d2w0w1000	12(0)	12(1)	22(1)	10(1)	16(2)	12(3)	13(3)	29(4)	31(4)	25(4)	86(10)	126(20)	229(49)
gn11e27d2w0w1000	0(0)	0(1)	0(1)	0(2)	0(2)	0(4)	0(4)	0(5)	0(5)	0(6)	0(12)	0(32)	10(65)
gn11e41d2w0w1000	0(1)	0(1)	0(2)	0(4)	0(4)	0(5)	0(5)	0(8)	0(7)	0(9)	0(18)	4(46)	16(90)
gn12e16d2w0w1000	330(1)	350(1)	355(1)	346(1)	360(1)	498(2)	473(2)	534(2)	555(2)	503(3)	1000(7)	1000(13)	1000(23)
gn12e24d2w0w1000	0(0)	0(2)	0(2)	1(2)	1(3)	0(4)	3(5)	7(5)	4(5)	4(6)	7(12)	18(30)	30(57)
gn12e33d2w0w1000	0(1)	0(2)	0(2)	0(3)	0(3)	0(4)	0(5)	0(6)	0(7)	0(7)	0(14)	0(44)	4(81)
gn13e19d2w0w1000	0(0)	2(1)	1(2)	3(2)	5(3)	29(3)	29(4)	30(4)	25(4)	24(5)	105(10)	345(26)	391(42)
gn13e29d2w0w1000	0(1)	0(2)	1(2)	3(2)	0(4)	2(5)	1(5)	0(6)	1(6)	1(8)	3(16)	8(40)	4(79)
gn13e39d2w0w1000	0(1)	0(2)	0(3)	0(4)	0(5)	0(6)	0(7)	2(8)	2(9)	1(10)	7(19)	5(53)	3(104)

TABLE 4.31. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior rational non-integer point that is the image of some base. Here $d = 3$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d3w0w20	0(0)	50(1)	32(1)	50(2)	171(2)	172(1)	164(3)	173(3)	396(3)	437(4)	587(7)	1000(15)	1000(25)
gn9e27d3w0w20	0(0)	0(1)	0(2)	2(2)	28(3)	28(3)	41(4)	54(4)	42(5)	52(5)	189(11)	295(25)	557(55)
gn10e22d3w0w20	0(0)	15(2)	8(2)	22(2)	32(2)	39(3)	64(3)	50(4)	47(4)	41(4)	54(11)	174(30)	205(49)
gn10e28d3w0w20	0(0)	0(1)	0(2)	14(2)	23(4)	24(4)	23(4)	19(4)	32(5)	31(6)	34(13)	96(29)	381(62)
gn10e33d3w0w20	0(1)	2(1)	0(2)	1(3)	3(3)	12(4)	9(4)	4(6)	13(6)	14(6)	47(13)	77(28)	125(49)
gn11e13d3w0w20	177(0)	309(0)	366(1)	393(1)	389(1)	387(1)	386(1)	374(2)	394(2)	369(3)	384(4)	356(10)	359(18)
gn11e20d3w0w20	1(0)	7(1)	8(2)	14(1)	69(2)	98(3)	120(3)	180(4)	166(4)	212(5)	449(10)	761(22)	1000(42)
gn11e27d3w0w20	0(0)	2(1)	2(2)	22(2)	30(3)	32(4)	34(5)	38(6)	29(6)	48(6)	94(13)	213(30)	412(57)
gn11e41d3w0w20	0(1)	17(2)	34(3)	25(4)	25(5)	40(5)	42(7)	126(8)	158(9)	147(9)	213(19)	420(48)	645(94)
gn12e16d3w0w20	56(0)	223(1)	228(1)	216(1)	191(1)	609(2)	600(2)	609(2)	880(3)	861(3)	839(6)	857(13)	850(24)
gn12e24d3w0w20	10(0)	34(1)	41(2)	41(3)	36(3)	68(4)	46(5)	45(5)	52(5)	55(7)	73(32)	166(62)	
gn12e33d3w0w20	0(0)	12(2)	12(3)	33(4)	30(4)	18(5)	43(6)	46(7)	46(7)	49(8)	116(15)	257(37)	513(81)
gn13e19d3w0w20	1(0)	27(2)	16(2)	19(2)	19(2)	14(3)	32(3)	38(4)	38(5)	51(5)	292(14)	862(31)	910(50)
gn13e29d3w0w20	0(0)	31(1)	28(3)	55(3)	50(4)	64(5)	57(6)	68(7)	80(7)	78(8)	116(16)	253(38)	413(73)
gn13e39d3w0w20	0(0)	7(2)	17(4)	35(5)	37(5)	39(7)	48(7)	72(9)	65(10)	74(10)	148(21)	345(52)	554(106)
gn9e18d3w0w100	0(0)	35(1)	31(1)	117(1)	122(1)	113(2)	124(2)	128(2)	139(3)	128(3)	209(5)	639(15)	765(26)
gn9e27d3w0w100	0(0)	0(1)	0(1)	0(2)	0(3)	0(3)	0(4)	0(4)	0(5)	0(6)	0(9)	99(26)	94(47)
gn10e22d3w0w100	0(1)	2(1)	1(1)	20(2)	22(2)	13(3)	20(4)	22(4)	30(4)	25(4)	25(10)	109(24)	233(46)
gn10e28d3w0w100	0(0)	0(1)	0(2)	0(2)	1(3)	1(4)	4(5)	4(5)	9(5)	8(7)	12(11)	38(26)	32(49)
gn10e33d3w0w100	0(0)	0(1)	0(2)	0(2)	0(4)	0(4)	0(4)	0(5)	0(6)	0(6)	0(16)	0(35)	0(79)
gn11e13d3w0w100	249(1)	703(0)	1000(1)	1000(1)	1000(2)	1000(1)	1000(2)	1000(2)	1000(2)	1000(3)	1000(4)	1000(9)	1000(19)
gn11e20d3w0w100	0(1)	0(1)	0(2)	2(2)	45(2)	44(3)	43(3)	32(4)	42(4)	38(5)	42(9)	130(20)	202(39)
gn11e27d3w0w100	0(0)	0(1)	0(2)	0(3)	0(4)	0(4)	3(5)	2(5)	2(6)	10(7)	11(13)	33(31)	109(59)
gn11e41d3w0w100	0(0)	0(2)	0(3)	0(4)	0(5)	0(6)	0(7)	0(8)	0(8)	0(9)	0(21)	0(52)	2(112)
gn12e16d3w0w100	59(1)	829(1)	858(2)	978(2)	990(2)	983(2)	989(3)	986(2)	1000(3)	1000(3)	1000(5)	1000(12)	1000(23)
gn12e24d3w0w100	0(0)	1(1)	4(2)	1(2)	5(4)	4(4)	12(4)	13(5)	10(6)	15(6)	14(12)	15(29)	9(53)
gn12e33d3w0w100	0(1)	0(2)	0(3)	0(3)	0(5)	0(5)	0(6)	0(8)	0(8)	0(9)	0(17)	0(44)	0(92)
gn13e19d3w0w100	2(1)	70(1)	75(2)	127(3)	154(3)	235(3)	276(4)	290(5)	281(5)	269(5)	419(10)	962(28)	1000(45)
gn13e29d3w0w100	0(1)	0(2)	0(2)	0(4)	0(4)	0(5)	0(6)	0(6)	0(6)	0(8)	0(14)	0(31)	0(57)
gn13e39d3w0w100	0(0)	0(2)	0(3)	0(4)	0(5)	0(6)	0(7)	0(8)	0(9)	0(10)	0(21)	0(48)	0(85)
gn9e18d3w0w1000	0(0)	225(1)	279(2)	281(1)	277(1)	275(2)	280(3)	281(3)	272(3)	268(3)	404(5)	547(13)	719(26)
gn9e27d3w0w1000	0(1)	0(1)	9(2)	15(2)	25(2)	21(3)	11(3)	11(4)	13(4)	11(5)	20(10)	13(27)	17(48)
gn10e22d3w0w1000	1(0)	12(1)	12(2)	18(2)	12(3)	18(3)	59(3)	58(3)	44(5)	63(4)	77(9)	89(19)	134(34)
gn10e28d3w0w1000	0(0)	0(1)	0(2)	0(3)	0(3)	0(4)	0(5)	0(5)	0(5)	0(6)	0(13)	1(32)	0(70)
gn10e33d3w0w1000	0(1)	0(1)	0(2)	0(3)	0(3)	0(4)	0(5)	0(5)	0(7)	0(8)	0(15)	12(37)	10(71)
gn11e13d3w0w1000	300(0)	403(1)	401(1)	398(0)	437(1)	795(2)	794(2)	790(2)	803(2)	841(2)	1000(5)	1000(10)	1000(19)
gn11e20d3w0w1000	0(0)	0(1)	0(1)	0(2)	0(3)	7(3)	13(3)	9(4)	8(4)	52(4)	68(8)	73(16)	82(30)
gn11e27d3w0w1000	0(1)	5(2)	3(2)	4(2)	5(3)	3(4)	2(4)	3(5)	2(6)	1(6)	4(12)	3(35)	2(71)
gn11e41d3w0w1000	0(0)	0(2)	0(3)	0(4)	0(5)	0(5)	0(6)	0(8)	0(9)	0(10)	0(20)	0(55)	0(103)
gn12e16d3w0w1000	65(1)	570(1)	576(1)	629(2)	650(2)	645(2)	936(3)	924(3)	921(3)	1000(3)	1000(6)	1000(12)	1000(25)
gn12e24d3w0w1000	0(1)	1(2)	0(2)	5(2)	6(3)	4(4)	18(4)	11(6)	13(6)	20(7)	13(36)	10(71)	13(77)
gn12e33d3w0w1000	0(0)	0(1)	0(2)	0(4)	0(4)	0(4)	0(6)	0(6)	0(8)	0(8)	2(41)	13(77)	
gn13e19d3w0w1000	5(0)	45(1)	38(1)	36(2)	134(3)	274(4)	228(4)	261(5)	300(5)	300(6)	398(11)	915(28)	1000(46)
gn13e29d3w0w1000	0(1)	0(1)	0(3)	1(3)	0(4)	1(5)	1(6)	2(7)	0(7)	0(8)	1(16)	0(37)	1(65)
gn13e39d3w0w1000	0(1)	0(2)	0(4)	0(5)	0(5)	0(7)	0(7)	0(9)	0(10)	0(11)	0(23)	0(56)	0(106)

TABLE 4.32. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^2$, where for each instance, $\hat{x}$ is an interior rational non-integer point that is the image of some base. Here $d = 5$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d5w0w1	0(0)	630(1)	741(1)	934(1)	941(2)	946(2)	925(2)	1000(3)	1000(3)	1000(3)	1000(6)	1000(12)	1000(23)
gn9e27d5w0w1	44(0)	776(2)	870(2)	887(2)	906(3)	940(3)	939(3)	936(4)	921(4)	937(4)	1000(9)	1000(19)	1000(37)
gn10e22d5w0w1	91(1)	724(1)	816(1)	851(2)	949(2)	960(2)	954(3)	962(3)	968(4)	983(5)	1000(8)	1000(16)	1000(33)
gn10e28d5w0w1	125(0)	755(2)	757(2)	834(3)	877(3)	901(4)	965(4)	962(4)	976(4)	998(5)	1000(9)	1000(22)	1000(43)
gn10e33d5w0w1	70(0)	961(1)	1000(2)	1000(3)	1000(3)	1000(4)	1000(4)	1000(5)	1000(5)	1000(10)	1000(10)	1000(24)	1000(47)
gn11e13d5w0w1	125(0)	530(0)	1000(1)	1000(2)	1000(2)	1000(2)	1000(1)	1000(2)	1000(2)	1000(2)	1000(4)	1000(10)	1000(19)
gn11e20d5w0w1	99(0)	531(1)	665(1)	747(1)	728(2)	753(3)	781(3)	932(3)	953(4)	991(4)	1000(8)	1000(17)	1000(32)
gn11e27d5w0w1	42(1)	601(1)	743(2)	796(3)	785(3)	826(3)	829(4)	857(4)	836(4)	844(5)	1000(14)	1000(24)	1000(47)
gn11e41d5w0w1	40(0)	957(3)	973(3)	979(4)	994(4)	997(5)	997(6)	1000(7)	1000(7)	1000(7)	1000(14)	1000(36)	1000(70)
gn12e16d5w0w1	77(1)	309(1)	335(1)	562(2)	565(2)	556(2)	579(3)	567(3)	567(3)	545(3)	1000(6)	1000(15)	1000(28)
gn12e24d5w0w1	32(1)	866(1)	956(3)	966(3)	975(3)	990(4)	1000(4)	1000(5)	1000(6)	1000(6)	1000(10)	1000(24)	1000(45)
gn12e33d5w0w1	65(0)	954(2)	993(3)	996(3)	996(4)	993(4)	998(5)	1000(5)	1000(6)	1000(7)	1000(12)	1000(30)	1000(58)
gn13e19d5w0w1	118(1)	478(1)	628(1)	703(2)	719(3)	773(3)	782(3)	783(4)	797(4)	776(5)	782(7)	779(18)	790(35)
gn13e29d5w0w1	85(1)	767(2)	850(3)	867(4)	905(4)	950(5)	936(5)	957(6)	961(7)	963(7)	986(12)	986(30)	978(57)
gn13e39d5w0w1	19(1)	593(3)	666(3)	723(4)	734(6)	792(6)	817(7)	846(8)	838(9)	856(10)	942(18)	979(41)	1000(80)
gn9e18d5w0w2	61(0)	293(1)	354(1)	819(1)	859(2)	862(2)	962(3)	964(3)	1000(4)	1000(3)	1000(6)	1000(12)	1000(23)
gn9e27d5w0w2	0(0)	323(1)	548(2)	702(3)	745(3)	786(3)	820(4)	841(4)	840(5)	899(5)	946(9)	965(20)	965(37)
gn10e22d5w0w2	1(0)	80(1)	120(2)	130(2)	157(2)	132(2)	192(4)	185(3)	216(4)	227(4)	283(8)	355(17)	371(33)
gn10e28d5w0w2	20(0)	245(1)	345(2)	446(2)	442(3)	598(4)	640(4)	642(5)	688(5)	726(6)	851(11)	984(25)	992(45)
gn10e33d5w0w2	6(1)	193(2)	453(3)	537(3)	629(5)	709(5)	724(5)	784(6)	836(7)	838(7)	879(12)	892(26)	896(48)
gn11e13d5w0w2	331(1)	760(1)	1000(1)	1000(2)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(3)	1000(4)	1000(10)	1000(19)
gn11e20d5w0w2	10(0)	165(1)	271(2)	338(2)	697(3)	701(3)	851(4)	868(5)	938(5)	945(5)	994(9)	1000(19)	1000(35)
gn11e27d5w0w2	17(0)	256(2)	290(2)	414(3)	479(4)	506(4)	514(5)	517(5)	549(6)	622(7)	802(12)	929(26)	933(48)
gn11e41d5w0w2	7(0)	428(2)	478(4)	601(4)	643(6)	670(6)	757(7)	802(8)	833(9)	870(10)	924(18)	956(38)	956(74)
gn12e16d5w0w2	51(0)	239(1)	245(1)	230(1)	412(3)	410(3)	961(3)	1000(4)	1000(4)	1000(4)	1000(7)	1000(14)	1000(26)
gn12e24d5w0w2	6(1)	145(2)	233(2)	350(3)	356(4)	407(4)	494(6)	494(6)	564(6)	567(7)	777(13)	938(29)	942(50)
gn12e33d5w0w2	101(1)	401(2)	544(2)	562(4)	594(4)	647(4)	752(6)	812(6)	816(7)	827(8)	872(14)	901(30)	920(57)
gn13e19d5w0w2	0(0)	30(1)	25(1)	43(3)	34(3)	39(3)	39(4)	50(5)	44(4)	40(6)	34(11)	303(30)	304(47)
gn13e29d5w0w2	4(1)	302(2)	416(3)	531(4)	607(5)	685(6)	761(7)	795(7)	815(8)	847(9)	958(15)	998(32)	1000(59)
gn13e39d5w0w2	3(1)	363(3)	492(4)	577(5)	616(6)	638(7)	668(8)	707(8)	730(9)	761(10)	896(20)	975(43)	981(83)
gn9e18d5w0w5	0(0)	33(1)	34(1)	35(1)	39(1)	47(3)	36(3)	43(3)	27(3)	35(4)	321(8)	581(17)	744(33)
gn9e27d5w0w5	0(1)	0(1)	4(2)	12(3)	4(3)	7(4)	7(5)	45(6)	42(6)	48(7)	77(11)	154(31)	298(58)
gn10e22d5w0w5	0(0)	0(1)	0(2)	0(2)	0(3)	0(4)	9(4)	10(5)	6(5)	13(6)	70(10)	142(22)	143(37)
gn10e28d5w0w5	0(1)	4(1)	8(3)	25(3)	39(4)	48(4)	48(5)	51(7)	41(7)	53(8)	118(16)	313(39)	643(80)
gn10e33d5w0w5	0(0)	1(2)	14(3)	10(4)	20(4)	25(5)	33(6)	42(7)	56(9)	64(9)	115(17)	174(38)	566(78)
gn11e13d5w0w5	313(0)	587(1)	613(1)	851(1)	838(2)	905(1)	913(2)	887(2)	909(2)	917(2)	903(4)	915(10)	902(19)
gn11e20d5w0w5	0(1)	17(2)	17(2)	18(2)	14(3)	66(3)	66(4)	57(4)	68(5)	67(5)	44(10)	63(23)	270(44)
gn11e27d5w0w5	0(0)	0(2)	0(3)	5(3)	4(4)	7(5)	8(5)	7(6)	4(7)	7(8)	80(16)	352(39)	697(80)
gn11e41d5w0w5	0(0)	30(3)	39(4)	39(5)	47(6)	53(7)	73(9)	52(9)	73(11)	80(12)	144(25)	308(61)	555(127)
gn12e16d5w0w5	59(1)	190(0)	516(1)	705(1)	985(2)	1000(2)	1000(3)	1000(3)	1000(3)	1000(4)	1000(6)	1000(14)	1000(25)
gn12e24d5w0w5	0(0)	1(2)	1(3)	2(3)	0(5)	0(4)	1(5)	0(6)	2(7)	0(8)	5(18)	395(44)	527(76)
gn12e33d5w0w5	0(1)	30(2)	35(4)	35(4)	36(5)	56(7)	49(7)	43(8)	51(9)	61(10)	103(20)	190(53)	280(95)
gn13e19d5w0w5	0(0)	37(1)	90(2)	179(2)	138(3)	178(4)	232(4)	426(5)	516(5)	513(6)	836(11)	952(23)	950(41)
gn13e29d5w0w5	0(0)	16(3)	24(3)	35(4)	31(5)	61(6)	55(7)	56(9)	60(9)	70(10)	104(19)	309(48)	531(99)
gn13e39d5w0w5	0(1)	11(3)	26(4)	30(6)	45(7)	54(8)	51(10)	60(10)	94(12)	93(14)	180(27)	380(66)	567(118)

TABLE 4.33. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior rational non-integer point that is the image of some base. Here $d = 2$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d2w0w20	32(0)	158(1)	162(1)	238(1)	210(2)	233(2)	527(2)	541(3)	631(3)	584(3)	811(6)	1000(13)	1000(23)
gn9e27d2w0w20	11(0)	187(1)	285(1)	305(2)	402(2)	461(3)	520(3)	576(4)	609(4)	621(4)	805(8)	1000(19)	1000(34)
gn10e22d2w0w20	9(0)	194(1)	234(1)	443(2)	512(2)	532(2)	541(3)	555(3)	632(4)	646(4)	715(7)	768(15)	767(29)
gn10e28d2w0w20	32(1)	358(1)	477(2)	568(3)	625(3)	720(4)	743(3)	796(4)	836(4)	847(6)	990(9)	998(20)	1000(37)
gn10e33d2w0w20	15(1)	232(1)	333(2)	449(2)	581(3)	591(3)	598(4)	674(5)	693(6)	716(6)	841(11)	997(24)	999(44)
gn11e13d2w0w20	239(0)	251(1)	432(1)	475(1)	645(1)	650(1)	648(1)	693(2)	800(3)	876(2)	1000(4)	1000(10)	1000(19)
gn11e20d2w0w20	0(0)	123(1)	138(1)	168(2)	137(2)	203(2)	176(3)	208(3)	189(4)	231(4)	365(8)	358(16)	364(28)
gn11e27d2w0w20	24(0)	237(1)	288(1)	280(2)	379(3)	435(3)	470(4)	517(4)	511(5)	560(6)	857(11)	957(22)	982(42)
gn11e41d2w0w20	17(1)	398(1)	529(3)	555(3)	608(4)	623(4)	695(5)	796(6)	805(7)	846(7)	949(14)	1000(31)	1000(61)
gn12e16d2w0w20	52(0)	269(1)	257(1)	573(2)	820(2)	837(3)	834(2)	833(3)	833(3)	839(5)	824(12)	836(22)	824(12)
gn12e24d2w0w20	33(0)	147(1)	209(2)	264(2)	291(3)	363(4)	453(4)	437(4)	484(5)	475(6)	631(11)	741(23)	769(43)
gn12e33d2w0w20	11(1)	212(1)	347(2)	401(3)	454(3)	489(4)	589(4)	596(5)	583(6)	598(6)	821(12)	839(26)	855(49)
gn13e19d2w0w20	8(0)	77(1)	140(2)	147(2)	238(3)	242(3)	235(3)	251(4)	235(5)	271(5)	311(8)	321(18)	308(34)
gn13e29d2w0w20	27(1)	179(1)	254(2)	306(3)	360(4)	451(5)	475(5)	495(6)	529(7)	584(7)	825(14)	992(30)	996(55)
gn13e39d2w0w20	26(1)	217(2)	317(3)	437(4)	498(5)	598(6)	623(7)	666(8)	710(8)	738(8)	897(17)	996(37)	998(68)
gn9e18d2w0w100	58(1)	390(1)	405(1)	372(1)	387(1)	443(1)	454(2)	463(2)	447(3)	476(3)	699(6)	843(13)	1000(24)
gn9e27d2w0w100	0(0)	6(1)	6(1)	4(2)	3(2)	24(2)	14(3)	24(3)	15(4)	17(4)	34(8)	23(16)	31(31)
gn10e22d2w0w100	0(0)	50(1)	57(2)	52(2)	43(2)	48(3)	44(3)	60(4)	43(5)	38(5)	41(8)	48(22)	36(39)
gn10e28d2w0w100	0(0)	31(1)	28(2)	27(2)	30(3)	20(3)	30(4)	39(5)	39(6)	34(6)	74(12)	169(30)	329(65)
gn10e33d2w0w100	3(0)	8(1)	8(1)	10(3)	24(3)	32(4)	30(4)	20(4)	35(5)	20(6)	29(12)	131(30)	151(52)
gn11e13d2w0w100	208(1)	409(1)	905(1)	888(2)	1000(2)	1000(1)	1000(2)	1000(1)	1000(2)	1000(2)	1000(5)	1000(10)	1000(18)
gn11e20d2w0w100	0(1)	0(1)	21(1)	18(2)	59(2)	67(2)	56(3)	57(3)	58(3)	63(4)	229(9)	598(21)	897(43)
gn11e27d2w0w100	0(0)	0(1)	0(2)	0(2)	2(2)	4(3)	1(4)	3(5)	0(5)	5(6)	9(12)	64(32)	185(64)
gn11e41d2w0w100	10(1)	20(2)	28(2)	39(3)	41(5)	65(5)	55(6)	70(7)	67(7)	84(8)	156(19)	286(46)	564(92)
gn12e16d2w0w100	30(1)	492(1)	518(1)	518(1)	537(2)	537(2)	507(2)	562(3)	517(3)	531(3)	614(6)	638(12)	633(24)
gn12e24d2w0w100	0(0)	0(1)	0(2)	0(3)	0(3)	0(3)	0(4)	4(5)	3(6)	3(6)	14(13)	102(31)	122(55)
gn12e33d2w0w100	0(0)	6(1)	5(2)	22(3)	53(4)	51(5)	42(6)	34(7)	51(8)	72(8)	101(16)	344(42)	494(82)
gn13e19d2w0w100	25(1)	89(1)	82(2)	70(2)	84(3)	100(3)	109(3)	85(3)	92(4)	107(4)	88(7)	102(17)	85(34)
gn13e29d2w0w100	1(0)	11(1)	21(3)	13(3)	40(3)	23(4)	20(4)	21(5)	26(6)	40(7)	50(13)	58(26)	39(49)
gn13e39d2w0w100	3(1)	32(2)	29(3)	33(4)	45(5)	33(5)	39(7)	50(8)	49(9)	100(10)	153(21)	309(53)	522(104)
gn9e18d2w0w1000	0(0)	70(1)	65(1)	75(1)	169(2)	150(2)	156(3)	167(3)	257(3)	314(3)	308(5)	367(12)	929(32)
gn9e27d2w0w1000	0(1)	0(1)	0(2)	0(2)	0(3)	0(3)	0(4)	0(4)	0(4)	0(4)	0(11)	0(28)	0(50)
gn10e22d2w0w1000	0(0)	0(1)	0(1)	0(2)	0(2)	1(3)	11(4)	8(4)	8(4)	21(5)	25(9)	34(22)	112(46)
gn10e28d2w0w1000	0(1)	0(2)	0(2)	0(2)	0(3)	0(4)	0(4)	0(5)	0(5)	0(6)	0(12)	0(30)	0(59)
gn10e33d2w0w1000	0(0)	0(1)	0(2)	0(2)	0(3)	0(4)	0(4)	0(6)	0(6)	0(6)	0(12)	0(35)	0(73)
gn11e13d2w0w1000	291(0)	442(0)	877(0)	863(1)	1000(1)	1000(2)	1000(1)	1000(2)	1000(2)	1000(2)	1000(4)	1000(9)	1000(18)
gn11e20d2w0w1000	6(0)	17(1)	20(1)	11(2)	20(2)	10(3)	10(2)	24(4)	28(4)	36(4)	52(9)	248(22)	297(40)
gn11e27d2w0w1000	0(0)	0(2)	0(1)	0(3)	0(3)	0(4)	0(4)	0(5)	0(6)	0(6)	0(12)	0(31)	0(61)
gn11e41d2w0w1000	0(1)	0(2)	0(3)	0(4)	0(4)	0(6)	0(6)	0(7)	0(8)	0(9)	0(20)	1(54)	1(106)
gn12e16d2w0w1000	345(0)	361(1)	376(1)	359(2)	355(2)	351(1)	360(2)	354(2)	354(2)	390(3)	775(7)	1000(15)	1000(26)
gn12e24d2w0w1000	0(0)	0(2)	0(2)	0(2)	2(3)	1(3)	4(4)	5(5)	10(5)	12(6)	7(12)	26(31)	39(63)
gn12e33d2w0w1000	0(1)	0(2)	0(2)	0(3)	0(3)	0(5)	2(6)	1(6)	1(6)	2(7)	2(15)	7(48)	7(92)
gn13e19d2w0w1000	0(0)	15(1)	6(1)	15(3)	14(2)	32(3)	38(3)	48(4)	69(4)	74(5)	112(10)	142(22)	568(48)
gn13e29d2w0w1000	0(0)	0(2)	0(2)	1(3)	1(4)	4(5)	2(5)	0(7)	2(7)	1(16)	0(41)	0(41)	2(81)
gn13e39d2w0w1000	0(1)	0(2)	0(3)	0(4)	0(6)	0(6)	0(7)	0(8)	0(10)	0(10)	2(22)	8(54)	24(114)

TABLE 4.34. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior rational non-integer point that is the image of some base. Here $d = 3$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d3w0w1000	0(0)	0(0)	0(1)	0(1)	0(2)	0(2)	0(3)	0(4)	125(4)	141(5)	703(9)	830(17)	1000(29)
gn9e27d3w0w1000	0(0)	0(1)	1(2)	6(2)	23(3)	39(3)	43(5)	45(4)	57(6)	61(6)	215(12)	445(29)	608(53)
gn10e22d3w0w1000	0(1)	15(1)	16(1)	13(2)	19(2)	16(3)	16(4)	14(4)	13(5)	17(5)	43(11)	109(25)	98(46)
gn10e28d3w0w1000	0(0)	0(2)	0(2)	10(3)	25(3)	21(4)	18(5)	25(6)	23(6)	13(6)	23(12)	65(30)	169(59)
gn10e33d3w0w1000	0(0)	3(1)	0(2)	1(3)	0(3)	11(5)	8(5)	15(6)	7(6)	9(7)	28(14)	60(34)	138(62)
gn11e13d3w0w1000	175(1)	285(1)	345(0)	387(1)	382(2)	377(1)	393(1)	396(2)	354(2)	407(3)	412(4)	398(10)	397(19)
gn11e20d3w0w1000	3(1)	11(1)	11(1)	9(2)	11(3)	134(4)	151(4)	208(4)	256(5)	229(5)	412(10)	1000(25)	1000(40)
gn11e27d3w0w1000	0(1)	2(2)	5(2)	24(3)	38(3)	46(4)	37(4)	54(5)	63(6)	61(6)	119(14)	242(31)	352(63)
gn11e41d3w0w1000	0(1)	29(2)	22(3)	27(4)	29(5)	25(6)	23(8)	27(9)	53(9)	92(11)	144(22)	359(53)	718(106)
gn12e16d3w0w1000	52(1)	130(0)	176(1)	183(1)	184(2)	578(2)	622(3)	570(3)	860(4)	870(4)	856(6)	844(14)	868(26)
gn12e24d3w0w1000	3(0)	18(1)	32(2)	31(3)	36(3)	44(4)	38(4)	35(5)	37(6)	56(7)	56(14)	65(33)	76(58)
gn12e33d3w0w1000	0(1)	14(2)	13(3)	21(3)	35(5)	27(5)	45(7)	59(7)	93(8)	79(9)	151(16)	375(42)	479(75)
gn13e19d3w0w1000	1(1)	25(1)	22(2)	20(2)	17(3)	26(3)	23(4)	17(4)	21(5)	37(5)	165(12)	697(30)	966(55)
gn13e29d3w0w1000	2(1)	20(2)	30(2)	51(3)	40(4)	64(5)	62(6)	68(7)	54(8)	70(9)	118(17)	197(41)	285(78)
gn13e39d3w0w1000	0(0)	10(3)	27(4)	31(4)	38(5)	48(7)	52(8)	70(9)	77(10)	98(11)	166(23)	423(60)	603(110)
gn9e18d3w0w100	0(1)	41(0)	37(1)	118(2)	107(2)	102(2)	108(2)	106(2)	114(3)	139(3)	175(6)	392(14)	763(30)
gn9e27d3w0w100	0(0)	2(1)	4(2)	4(2)	4(3)	3(4)	2(4)	1(4)	3(5)	3(5)	2(11)	37(30)	63(59)
gn10e22d3w0w100	0(0)	2(1)	1(2)	13(3)	18(3)	14(3)	14(3)	18(4)	21(4)	25(6)	19(11)	24(26)	245(50)
gn10e28d3w0w100	0(1)	0(1)	0(2)	0(2)	0(4)	0(4)	1(5)	3(6)	9(6)	5(7)	32(13)	61(29)	110(52)
gn10e33d3w0w100	0(1)	0(2)	0(2)	0(2)	0(4)	0(4)	0(5)	0(6)	0(7)	0(8)	0(16)	0(36)	0(79)
gn11e13d3w0w100	244(0)	715(1)	1000(0)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(2)	1000(4)	1000(10)	1000(19)
gn11e20d3w0w100	0(0)	0(1)	0(1)	0(2)	36(3)	33(3)	44(4)	28(5)	35(4)	49(5)	45(10)	30(18)	360(44)
gn11e27d3w0w100	0(0)	0(1)	0(2)	0(3)	0(4)	0(4)	0(5)	0(6)	0(6)	0(6)	0(13)	29(33)	52(56)
gn11e41d3w0w100	0(1)	0(2)	0(3)	0(4)	0(5)	0(6)	0(7)	0(8)	1(9)	0(11)	1(20)	0(53)	10(114)
gn12e16d3w0w100	59(0)	677(1)	743(2)	887(2)	898(2)	889(2)	895(2)	900(2)	894(3)	893(3)	1000(5)	1000(13)	1000(25)
gn12e24d3w0w100	0(1)	0(1)	0(2)	1(3)	0(3)	1(4)	4(5)	1(5)	1(6)	2(7)	18(13)	18(30)	16(53)
gn12e33d3w0w100	0(0)	0(1)	0(3)	0(4)	0(4)	0(6)	0(6)	0(8)	0(8)	0(9)	0(18)	0(48)	0(90)
gn13e19d3w0w100	6(0)	14(1)	16(2)	9(3)	13(3)	30(3)	23(4)	18(5)	18(5)	57(6)	297(12)	452(29)	781(54)
gn13e29d3w0w100	0(0)	0(2)	0(3)	0(4)	0(4)	3(5)	3(6)	2(7)	4(7)	3(8)	1(15)	6(32)	3(58)
gn13e39d3w0w100	0(1)	0(2)	0(3)	0(5)	0(5)	0(7)	0(8)	0(9)	0(9)	0(10)	0(21)	0(47)	0(85)
gn9e18d3w0w1000	0(0)	139(1)	181(1)	217(2)	239(2)	248(3)	221(3)	213(3)	234(3)	220(3)	340(7)	488(16)	858(32)
gn9e27d3w0w1000	0(0)	0(1)	14(1)	17(2)	21(3)	18(3)	22(3)	16(4)	23(5)	21(5)	28(11)	54(30)	51(53)
gn10e22d3w0w1000	2(1)	14(1)	15(2)	9(2)	23(3)	22(3)	26(4)	25(4)	21(4)	25(4)	73(9)	180(20)	256(37)
gn10e28d3w0w1000	0(0)	0(2)	0(2)	0(3)	0(3)	0(4)	0(5)	0(5)	0(6)	0(7)	1(13)	60(32)	133(64)
gn10e33d3w0w1000	0(0)	0(1)	0(2)	0(3)	0(4)	0(4)	0(5)	0(6)	0(6)	0(8)	0(17)	0(39)	0(78)
gn11e13d3w0w1000	318(0)	406(0)	411(0)	415(1)	392(2)	409(2)	528(2)	539(2)	552(2)	552(3)	726(4)	714(11)	733(20)
gn11e20d3w0w1000	0(1)	0(1)	0(1)	0(2)	0(3)	11(4)	7(4)	10(4)	8(5)	73(5)	123(9)	122(18)	117(33)
gn11e27d3w0w1000	0(1)	1(2)	0(3)	0(3)	2(3)	2(4)	1(5)	0(6)	2(7)	0(7)	5(14)	2(33)	2(69)
gn11e41d3w0w1000	0(0)	0(2)	0(3)	0(4)	0(5)	0(7)	0(7)	0(9)	0(10)	0(10)	0(22)	0(59)	0(114)
gn12e16d3w0w1000	63(1)	587(1)	576(1)	651(1)	654(1)	616(2)	926(3)	927(3)	935(4)	1000(4)	1000(6)	1000(14)	1000(26)
gn12e24d3w0w1000	0(0)	0(1)	0(2)	0(3)	1(4)	1(4)	0(5)	0(6)	1(7)	0(7)	1(15)	1(34)	0(71)
gn12e33d3w0w1000	0(0)	0(2)	0(2)	0(4)	0(4)	0(5)	0(6)	0(7)	0(8)	0(10)	0(19)	0(47)	0(88)
gn13e19d3w0w1000	0(1)	7(2)	17(1)	73(2)	68(2)	85(4)	63(5)	68(5)	90(6)	73(7)	85(11)	480(32)	809(59)
gn13e29d3w0w1000	0(1)	0(2)	0(3)	0(3)	0(5)	0(5)	0(6)	0(7)	0(8)	0(9)	0(17)	0(37)	0(67)
gn13e39d3w0w1000	0(1)	0(2)	0(3)	0(5)	0(6)	0(7)	0(8)	0(9)	0(10)	0(11)	0(24)	0(58)	0(109)

TABLE 4.35. Calibration test set: Tabu Search **Heuristic 85** minimizing $(x - \hat{x})^4$, where for each instance, $\hat{x}$ is an interior rational non-integer point that is the image of some base. Here $d = 5$ and all tests run on Fuzzy. Shown are the number of successes out of 1000 Tabu Searches for Tabu Search limits of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100 and seconds (in parenthesis) to perform the 1000 searches.

Name	Tabu Limit 1	Tabu Limit 2	Tabu Limit 3	Tabu Limit 4	Tabu Limit 5	Tabu Limit 6	Tabu Limit 7	Tabu Limit 8	Tabu Limit 9	Tabu Limit 10	Tabu Limit 20	Tabu Limit 50	Tabu Limit 100
gn9e18d5w0w1	0(0)	398(1)	730(2)	927(2)	1000(2)	1000(3)	1000(3)	1000(3)	1000(3)	1000(3)	1000(6)	1000(13)	1000(24)
gn9e27d5w0w1	45(0)	723(2)	800(2)	886(2)	889(3)	910(3)	915(4)	927(4)	910(4)	925(5)	1000(10)	1000(22)	1000(41)
gn10e22d5w0w1	71(0)	729(1)	848(2)	871(2)	941(3)	956(3)	964(3)	963(4)	959(4)	985(4)	1000(8)	1000(18)	1000(35)
gn10e28d5w0w1	126(1)	733(1)	795(3)	830(2)	891(3)	900(4)	962(4)	963(4)	985(5)	999(7)	1000(10)	1000(24)	1000(46)
gn10e33d5w0w1	81(1)	958(1)	999(2)	994(3)	1000(4)	1000(4)	1000(5)	1000(5)	1000(6)	1000(6)	1000(11)	1000(26)	1000(51)
gn11e13d5w0w1	146(1)	525(1)	1000(1)	1000(2)	1000(2)	1000(2)	1000(3)	1000(2)	1000(2)	1000(3)	1000(5)	1000(10)	1000(21)
gn11e20d5w0w1	103(0)	565(1)	661(1)	743(2)	725(2)	768(2)	773(3)	945(4)	946(4)	991(5)	1000(8)	1000(18)	1000(33)
gn11e27d5w0w1	36(1)	624(2)	744(3)	782(3)	784(4)	818(4)	838(5)	826(5)	853(6)	865(6)	968(11)	1000(25)	1000(49)
gn11e41d5w0w1	28(0)	945(2)	965(4)	984(4)	998(5)	995(6)	1000(6)	1000(7)	1000(8)	1000(8)	1000(16)	1000(38)	1000(76)
gn12e16d5w0w1	64(0)	310(0)	328(1)	337(2)	557(2)	569(2)	562(2)	549(2)	570(3)	554(4)	1000(8)	1000(16)	1000(29)
gn12e24d5w0w1	22(1)	869(2)	953(3)	979(3)	973(4)	994(4)	1000(5)	1000(5)	1000(5)	1000(6)	1000(11)	1000(25)	1000(49)
gn12e33d5w0w1	61(0)	965(2)	988(3)	991(4)	993(4)	997(5)	1000(5)	999(6)	1000(7)	999(7)	1000(14)	1000(32)	1000(63)
gn13e19d5w0w1	151(0)	457(2)	657(2)	724(2)	765(3)	740(3)	766(4)	793(4)	760(4)	778(5)	766(9)	775(19)	798(37)
gn13e29d5w0w1	81(1)	809(3)	872(3)	891(4)	934(4)	969(5)	986(6)	993(7)	989(6)	996(7)	999(13)	999(31)	1000(62)
gn13e39d5w0w1	17(1)	606(3)	674(4)	745(5)	745(6)	803(7)	789(7)	859(9)	845(9)	877(10)	940(19)	974(45)	1000(87)
gn9e18d5w0w2	63(1)	267(1)	432(1)	487(1)	601(2)	846(3)	879(3)	861(3)	889(4)	885(4)	910(6)	907(14)	910(24)
gn9e27d5w0w2	0(0)	330(1)	512(2)	592(3)	662(3)	741(4)	740(4)	805(5)	815(6)	886(6)	959(10)	948(21)	943(40)
gn10e22d5w0w2	1(1)	63(1)	153(1)	203(2)	208(2)	246(3)	247(3)	289(4)	299(4)	300(5)	389(8)	377(18)	392(35)
gn10e28d5w0w2	18(0)	222(1)	311(2)	440(3)	463(4)	517(4)	558(4)	618(5)	660(6)	625(7)	782(12)	905(27)	912(49)
gn10e33d5w0w2	5(0)	217(1)	403(3)	473(3)	568(4)	580(5)	693(6)	740(7)	768(7)	809(8)	880(14)	879(29)	860(53)
gn11e13d5w0w2	292(0)	755(0)	1000(1)	1000(1)	1000(1)	1000(2)	1000(1)	1000(2)	1000(3)	1000(2)	1000(4)	1000(10)	1000(20)
gn11e20d5w0w2	13(0)	105(1)	143(2)	182(2)	521(4)	614(3)	690(4)	680(4)	693(5)	741(6)	864(9)	1000(21)	1000(39)
gn11e27d5w0w2	15(0)	246(1)	305(2)	434(4)	481(4)	532(5)	553(5)	588(6)	662(6)	669(7)	818(13)	937(27)	929(50)
gn11e41d5w0w2	3(0)	452(3)	502(3)	611(5)	653(6)	701(7)	756(8)	801(9)	853(10)	883(11)	960(19)	988(42)	988(79)
gn12e16d5w0w2	55(0)	192(0)	206(1)	215(2)	991(3)	1000(4)	1000(4)	1000(4)	1000(4)	1000(4)	1000(7)	1000(15)	1000(28)
gn12e24d5w0w2	5(0)	136(2)	206(3)	300(3)	311(4)	360(5)	454(5)	494(6)	550(7)	601(8)	683(14)	958(32)	941(55)
gn12e33d5w0w2	119(1)	409(2)	509(3)	565(4)	589(4)	627(5)	680(6)	725(7)	768(8)	822(9)	851(15)	894(34)	890(62)
gn13e19d5w0w2	0(1)	33(1)	22(2)	33(3)	39(4)	44(4)	50(4)	39(5)	47(6)	47(6)	47(12)	54(26)	61(44)
gn13e29d5w0w2	1(0)	280(2)	400(4)	520(4)	561(6)	709(6)	726(7)	735(8)	779(8)	778(9)	964(17)	995(34)	994(65)
gn13e39d5w0w2	3(1)	356(3)	473(4)	529(6)	577(6)	650(7)	664(9)	736(9)	753(11)	755(12)	914(21)	986(46)	993(87)
gn9e18d5w0w5	0(0)	14(1)	17(1)	279(2)	279(2)	271(3)	282(3)	271(4)	362(4)	345(5)	369(7)	933(20)	1000(33)
gn9e27d5w0w5	0(0)	0(2)	0(2)	0(3)	0(4)	0(4)	0(5)	74(6)	79(6)	82(7)	213(14)	670(39)	804(61)
gn10e22d5w0w5	0(0)	0(1)	0(2)	0(2)	0(3)	0(4)	0(4)	0(5)	0(6)	1(6)	22(11)	18(20)	30(36)
gn10e28d5w0w5	0(1)	40(2)	51(3)	50(3)	84(4)	83(5)	114(6)	81(6)	104(7)	149(8)	247(18)	446(40)	551(69)
gn10e33d5w0w5	0(1)	13(2)	15(3)	11(4)	101(5)	131(5)	151(6)	179(7)	208(9)	263(9)	280(18)	603(45)	871(84)
gn11e13d5w0w5	292(1)	589(0)	749(1)	959(1)	1000(1)	1000(1)	1000(1)	1000(2)	1000(2)	1000(3)	1000(4)	1000(11)	1000(20)
gn11e20d5w0w5	0(0)	9(1)	10(2)	5(2)	7(3)	71(4)	85(3)	88(5)	93(5)	127(5)	113(11)	141(23)	228(42)
gn11e27d5w0w5	0(1)	0(2)	0(3)	5(3)	5(4)	6(5)	8(6)	13(7)	11(8)	18(9)	52(17)	382(45)	644(82)
gn11e41d5w0w5	0(1)	42(3)	51(5)	43(5)	48(6)	46(7)	77(9)	115(10)	103(12)	115(13)	208(26)	383(61)	560(118)
gn12e16d5w0w5	47(1)	169(1)	333(1)	574(2)	745(2)	886(2)	876(3)	985(4)	1000(4)	1000(4)	1000(7)	1000(14)	1000(28)
gn12e24d5w0w5	0(0)	5(2)	1(3)	0(3)	0(4)	2(5)	3(6)	1(7)	2(7)	1(7)	2(16)	0(43)	27(75)
gn12e33d5w0w5	0(1)	14(3)	15(4)	16(4)	33(6)	30(7)	82(8)	74(9)	88(9)	94(10)	144(21)	206(50)	278(90)
gn13e19d5w0w5	0(0)	73(1)	103(2)	154(3)	166(3)	160(3)	235(4)	242(4)	290(5)	384(6)	638(12)	902(26)	899(45)
gn13e29d5w0w5	0(1)	12(3)	14(3)	19(5)	20(5)	33(7)	44(7)	46(8)	48(10)	53(11)	71(20)	186(54)	286(97)
gn13e39d5w0w5	0(1)	12(3)	23(4)	26(6)	36(7)	41(9)	66(9)	77(12)	82(13)	106(14)	165(28)	422(67)	699(134)

TABLE 4.36. Two experimental design problems P_{20} and P_{28} and comparing run times of IBM Blue Gene/L vs Fuzzy to find all projected bases. We used **Heuristic 92** (DFBFS) with 100 searches, boundary retry limit 100, interior retry limit 10,000 and truncation depth 8.

Name	P_{20}	P_{28}
Columns(#matroid elements)	100	100
Rows(matroid rank)	20	28
Projected bases	13,816	19,193
Seconds for IBM Blue Gene/L	23,773	56,763
# Projected bases found by run 1 of Heuristic 92	13,757	19,193
Seconds for run 1 of Heuristic 92	923	2,557
# Projected bases found by run 2 of Heuristic 92	13,816	19,191
Seconds for run 2 of Heuristic 92	945	2,645

TABLE 4.37. Experimental design problem P_{20} and comparing run times of IBM Blue Gene/L vs Fuzzy to find all projected bases. We partitioned the region of feasible projected bases into 12 disjoint regions and ran 12 separate concurrently running instances of **Heuristic 86** (PT) on Fuzzy to find as many projected bases as possible. When running **Heuristic 86** (PT) we used **Heuristic 85** (TS) and tried each point 10 times with a Tabu limit of 20.

Region	#Projected Bases Found	Seconds
Box1	1,044	9,544
Box2	1,122	5,086
Box3	1,122	4,966
Box4	1,135	8,066
Box5	1,122	5,414
Box6	1,122	4,540
Box7	1,122	4,528
Box8	1,188	5,614
Box9	1,147	9,181
Box10	1,124	5,232
Box11	1,124	5,252
Box12	1,249	8,173
TOTAL	13,816	(Longest time)9,544

TABLE 4.38. Machines Used in Experiments.

Name	#CPUS	Clock Speed	Memory
Fuzzy	16 AMD	2400 MHz	64GB
Truth	16 AMD	2400 MHz	64GB
Mathocr01	2 AMD	2000 MHz	4GB
Mathocr02	2 AMD	2000 MHz	4GB
Mathocr03	2 AMD	2000 MHz	4GB
Mathocr04	2 AMD	2000 MHz	12GB

Bibliography

- [ABB⁺99] E. Anderson, Z. Bai, C. Bischof, S. Blackford, J. Demmel, J. Dongarra, J. Du Croz, A. Greenbaum, S. Hammarling, A. McKenney, and D. Sorensen, *LAPACK users' guide*, third ed., Society for Industrial and Applied Mathematics, Philadelphia, PA, 1999.
- [Bar94] A. I. Barvinok, *Polynomial time algorithm for counting integral points in polyhedra when the dimension is fixed*, Mathematics of Operations Research **19** (1994), 769–779.
- [Bar06] ———, *Computing the Ehrhart quasi-polynomial of a rational simplex*, Math. Comp. **75** (2006), no. 255, 1449–1466 (electronic).
- [BGW07] A. V. Borovik, I. M. Gelfand, and N. White, *Coxeter matroid polytopes*, Annals of Combinatorics **1** (2007), no. 1, 123–134.
- [BHS09] M. Beck, C. Haase, and F. Sottile, *Formulas of Brion, Lawrence, and Varchenko on rational generating functions for cones*, The Mathematical Intelligencer **31** (2009), no. 1, 9–17.
- [BJR06] L. J. Billera, N. Jia, and V. Reiner, *A quasisymmetric function for matroids*, eprint arXiv:math/0606646, 2006.
- [Bla08] J. Blasiak, *The toric ideal of a graphic matroid is generated by quadrics*, Combinatorica **28** (2008), no. 3, 283–297.
- [BLMA⁺08] Y. Berstein, J. Lee, H. Maruri-Aguilar, S. Onn, E. Riccomagno, R. Weismantel, and H. Wynn, *Nonlinear matroid optimization and experimental design*, SIAM Journal on Discrete Mathematics **22** (2008), 901.
- [BLOW08] Y. Berstein, J. Lee, S. Onn, and R. Weismantel, *Nonlinear optimization for matroid intersection and extensions*, 2008.
- [BO08] Y. Berstein and S. Onn, *Nonlinear bipartite matching*, Discrete Optimization **5** (2008), 53.

-
- [BP99] A. I. Barvinok and J. E. Pommersheim, *An algorithmic theory of lattice points in polyhedra*, New Perspectives in Algebraic Combinatorics (L. J. Billera, A. Björner, C. Greene, R. E. Simion, and R. P. Stanley, eds.), Math. Sci. Res. Inst. Publ., vol. 38, Cambridge Univ. Press, Cambridge, 1999, pp. 91–147.
 - [BR07] M. Beck and S. Robins, *Computing the continuous discretely: Integer-point enumeration in polyhedra*, Springer, 2007.
 - [Bri88] M. Brion, *Points entiers dans les polyédres convexes*, Ann. Sci. École Norm. Sup. **21** (1988), no. 4, 653–663.
 - [BS07a] A. Barvinok and A. Samorodnitsky, *Random weighting, asymptotic counting, and inverse isoperimetry*, Israel Journal of Mathematics **158** (2007), 159–191.
 - [BS07b] M. Beck and F. Sottile, *Irrational proofs for three theorems of Stanley*, European Journal of Combinatorics **28** (2007), no. 1, 403–409, math.CO/0501359.
 - [BW91] G. Brightwell and P. Winkler, *Counting linear extensions*, Order **8** (1991), no. 3, 225–242.
 - [BW03] A. I. Barvinok and K. Woods, *Short rational generating functions for lattice point problems*, Journal of the AMS **16** (2003), no. 4, 957–979.
 - [CS86] W. Cook and J. F. A. Schrijver, *An integer analogue of caratheodory’s theorem*, Journal of Combinatorial Theory **510** (1986), 179–185.
 - [DF88] M. E. Dyer and A. M. Frieze, *On the complexity of computing the volume of a polyhedron*, SIAM J. Comput. **17** (1988), no. 5, 967–974.
 - [DHLO09] J. De Loera, D. C. Haws, J. Lee, and A. O’Hair, *Matroids Optimization Combinatorics Heuristics and Algorithms*, Available from URL <http://math.ucdavis.edu/~haws/MOCHA/>, 2009.
 - [DLHH⁺04] J. A. De Loera, D. Haws, R. Hemmecke, P. Huggins, B. Sturmfels, and R. Yoshida, *Short rational functions for toric algebra and applications*, Journal of Symbolic Computation **38** (2004), no. 2, 959–973.
 - [DLHH⁺05] J. A. De Loera, D. C. Haws, R. Hemmecke, P. Huggins, J. Tauzer, and R. Yoshida, *Software and user’s guide for latte v.1.1*, 2005.
 - [DLHK08] J. A. De Loera, D. C. Haws, and M. Köppe, *Ehrhart polynomials of matroid polytopes and polymatroids*, Discrete and Computational Geometry (2008).

-
- [DLHTY04] J. A. De Loera, R. Hemmecke, J. Tauzer, and R. Yoshida, *Effective lattice point counting in rational convex polytopes*, Journal of Symbolic Computation **38** (2004), no. 4, 1273–1302.
- [DLRS09] J. A. De Loera, J. Rambau, and F. Santos, *Triangulations: Applications, structures, algorithms*, Book manuscript, 2009.
- [DNH97] E. De Negri and T. Hibi, *Gorenstein algebras of Veronese type*, Journal of Algebra **193** (1997), no. 2, 629–639.
- [dPS03] J. C. de Pina and J. Soares, *Improved bound for the carathéodory rank of the bases of a matroid*, Journal of Combinatorial Theory **B** (2003), no. 88, 323–327.
- [Edm03] J. Edmonds, *Submodular functions, matroids, and certain polyhedra*, Combinatorial Optimization – Eureka, You Shrink!: Papers Dedicated to Jack Edmonds. 5th International Workshop, Aussois, France, March 5–9, 2001, Revised Papers (M. Jünger, G. Reinelt, and G. Rinaldi, eds.), Lecture notes in computer science, vol. 2570, Springer-Verlag, Berlin, 2003, pp. 11–26.
- [EG00] M. Ehrgott and X. Gandibleux, *A survey and annotated bibliography of multi-objective combinatorial optimization*, OR Spektrum **22** (2000), no. 4, 425–460.
- [Ehr96] M. Ehrgott, *On matroids with multiple objectives*, Optimization **38** (1996), no. 1, 73–84, Multicriteria optimization and decision theory (Holzhau, 1994).
- [Ele86] G. Elekes, *A geometric inequality and the complexity of computing volume*, Discrete Comput. Geom. **1** (1986), no. 4, 289–292.
- [FH80] A. Fries and W. Hunter, *Minimum aberration 2^{k-p} designs*, Techno. (1980), no. 22, 601–608.
- [FS05] E. M. Feichtner and B. Sturmfels, *Matroid polytopes, nested sets and Bergman fans*, Port. Math. **62** (2005), 437–468.
- [Fuk06] K. Fukuda, *cdd+, a C++ implementation of the double description method of Motzkin et al.*, Available from URL http://www.ifor.math.ethz.ch/~fukuda/cdd_home/cdd.html, 2006.
- [Gea09] T. Granlund and et al., *GNU multiple precision arithmetic library*, 2009, <http://gmp.lib.org/>.
- [GGMS87] I. M. Gelfand, M. Goresky, R. D. MacPherson, and V. V. Serganova, *Combinatorial geometries, convex polyhedra, and Schubert cells*, Adv. Math. **63** (1987),

301–316.

- [GJ79] M. Garey and D. Johnson, *Computers and intractability: A guide to the theory of NP-completeness*, W.H. Freeman and Company, San Francisco, 1979.
- [GLLR79] R. Graham, E. Lawler, J. Lenstra, and A. Rinnooy Kan, *Optimization and approximation in deterministic sequencing and scheduling: A survey*, Annals of Discrete Mathematics (1979), no. 5, 287–326.
- [GLM08] J. Gunnels, J. Lee, and S. Margulies, *IBM research report*, Preprint (2008).
- [Glo86] F. Glover, *Future paths for integer programming and links to artificial intelligence*, Computers and Operations Research **13** (1986), 533–549.
- [GO97] J. E. Goodman and J. O’Rourke (eds.), *Handbook of discrete and computational geometry*, CRC Press, Inc., Boca Raton, FL, USA, 1997.
- [Haw09] D. C. Haws, *Matroid polytopes*, <http://math.ucdavis.edu/~haws/Matroids/>, 2009.
- [HH72] C. Hotzmann and F. Harary, *On the tree graph of a matroid*, SIAM Journal of Applied Math **22** (1972), no. 2.
- [Hib92] T. Hibi, *Algebraic combinatorics on convex polytopes*, Carslaw Publications, Glebe, Australia, 1992.
- [Kat05] M. Katzman, *The Hilbert series of algebras of Veronese type*, Communications in Algebra **33** (2005), 1141–1146, math/0408038.
- [KC02] J. D. Knowles and D. W. Corne, *Enumeration of pareto optimal multi-criteria spanning trees - a proof of the incorrectness of zhou and gen’s proposed algorithm*, European Journal of Operational Research **143** (2002), no. 3, 543 – 547.
- [Kha93] L. Khachiyan, *Complexity of polytope volume computation*, New Trends in Discrete and Computational Geometry, Algorithms Combin., vol. 10, Springer, Berlin, 1993, pp. 91–101.
- [Köp07a] M. Köppe, *LattE macchiato, version 1.2-mk-0.9, an improved version of De Loera et al.’s LattE program for counting integer points in polyhedra with variants of Barvinok’s algorithm*, Available from URL <http://www.math.uni-magdeburg.de/~mkoeppel/latte/>, 2007.

-
- [Köp07b] M. Köppe, *A primal Barvinok algorithm based on irrational decompositions*, SIAM Journal on Discrete Mathematics **21** (2007), no. 1, 220–236, math.CO/0603308.
- [KV08] M. Köppe and S. Verdoolaege, *Computing parametric rational generating functions with a primal Barvinok algorithm*, The Electronic Journal of Combinatorics **15** (2008), #R16.
- [Law91a] J. Lawrence, *Polytope volume computation*, Math. Comp. **57** (1991), no. 195, 259–271.
- [Law91b] ———, *Rational-function-valued valuations on polyhedra*, Discrete and Computational Geometry (New Brunswick, NJ, 1989/1990), DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 6, Amer. Math. Soc., Providence, RI, 1991, pp. 199–208.
- [Mat97] T. Matsui, *A flexible algorithm for generating all the spanning trees in undirected graphs*, Algorithmica **18** (1997), no. 4, 530–543.
- [May08] D. Mayhew, *Matroid complexity and nonsuccinct descriptions*, SIAM J. Discret. Math. **22** (2008), no. 2, 455–466.
- [MRar] D. Mayhew and G. Royle, *Matroids with nine elements*, Journal of Combinatorial Theory ((To appear)).
- [Onn03] S. Onn, *Convex matroid optimization*, SIAM Journal on Discrete Mathematics **17** (2003), 249.
- [Oxl92] J. Oxley, *Matroid theory*, Oxford University Press, New York, NY, USA, 1992.
- [PWZ96] M. Petkovšek, H. S. Wilf, and D. Zeilberger, *$A = B$* , AK Peters, Ltd., 1996.
- [Ram02] J. Rambau, *TOPCOM: Triangulations of point configurations and oriented matroids*, Mathematical Software—ICMS 2002 (A. M. Cohen, X.-S. Gao, and N. Takayama, eds.), World Scientific, 2002, pp. 330–340.
- [Sch86a] A. Schrijver, *Theory of linear and integer programming*, Wiley-Interscience, 1986.
- [Sch86b] A. Schrijver, *Theory of linear and integer programming*, John Wiley & Sons, Inc., New York, NY, USA, 1986.
- [Sch03] A. Schrijver, *Combinatorial optimization: Polyhedra and efficiency*, Springer, 2003.

-
- [Seb90] A. Sebö, *Hilbert bases, Carathéodory's theorem and combinatorial optimization*, University of Waterloo Press, R. Kannan and W. Pullyblanks eds, 1990.
 - [Spe06] D. E. Speyer, *A matroid invariant via the K-theory of the Grassmannian*, eprint arXiv:math/0603551v1, 2006.
 - [Sta96] R. P. Stanley, *Combinatorics and commutative algebra: Second edition*, 2nd ed., Birkhäuser, Boston, 1996.
 - [Sta97] R. P. Stanley, *Enumerative combinatorics*, vol. 1, Cambridge University Press, 1997.
 - [Stu96] B. Sturmfels, *Gröbner bases and convex polytopes*, University Lecture Series, vol. 8, American Mathematical Society, 1996.
 - [Top84] D. M. Topkis, *Adjacency on polymatroids*, Mathematical Programming **30** (1984), no. 2, 229–237.
 - [VW08] S. Verdoolaege and K. M. Woods, *Counting with rational generating functions*, J. Symb. Comput. **43** (2008), no. 2, 75–91.
 - [War85] A. Warburton, *Worse case analysis of greedy and related heuristics for some min-max combinatorial optimization problems*, Mathematical Programming **33** (1985), no. 2, 234–241.
 - [Wel76] D. Welsh, *Matroid theory*, Academic Press, Inc., 1976.
 - [Whi80] N. White, *A unique exchange property for bases*, Linear Algebra and its Applications **31** (1980), 81–91.
 - [Woo04] K. Woods, *Rational generating functions and lattice point sets*, Ph.D. thesis, University of Michigan, 2004.